

US00RE44962E

(19) **United States**
(12) **Reissued Patent**
Leemans et al.

(10) **Patent Number:** **US RE44,962 E**
(45) **Date of Reissued Patent:** **Jun. 24, 2014**

(54) **GENETICALLY ENGINEERED PLANT CELLS AND PLANTS EXHIBITING RESISTANCE TO GLUTAMINE SYNTHETASE INHIBITORS, DNA FRAGMENTS AND RECOMBINANTS FOR USE IN THE PRODUCTION OF SAID CELLS AND PLANTS**

(71) Applicants: **Bayer Cropscience N.V.**, Diegem (BE); **Biogen Idec MA, Inc.**, Cambridge, MA (US)

(72) Inventors: **Jan Leemans**, Heusden (BE); **Johan Botterman**, Zwijnaarde (BE); **Marc De Block**, Gentbrugge (BE); **Charles Thompson**, Grand Lancy/Geneve (CH); **Rao Mouva**, Geneva (CH)

(73) Assignees: **Bayer Cropscience N.V.**, Diegem (BE); **Biogen Idec MA Inc.**, Cambridge, MA (US)

(21) Appl. No.: **14/017,108**

(22) Filed: **Sep. 3, 2013**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **7,112,665**
Issued: **Sep. 26, 2006**
Appl. No.: **08/465,219**
Filed: **Jun. 5, 1995**

U.S. Applications:

(60) Division of application No. 07/525,300, filed on May 17, 1990, now Pat. No. 5,561,236, which is a continuation of application No. 07/131,140, filed on Nov. 5, 1987, now abandoned, which is a continuation of application No. PCT/EP87/00141, filed on Mar. 11, 1987.

(30) **Foreign Application Priority Data**

Mar. 11, 1986 (GB) 86400521.0
Jan. 21, 1987 (GB) 87400141.5

(51) **Int. Cl.**
C12N 15/52 (2006.01)
C12N 15/31 (2006.01)

(52) **U.S. Cl.**
USPC **536/23.2**; 536/23.7

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,407,956 A 10/1983 Howell
4,459,355 A 7/1984 Cello et al.
4,769,061 A 9/1988 Comai
5,273,894 A 12/1993 Strauch et al.
5,276,268 A 1/1994 Strauch et al.
5,637,489 A 6/1997 Strauch et al.
5,648,477 A 7/1997 Leemans et al.
5,879,903 A 3/1999 Strauch et al.

FOREIGN PATENT DOCUMENTS

EP 0 173 327 3/1986
EP 0 173 327 A2 * 3/1986
EP 196375 10/1986
GB 2007976 10/1978
GB 2007976 5/1979
WO WO 84/02913 8/1984
WO WO 84/02920 8/1984
WO WO 86/02097 4/1986

OTHER PUBLICATIONS

“Report On The Filing Or Determination Of An Action Regarding A Patent Or Trademark”, U.S. District Court in the Eastern District of Virginia; Case No. 2:12cv46, dated Jan. 25, 2012, 2 pages.

Bayer et al., *Helvetica Chimica Acta*, vol. 55, No. 25, pp. 224-239 (1972).

Bowie et al 1990, *Science* 247:1306-1310.

Browun et al 1998, *Science* 282:1315-1317.

Cashmore, *Proc. Natl. Acad. Sci. USA*, vol. 81, pp. 2960-2964 (1984).

Comai et al 1985, *Nature* 317:741-744.

Coruzzi et al. “Nucleotide Sequences of Two Pea cDNA Clones Encoding the Small Subunit of Riboluse 1,5-Bisphosphate Carboxylase . . .”, *Journal of Biological Chem.* vol. 258, No. 3, Feb. 10, 1983, pp. 1399-1402.

Cullimore et al., “Purification and Properties of two forms of glutamine synthetase from the plant fraction of Phaseolus root nodules”, *Planta Journal*, 1983, 157: 245-253.

De Block et al., “Engineering herbicide resistance in plants by expression of a detoxifying enzyme”, *EMBO Journal* vol. 6, No. 9, 1987, pp. 2513-2518.

De Cleene, M., “The Susceptibility of Monocotyledons to *Agrobacterium tumefaciens*”, *Pythopathology*, 1985, 113: 81-89.

Donn et al., “Herbicide-Resistant Alfalfa Cells: An Example of Gene Amplification in Plants”, *Journal of Molecular and Applied Genetics*, 1984, vol. 2, No. 6, pp. 621-635.

Donn et al., *Chem Abstracts*, 1985, vol. 102, No. 144025J: 140.

E. Strauch et al., “Cloning of a Phosphinothricin N-acetyltransferase Gene from *Streptomyces viridochromogenes* Tu494 and its Expression in *Streptomyces lividans* and *Escherichia coli*”, *Gene*, 63:65-74 (1988), Elsevier Science, Oxford, United Kingdom.

Erickson et al., *Chemical Abstracts*, vol. 104, No. 9, p. 311, 64619g (Mar. 3, 1986).

Fayerman et al., *Biotechnology*, vol. 4, No. 9, pp. 786-789 (Sep. 1986).

Goodman et al., “Gene Transfer in Crop Improvement”, *Science*, Apr. 3, 1987, vol. 236, No. 4797, pp. 48-54.

(Continued)

Primary Examiner — David H Kruse

(74) *Attorney, Agent, or Firm* — Birch, Stewart, Kolasch & Birch, LLP

(57) **ABSTRACT**

The invention relates to a DNA fragment containing a determined gene, the expression of which inhibits the antibiotic and herbicidal effects of Bialaphos and related products. It also relates to recombinant vectors, containing such DNA fragment, which enable this protective gene to be introduced and expressed into cells and plant cells.

2 Claims, 12 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

Hill et al 1998, Biochem. Biophys. Res. Comm. 244:573-577.

Jones et al., Chemical Abstracts, vol. 104, No. 5, p. 152, 29747a (Feb. 1986).

Journal of Cellular Biochemistry, Abstracts, Mar. 31-Apr. 29, 1984, Alan R. Liss, Inc., New York.

Kobayashi et al., The Journal of Antibiotics, vol. 49, No. 5, pp. 688-693 (May 1986).

Langelueddeke et al. Chemical Abstracts, vol. 98, p. 242, 48585v (1982).

Larkin et al., "Somaclonal Variation—a Novel Source of Variability from Cell Cultures for Plant Improvement", Journal for Molecular Appl Genet. vol. 60, 1981, pp. 197-214.

Lazar et al., "Transforming Growth Factor: Mutation of Aspartic Acid 47 and Leucine 48 Results in Different Biological Activities", Molecular and Cellular Biology, vol. 8, Mar. 1988, pp. 1247-1252.

Leason et al., "Inhibition of Pea Leaf Glutamine Synthetase by Methionine Sulphoximine, Phosphinothricin and Other Glutamate Analogues", Phytochemistry, vol. 21, No. 4, pp. 855-857, 1982.

Mason et al., Phytochemistry, vol. 21, No. 4, pp. 855-857 (1982).

Murakami et al., "The bialaphos biosynthetic genes of *Streptomyces hygroscopicus*: Molecular cloning and characterization of the gene cluster", Mol Gen Genet. vol. 205, 1986, pp. 42-50.

Murakami et al., Chemical Abstracts, vol. 106, p. 1149, 1151u (1987).

Sanders et al., "Amplification and cloning of the Chinese hamster glutamine synthetase gene", The EMBO Journal, vol. 3, No. 1, pp. 65-71, (1984).

Schreier et al. EMBO J., vol. 4, pp. 25-32, (1985).

Scolnik et al., "The Wild-Type Gene for Glutamine Synthetase Restores Ammonia Control of the Nitrogen Fixation . . .", Journal of Bacteriology, vol. 155, No. 1, Jul. 1983, pp. 180-185.

Shields, "Engineering Herbicide Resistance", Nature, vol. 317, Oct. 1985, p. 668.

Stalker et al., "Herbicide Resistance in Transgenic Plants Expressing a Bacterial Detoxification Gene", Science, Oct. 21, 1988, pp. 419-423.

T. W. Goodwin and E. I. Mercer, "Introduction to Plant Biochemistry", 1983, 2nd ed., pp. 349-353.

Thompson (Sep. 3-8, 1984) Genetic Engineering and Antibiotic Production, Rev. Biol., vol. 77, No. 4, pp. 567-568.

Thompson et al 1987, The EMBO Journal 6(9):2519-2523.

Thompson et al., Journal of Bacteriology, vol. 151, No. 2, pp. 678-685 (1982).

Thompson et al., Mol. Gen. Genet., vol. 195, pp. 39-43 (1984).

Thompson et al., Nature, vol. 286, No. 5772, pp. 525-527 (1980).

Thompson et al., Proc. Natl. Acad. Sci. USA, vol. 80, pp. 5190-5194 (1983).

Thompson, et al., Gene, vol. 20, pp. 51-62 (1982).

Vara et al., "Cloning and expression of a puromycin N-acetyl transferase gene from *Streptomyces alboniger* in *Streptomyces lividans* and *Escherichia coli*", Gene, vol. 33, No. 2, 1985, pp. 197-206.

Vasil, "Progress In The Regeneration and Genetic Manipulation of Cereal Crops", Biotechnology vol. 6, Apr. 1988, pp. 397-402.

Velten et al. Nucleic Acids Research, vol. 13, No. 19, pp. 6981-6998 (1985).

W. Wohlleben et al., "Nucleotide Sequence of the Phosphinothricin N-acetyltransferase Gene from *Streptomyces viridochromogenes* Tu 494 and its Expression in *Nicotiana tabacum*", Gene, 70:25-37 (1988), Elsevier Science, Oxford, United Kingdom.

Wedler et al., "Interaction of a new γ -glutamyl-phosphate analog, 4-(phosphonoacetyl) . . .", Archives of Biochemistry and Biophysics, vol. 202, No. 2, Jul. 1980, pp. 482-490.

Winnacker, Ernst-L., "From Genes to Clones: Introduction to Gene Technology", 1987, pp. 413-415.

Worthing (Ed.), The Pesticide Manual, 7th Ed., p. 302, 1983.

Zalacain et al. Nucleic Acids Research, vol. 14, No. 4, pp. 1565-1581 (Feb. 1986).

* cited by examiner

FIG. 1





FIG. 2







FIG. 3






FIG.4B





FIG. 5A




FIG. 5B




FIG. 5C






FIG. 6A








Fig. 8.

TAAAGAGGTGCCCCGCCACCCGCTTTCCGAGAACACC GAAGGAGACCACAC
↓
GTGAGCCCAGAACGACGCCCGGTCGAGATCCGTCCCGCCACCGCCGCCGA
CATGGCGGCGGTCTGCGACATCGTCAATCACTACATCGAGACGAGCACGG
TCAACTTCCGTACGGAGCCGCAGACTCCGCAGGAGTGGATCGACGACCTG
GAGCGCCTCCAGGACCGCTACCCCTGGCTCGTCGCCGAGGTGGAGGGCGT
CGTCGCCGGCATCGCCTACGCCGGCCCCCTGGAAGGCCCGCAACGCCCTACG
ACTGGACCGTCGAGTCGACGGTGACGTCTCCACCGGCCACCAGCGGCTC
GGACTGGGCTCCACCCTCTACACCCACCTGCTGAAGTCCATGGAGGCCCA
GGGCTTCAAGAGCGTGGTCGCCGTCATCGGACTGCCCAAACGACCCGAGCG
TGCGCCTGCACGAGGCGCTCGGATACACCGCGCGCGGGACGCTGCGGGCA
GCCGGCTACAAGCACGGGGGCTGGCACGACGTGGGGTTCTGGCAGCGCGA
CTTCGAGCTGCCGGCCCCCGCCCCCGCCCCGTCCGGCCCGTCACACAGATCT
↑
GAGCGGAGAGCGCATGGC

FIG. 9

	10	20	30	40	50	
PBG 39	-----	MSPERRPADI	RRATEADMPA	VCTIVNHYIE	TSTVNFRTPE	QEPQEWTDL
PJS1	-----	VSPERRPVEI	RPATAADMAA	VCDIVNHYIE	TSTVNFRTPE	QTPQEWTDL
	60	70	80	90	100	
PBG 39	-----	VRLRERYPWL	VAEVDGEVAG	IAYAGPWKAR	NAYDWTAE	TSTVNFRTPE
PJS1	-----	ERLQDRYPWL	VAEVEGVVAG	IAYAGPWKAR	NAYDWTAE	TSTVNFRTPE
	110	120	130	140	150	
PBG 39	-----	GLGSTLYTHL	LKSLEAQGFK	SVVAVIGLPN	DPSVRMHEAL	GYAPRGMLRA
PJS1	-----	GLGSTLYTHL	LKSMEAQGFK	SVVAVIGLPN	DPSVRMHEAL	GYTARGTLRA
	160	170	180			
PBG 39	-----	AGFKHGNWHD	VGFWQLDFSL	PVPPRPVLPV	TEI★	
PJS1	-----	AGYKHGGWHD	VGFWQRDFEL	PAPPRPVRPV	TQI★	

FIG.10




FIG. 11

1

**GENETICALLY ENGINEERED PLANT
CELLS AND PLANTS EXHIBITING
RESISTANCE TO GLUTAMINE SYNTHETASE
INHIBITORS, DNA FRAGMENTS AND
RECOMBINANTS FOR USE IN THE
PRODUCTION OF SAID CELLS AND PLANTS**

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in *italics* indicates the additions made by reissue.

This application is a divisional of application Ser. No. 07/525,300, filed May 17, 1990, now U.S. Pat. No. 5,561,236, which is a continuation of application Ser. No. 07/131,140, filed Nov. 5, 1987, abandoned, which is a continuation under 35 USC §§363 and 120 of PCT/EP 87/00141, filed Mar. 11, 1987, published as WO 87/05629.

The invention relates to a process for protecting plant cells and plants against the action of glutamine synthetase inhibitors.

It also relates to applications of such process, particularly to the development of herbicide resistance into determined plants.

It relates further to non-biologically transformed plant cells and plants displaying resistance to glutamine synthetase inhibitors as well as to suitable DNA fragments and recombinants containing nucleotide sequences encoding resistance to glutamine synthetase inhibitors.


Glutamine synthetase (hereafter simply designated by GS) constitutes in most plants one of the essential enzymes for the development and life of plant cells. It is known that GS converts glutamate into glutamine. GS is involved in an efficient pathway (the only one known nowadays) in most plants for the detoxification of ammonia released by nitrate reduction, aminoacid degradation or photorespiration. Therefore potent inhibitors of GS are very toxic to plant cells. A particular class of herbicides has been developed, based on the toxic effect due to inhibit inhibition of GS in plants.

These herbicides comprise as active ingredient a GS inhibitor.

There are at least two possible ways which might lead to plants resistant to the inhibitors of the action of glutamine synthetase; (1) by changing the target. It can be envisaged that mutations in the GS enzyme can lead to insensitivity towards the herbicide; (2) by inactivation of the herbicide. Breakdown or modification of the herbicide inside the plant could lead to resistance.


Bialaphos and phosphinothricin (hereafter simply designated by PPT) are two such inhibitors of the action of GS, (ref. 16, 17) and have been shown to possess excellent herbicidal properties (see more particularly ref. 2 as concerns Bialaphos).

Bialaphos has the following formula (I)



2

PPT has the following formula (II)



Thus the structural difference between PPT and Bialaphos resides in the absence of two alanine aminoacids in the case of PPT.

These two herbicides are non selective. They inhibit growth of all the different species of plants present on the soil, accordingly cause their total destruction.

Bialaphos was first disclosed as having antibiotic properties, which enabled it to be used as a pesticide or a fungicide. Bialaphos can be produced according to the process disclosed in U.S. Pat. No. 3,832,394, assigned to MEIJI SEIKA KAI-SHA LTD., which patent is incorporated herein by reference. It comprises cultivating *Streptomyces hygroscopicus*, such as the strain available at the American Type Culture Collection, under the ATCC number 21,705, and recovering Bialaphos from its culture medium. However, other strains, such as *Streptomyces viridochromogenes*, also produce this compound (ref. 1).

Other tripeptide antibiotics which contain a PPT moiety are or might be discovered in nature as well, e.g. phosalacin (ref. 15).

PPT is also obtained by chemical synthesis and is commercially distributed by the industrial Company HOECHST.

A number of *Streptomyces* species have been disclosed which produce highly active antibiotics which are known to incapacitate procaryotic cell functions or enzymes. The *Streptomyces* species which produce these antibiotics would themselves be destroyed if they had not a self defence mechanism against these antibiotics. This self defence mechanism has been found in several instances to comprise an enzyme capable of inhibiting the antibiotic effect, thus of avoiding autotoxicity for the *Streptomyces* species concerned. This modification is generally reversed when the molecule is exported from the cell.

The existence of a gene which encodes an enzyme able to modify the antibiotic so as to inhibit the antibiotic effect against the host has been demonstrated in several *Streptomyces* producing antibiotics, for example, in *S. fradiae*, *S. azureus*, *S. vinaceus*, *S. erythreus*, producing neomycin, thiostrepton, viomycin, and MLS (Macrolide Lincosamide Streptogramin) antibiotics respectively (ref. 4), (ref. 5), (ref. 6), (ref. 14 by CHATER et al., 1982 describes standard techniques which can be used for bringing these effects to light).

In accordance with the present invention, it has been found that *Streptomyces hygroscopicus* ATCC 21,705, also possesses a gene encoding an enzyme responsible of the inactivation of the antibiotic properties of Bialaphos. Experiments carried out by the applicants have lead to the isolation of such a gene and its use in a process for controlling the action of GS inhibitors, based on PPT or derived products.

An object of the invention is to provide a new process for controlling the action in plant cells and plants of GS inhibitors.

Another object of the invention is to provide DNA fragments and DNA recombinants, particularly modified vectors containing said DNA fragments, which DNA fragments contain nucleotide sequences capable, when incorporated in plant cells and plants, to protect them against the action of GS inhibitors.

3

A further object of the invention is to provide non-biologically transformed plant cells and plants capable of neutralizing or inactivating GS inhibitors.

A further object of the invention is to provide a process for selectively protecting plant species against herbicides of a GS inhibitor type.

More specifically an object of the invention is to provide a DNA fragment transferable to plant cells- and to whole plants—capable of protecting them against the herbicidal effects of Bialaphos and of structurally analogous herbicides.

A further object of the invention is to provide plant cells resistant to the products of the class exemplified by Bialaphos, which products possess the PPT unit in their structure.

The process according to the invention for controlling the action in plant cells and plants of a GS inhibitor when contacted therewith, comprises providing said plants with a heterologous DNA fragment including a foreign nucleotide sequence, capable of being expressed in the form of a protein

in said plant cells and plants, under condition such as to cause said heterologus DNA fragment to be integrated stably through generations in the cells of said plants, and wherein said protein has an enzymatic activity capable of inactivating or neutralization of said glutamine synthetase inhibitor.

A preferred DNA fragment is one derived from an antibiotic-producing-Streptomyces strain (or a sequence compris-

4

ing a nucleotide sequence encoding the same activity) and which encodes resistance to a said GS inhibitors.

Preferred nucleotide sequences for use in this invention encode a protein which has acetyl tranferase activity with respect to said GS inhibitors.

A most preferred DNA fragment according to the invention comprises a nucleotide sequence coding for a polypeptide having a PPT acetyl transferase activity.

A particular DNA fragment according to the invention, for the subsequent transformation of plant cells, consists of a nucleotide sequence coding for at least part of a polypeptide having the following sequence (SEQ ID NO:1):

183 X SER PRO GLU

ARG ARG PRO ALA ASP ILE ARG ARG ALA THR GLU ALA ASP MET PRO 228

ALA VAL CYS THR ILE VAL ASN HIS TYR ILE GLU THR SER THR VAL 273

ASN PHE ARG THR GLU PRO GLN GLU PRO GLN GLU TRP THR ASP ASP 318

LEU VAL ARG LEU ARG GLU ARG TYR PRO TRP LEU VAL ALA GLU VAL 363

ASP GLY GLU VAL ALA GLY ILE ALA TYR ALA GLY PRO TRP LYS ALA 408

ARG ASN ALA TYR ASP TRP THR ALA GLU SER THR VAL TYR VAL SER 453

PRO ARG HIS GLN ARG THR GLY LEU GLY SER THR LEU TYR THR HIS 498

LEU LEU LYS SER LEU GLU ALA GLN GLY PHE LYS SER VAL VAL ALA 543

VAL ILE GLY LEU PRO ASN ASP PRO SER VAL ARG MET HIS GLU ALA 588

LEU GLY TYR ALA PRO ARG GLY MET LEU ARG ALA ALA GLY PHE LYS 633

HIS GLY ASN TRP HIS ASP VAL GLY PHE TRP GLN LEU ASP PHE SER 678

LEU PRO VAL PRO PRO ARG PRO VAL LEU PRO VAL THR GLU ILE 723

183 GTG AGC CCA GAA

CGA CGC CCG GCC GAC ATC CGC CGT GCC ACC GAG GCG GAC ATG CCG 228

in which X represents MET or VAL, which part of said polypeptide is of sufficient length to confer protection against Bialaphos to plant cells, when incorporated genetically and expressed therein, i.e. as termed hereafter “plant-protecting capability” against Bialaphos.

A preferred DNA fragment consists of the following nucleotide sequence (SEQ ID NO:2):

-continued

GCG GTC TGC ACC ATC GTC AAC CAC TAC ATC GAG ACA AGC ACG GTC
273

AAC TTC CGT ACC GAG CCG CAG GAA CCG CAG GAG TGG ACG GAC GAC
318

CTC GTC CGT CTG CGG GAG CGC TAT CCC TGG CTC GTC GCC GAG GTG
363

GAC GGC GAG GTC GCC GGC ATC GCC TAC GCG GGC CCC TGG AAG GCA
408

CGC AAC GCC TAC GAC TGG ACG GCC GAG TCG ACC GTG TAC GTC TCC
453

CCC CGC CAC CAG CGG ACG GGA CTG GGC TCC ACG CTC TAC ACC CAC
498

CTG CTG AAG TCC CTG GAG GCA CAG GGC TTC AAG AGC GTG GTC GCT
543

GTC ATC GGG CTG CCC AAC GAC CCG AGC GTG CGC ATG CAC GAG GCG
588

CTC GGA TAT GCC CCC CGC GGC ATG CTG CGG GCG GCC GGC TTC AAG
633

CAC GGG AAC TGG CAT GAC GTG GGT TTC TGG CAG CTG GAC TTC AGC
678

CTG CCG GTA CCG CCC CGT CCG GTC CTG CCC GTC ACC GAG ATC
723

or of a part thereof expressing a polypeptide having plant-
protecting capability against Bialaphos.



The invention also relates to any DNA fragment differing
from the preferred one indicated hereabove by the replace-
ment of any of its nucleotides by others, yet without modify-
ing the genetic information of the preferred DNA sequence
mentioned hereabove (normally within the meaning of the
universal genetic code), and furthermore to any equivalent
DNA sequence which would encode a polypeptide having the
same properties, particularly a Bialaphos-resistance-activity.

It will be understood that the man skilled in the art should
be capable of readily assessing those parts of the nucleotide
sequences that could be removed from either side of any of the
DNA fragments according to the invention, for instance by
removing terminal parts on either side of said DNA fragment,
such as by an exonucleolytic enzyme, for instance Bal31, by
recloning the remaining fragment in a suitable plasmid and by
assaying the capacity of the modified plasmid to transform
appropriate cells and to protect it against the Bialaphos anti-
biotic or herbicide as disclosed later, whichever assay is
appropriate.

For the easiness of language, these DNA fragments will be
termed hereafter as "Bialaphos-resistance DNA". In a similar
manner, the corresponding polypeptide will be termed as
"Bialaphos-resistance enzyme".

While in the preceding discussion particular emphasis has
been put on DNA fragments capable, when introduced into
plant cells and plants, to confer on them protection against
Bialaphos or PPT, it should be understood that the invention
should in no way be deemed as limited thereto.

In a same manner, the invention pertains to DNA fragments
which, when introduced into such plant cells, would also
confer on them a protection against other GS inhibitors, for
instance of intermediate products involved in the natural bio-
synthesis of phosphinotricin, such as the compounds design-
ated by the abbreviations MP101 (III), MP102 (IV), the
formula of which are indicated hereafter:



More generally, the invention has opened the route to the
production of DNA fragments which, upon proper incorpora-
tion into plant cells and plants, can protect them against GS
inhibitors when contacted therewith, as this will be shown in
a detailed manner in relation to Bialaphos and PPT in the
examples which will follow.

This having been established, it will be appreciated that any
fragment encoding an enzymatic activity which would pro-
tect plant cells and plants against said GS inhibitors, by inac-
tivating, should be viewed as an equivalent of the preferred
fragments which have been disclosed hereabove. This would
apply especially to any DNA fragments that would result
from genetic screening of the genomic DNAs of strains, par-
ticularly of antibiotic-producing strains, likely to possess
genes which, even-though structurally different, would
encode similar activity with respect to Bialaphos or PPT, or
even with respect to other GS inhibitors. This applies to any
gene in other strains producing a PPT derivative.

Therefore, it should be understood that the language
"Bialaphos-resistance DNA" or "Bialaphos-resistance
enzyme" used thereafter as a matter of convenience is
intended to relate not only to the DNAs and enzymes specifi-
cally concerned with resistance to PPT or most directly
related derivatives, but more generally with other DNAs and
enzymes which would be capable, under the same circum-
stances, of inactivating the action in plants of GS inhibitors.

The invention also relates to DNA recombinants containing the above defined Bialaphos-resistance DNA fragments recombined with heterologous DNA, said heterologous DNA containing regulation elements and said Bialaphos-resistance DNA being under the control of said regulation elements in such manner as to be expressible in a foreign cellular environment compatible with said regulation elements. Particularly the abovesaid Bialaphos-resistance-DNA fragments contained in said DNA recombinants are devoid of any DNA region involved in the biosynthesis of Bialaphos, when said Bialaphos-resistance-DNA fragment originate themselves from Bialaphos-producing strains.

By "heterologous DNA" is meant a DNA of an other origin than that from which said Bialaphos-resistance-DNA originated, e.g. is different from that of a *Streptomyces hygroscopicus* or *Streptomyces viridochromogenes* or even more preferably a DNA foreign to *Streptomyces* DNA. Particularly said regulation elements are those which are capable of controlling the transcription and translation of DNA sequences normally associated with them in said foreign environment. "Cellular" refers both to microorganisms and to cell cultures.

This heterologous DNA may be a bacterial DNA, particularly when it is desired to produce a large amount of the recombinant DNA, such as for amplification purposes. In that respect a preferred heterologous DNA consists of DNA of *E. coli* or of DNA compatible with *E. coli*. It may be DNA of the same origin as that of the cells concerned or other DNA, for instance viral or plasmidic DNA known as capable of replicating in the cells concerned.

Preferred recombinant DNA contains heterologous DNA compatible with plant cells, particularly Ti-plasmid DNA.

Particularly preferred recombinants are those which contain GS inhibitor inactivating DNA under the control of a promoter recognized by plant cells, particularly those plant cells on which inactivation of GS inhibitors is to be conferred.

Preferred recombinants according to the invention further relate to modified vectors, particularly plasmids, containing said GS-inhibitor-inactivating DNA so positioned with respect to regulation elements, including particularly promoter elements, that they enable said GS inhibitor-inactivating DNA to be transcribed and translated in the cellular environment which is compatible with said heterologous DNA. Advantageous vectors are those so engineered as to cause stable incorporation of said GS inhibitor inactivating DNA in foreign cells, particularly in their genomic DNA. Preferred modified vectors are those which enable the stable transformation of plant cells and which confer to the corresponding cells, the capability of inactivating GS inhibitors.

It seems that, as described later, the initiation codon of the Bialaphos-resistance-gene of the *Streptomyces hygroscopicus* strain used herein is a GTG codon. But in preferred recombinant DNAs or vectors, the Bialaphos-resistance-gene is modified by substitution of an ATG initiation codon for the initiation codon GTG, which ATG enables translation initiation in plant cells.

In the example which follows, the plant promoter sequence which has been used was constituted by a promoter of the 35 S cauliflower mosaic virus. Needless to say that the man skilled in the art will be capable of selecting other plant promoters, when more appropriate in relation to the plant species concerned.

According to an other preferred embodiment of the invention, particularly when it is desired to achieve transport of the enzyme encoded by the Bialaphos-resistance-DNA into the chloroplasts, the heterologous DNA fragment is fused to a gene or DNA fragment encoding a transit peptide, said last

mentioned fragment being then intercalated between the GS inhibitor inactivating gene and the plant promoter selected.

As concerns means capable of achieving such constructions, reference can be made to the following British applications 84 32757 filed on Dec. 28, 1984 and 85 00336 filed on Jan. 7, 1985 and to the related applications filed in the United-States of America (No. 06/755,173, filed on Jul. 15, 1985), in the European Patent Office (No. 85 402596.2, filed on Dec. 20, 1985), in Japan (No. 299 730, filed on Dec. 27, 1985), in Israel (No. 77 466 filed on Dec. 27, 1985) and in Australia (No. 5 165 485, filed on Dec. 24, 1985), all of which are incorporated herein by reference.

Reference can also be made to the scientific literature, particularly to the following articles:

VAN DEN BROECK et al., 1985, *Nature*, 313, 358-363;
SCHREIER and al., *Embo. J.*, vol. 4, No. 1, 25-32.

These articles are also incorporated herein by reference.

For the sake of the record, be it recalled here that under the expression "transit peptide", one refers to a polypeptide fragment which is normally associated with a chloroplast protein or a chloroplast protein sub-unit in a precursor protein encoded by plant cell nuclear DNA. The transit peptide then separates from the chloroplast protein or is proteolitically removed, during the translocation process of the latter protein into the chloroplasts. Examples of suitable transit peptides are those associated with the small subunit of ribulose-1,5 biphosphate (RuBP) carboxylase or that associated with the chlorophyll a/b binding proteins.

There is thus provided DNA fragments and DNA recombinants which are suitable for use in the process defined hereafter.

More particularly the invention also relates to a process, which can be generally defined as a process for producing plants and reproduction material of said plants including a heterologous genetic material stably integrated therein and capable of being expressed in said plants or reproduction material in the form of a protein capable of inactivating or neutralizing the activity of a glutamine synthetase-inhibitor, comprising the non biological steps of producing plants cells or plant tissue including said heterologous genetic material from starting plant cells or plant tissue not able to express that inhibiting or neutralizing activity, regenerating plants or reproduction material of said plants or both from said plant cells or plant tissue including said genetic material and, optionally, biologically replicating said last mentioned plants or reproduction material or both, wherein said non-biological steps of producing said plant cells or plant tissue including said heterologous genetic material, comprises transforming said starting plant cells or plant tissue with a DNA-recombinant containing a nucleotide sequence encoding said protein, as well as the regulatory elements selected among those which are capable of enabling the expression of said nucleotide sequence in said plant cells or plant tissue, and to cause the stable integration of said nucleotide sequence in said plant cells and tissue, as well as in the plant and reproduction material processed therefrom throughout generations.

The invention also relates to the cell cultures containing Bialaphos-resistance-DNA, or more generally said GS-inhibitor-inactivating DNA, which cell cultures have the property of being resistant to a composition containing a GS inhibitor, when cultured in a medium containing a such composition at dosages which would be destructive for non transformed cells.

The invention concerns more particularly those plant cells or cell cultures in which the Bialaphos-resistance DNA is stably integrated and which remains present over successive generations of said plant cells. Thus the resistance to a GS

inhibitor, more particularly Bialaphos or PPT, can also be considered as a way of characterizing the plant cells of this invention.

Optionally one may also resort to hybridization experiments between the genomic DNA obtained from said plant cells with a probe containing a GS inhibitor inactivating DNA sequence.

More generally the invention relates to plant cells, reproduction material, particularly seeds, as well as plants containing a foreign or heterologous DNA fragment stably integrated in their respective genomic DNAs, said fragments being transferred throughout generations of such plant cells, reproduction material, seeds and plants, wherein said DNA fragment encodes a protein inducing a non-variety-specific enzymatic activity capable of inactivating or neutralizing GS inhibitors, particularly Bialaphos and PPT, more particularly to confer on said plant cells, reproduction material, seeds and plants a corresponding non-variety-specific phenotype of resistance to GS inhibitors.

"Non-variety-specific" enzymatic activity or phenotype aims at referring to the fact that they are not characteristic of specific plant genes or species as this will be illustrated in a non-limitative way by the examples which will follow. They are induced in said plant materials by essentially non-biological processes applicable to plants belonging to species normally unrelated with one another and comprising the incorporation into said plant material of heterologous DNA, e.g. bacterial DNA or chemically synthesized DNA, which does not normally occur in said plant material or which normally cannot be incorporated therein by natural breeding processes, and which yet confers a common phenotype (e.g. herbicide resistance) to them.

The invention is of particular advantageous use in processes for protecting field-cultivated plant species against weeds, which processes comprise the step of treating the field with an herbicide, e.g. Bialaphos or PPT in a dosage effective to kill said weeds, wherein the cultivated plant species then contains in their genome a DNA fragment encoding a protein having an enzymatic activity capable of neutralizing or inactivating said GS inhibitor.

By way of illustration only, effective doses for use in the abovesaid process range from about 0.4 to about 1.6 kg/Hectare of Bialaphos or PPT.

There follows now a disclosure of how the preferred DNA fragment described hereabove was isolated starting from the *Streptomyces hygroscopicus* strain available at the American Type Culture Collection under deposition number ATCC 21705, by way of exemplification only.

The following disclosure also provides the technique which can be applied to other strains producing compounds with a PPT moiety.

The disclosure will then be completed with the description of the insertion of a preferred DNA fragment conferring to the transformed cells the capability of inactivating Bialaphos and PPT. Thus the Bialaphos-inactivating-DNA fragment designated thereafter by Bialaphos-resistance gene or "sfr" gene, isolated by the above described technique into plasmids which can be used for transforming plant cells and conferring to them a resistance against Bialaphos, also merely by way of example for non-limitative illustration purposes.

The following disclosure is made with reference to the drawings in which:

FIG. 1 is a restriction map of a plasmid containing a *Streptomyces hygroscopicus* DNA fragment encoding Bialaphos-resistance, which plasmid, designated hereafter as pBG1 has been constructed according to the disclosure which follows;

FIG. 2 shows the nucleotide sequence (SEQ ID NO:12) of a smaller fragment obtained from pBG1, subcloned into another plasmid (pBG39) and containing the resistance gene;

FIG. 3 shows the construction of a series of plasmids given by way of example, which plasmids aim at providing suitable adaptation means for the insertion therein of the Bialaphos-resistance gene or "sfr" gene;

FIGS. 4A and 4B show the construction of a series of plasmids given by way of example, which plasmids contain suitable plant cell promoter sequences able to initiate transcription and expression of the foreign gene inserted under their control into said plasmids;

FIG. 5A shows a determined fragment of the nucleotide sequence (SEQ ID NO:13) of the plasmid obtained in FIG. 3;

FIG. 5B shows the reconstruction of the first codons of a Bialaphos-resistance gene, from a FokI/BglII fragment obtained from pBG39 and the substitution of an ATG initiation codon for the GTG initiation codon of the natural "sfr" gene (SEQ ID NO:14);

FIG. 5C shows the reconstruction of the entire "sfr" gene, namely the last codons thereof (SEQ ID NO:15), and its insertion into a plasmid obtained in FIGS. 4A and 4B;

FIG. 6A shows an expression vector containing the "sfr" gene placed under the control of a plant cell promoter;

FIG. 6B shows another expression vector deriving from the one shown in FIG. 6A, by the substitution of some nucleotides.

FIG. 7 shows the construction of a series of plasmids given by way of examples, to ultimately produce plasmids containing the promoter region and the transit peptide sequence of a determined plant cell gene, for the insertion of the "sfr" gene under the control of said promoter region and downstream of said transit peptide sequence (SEQ ID NOS: 16 and 17).

FIGS. 8 to 11 will be referred to hereafter.

The following experiment was set up to isolate a Bialaphos-resistance-gene from *S. hygroscopicus*, according to standard techniques for cloning into *Streptomyces*.

2.5 μ g of *S. hygroscopicus* genomic DNA and 0.5 μ g of *Streptomyces* vector pIJ61 were cleaved with PstI according to the method described in ref. 6. The vector, fragments and genomic fragments were mixed and ligated (4 hours at 10° C. followed by 72 hours at 4° C. in ligation salts which contain 66 mM Tris-HCl (pH 7.5), 1 mM EDTA, 10 mM MgCl₂, 10 mM 2-mercaptoethanol and 0.1 mM ATP) at a total DNA concentration of 40 μ g ml⁻¹ with T4 DNA ligase. Ligation products were introduced into 3 \times 10⁹ *S. lividans* strain 66 protoplasts by a transformation procedure mediated by polyethylene-glycol (PEG) as described hereafter. These protoplasts gave rise to 5 \times 10⁷ colonies and 4 \times 10⁴ pocks after regeneration on 20 plates of R2 agar containing 0.5% of Difco yeast extract (R2 YE). Preparation and composition of the different mediums and buffers used in the disclosed experiments are described hereinafter. When these lawns were replica-plated on minimal medium plates containing 50 μ g ml⁻¹ Bialaphos, drug resistant colonies appeared at a frequency of 1 per 10⁴ transformants. After purification of the drug resistant colonies, there plasmid DNA was isolated and used to retransform *S. lividans* protoplasts. Non selective regeneration followed by replication to Bialaphos-containing-medium demonstrated a 100% correlation between pocks and Bialaphos resistant growth. The recombinant plasmids of several resistant clones all contained a 1.7 Kb PstI insert (see FIG. 1).

Subcloning of the Herbicide Resistance Gene

The 1.7 Kb PstI insert was then subcloned into the high copy number streptomycete vector pIJ385 to generate plasmid pBG20. *S. lividans* strains which contained pBG20 were

more than 500 times more resistant to Bialaphos. *S. lividans* growth is normally inhibited in minimal medium containing 1 µg/ml Bialaphos; growth of transformants containing pBG20 was not noticeably inhibited in a medium containing 500 µg/ml Bialaphos. The PstI fragment was also subcloned in either orientation into the PstI site of the plasmid pBR322, to produce plasmids pBG1 and pBG2, according to their orientation. A test on minimal M9 medium demonstrated that *E. coli* E8767 containing pBG1 or pBG2 was resistant to Bialaphos.

A ±1.65 Kb PstI-BamHI fragment was subcloned from pBG1 into the plasmid pUC19 to produce the plasmid pBG39, and conferred Bialaphos resistance to *E. coli* W3110, C600 and JM83.

Using an in vitro coupled transcription-translation system (ref. 5) from *S. lividans* extracts, the 1.65 Kb PstI-BamHI fragment in pBG39 was shown to direct the synthesis of a 22 Kd protein. In the following, this 1.65 Kb insert includes a fragment coding for a 22 Kd protein and will be called "sfr" gene.

Fine Mapping and Sequencing of the Gene

A 625 bp Sau3A fragment was subcloned from pBG39 into pUC19 and still conferred Bialaphos resistance to a *E. coli* W3110 host. The resulting clones were pBG93 and pBG94, according to the orientation.

The orientation of the gene in the Sau3A fragment was indicated by experiments which have shown that Bialaphos resistance could be induced with IPTG from the pUC19 lac promoter in pBG93. In the presence of IPTG (0.5 mM) the resistance of pBG93/W3110 increased from 5 to 50 µg/ml on a M9 medium containing Bialaphos. The W3110 host devoid of pBG93, did not grow on M9 medium containing 5 µg/ml Bialaphos. These experiments demonstrated that the Sau3A fragment could be subcloned without loss of activity. They also provided for the proper orientation as shown in the FIG. 2, enclosed thereafter. The protein encoded by these clones was detected by using coupled transcription-translation systems derived from extracts of *S. lividans* (ref. 7). Depending on the orientation of the Sau3A fragment, translation products of different sizes were observed; 22 Kd for pBG94 and ±28 Kd for pBG93. This indicated that the Sau3A fragment did not contain the entire resistance gene and that a fusion protein was formed which included a polypeptide sequence resulting from the translation of a pUC19 sequence.

In order to obtain large amounts of the protein, a 1.7 Kb PstI fragment from pBG1 was cloned into the high copy number *Streptomyces* replicon pIJ385. The obtained plasmid, pBG20, was used to transform *S. hygroscopicus*. Transformants which contained this plasmid had more than 5 times as much PPT acetylating activity and also had increased amounts of a 22 kd protein on sodium dodecylsulfate gels (SDS gels). Furthermore, both the acetyl transferase and the 22 kd protein appeared when the production of Bialaphos began. The correlation of the in vitro data, kinetics of synthesis, and amplified expression associated with pBG20 transformants strongly implied that this 22 Kd band was the gene product.

The complete nucleotide sequence of the 625 bp Sau3A fragment was determined as well as of flanking sequences. Computer analysis revealed the presence of an open reading frame over the entire length of the Sau3A fragment.

Characterization of the sfr Gene Product

A series of experiments were performed to determine that the open reading frame of the "sfr" gene indeed encoded the Bialaphos resistance enzyme. To determine the 5' end of the resistance gene, the NH₂-terminal sequence of the enzyme was determined. As concerns more particularly the technique

used to determine the said sequence, reference is made to the technique developed by J. VANDEKERCKHOVE, Eur. J. Bioc. 152, p. 9-19, 1985, and to French patent applications No. 85 14579 filed on Oct. 1, 1985 and No. 85 13046 filed on September 2, 1985, all of which are incorporated herein by reference.

This technique allows the immobilization on glass fibre sheets coated with the polyquaternary amine commercially available under the registered trademark POLYBRENE of proteins and of nucleic acids previously separated on a sodium dodecylsulfate containing polyacrylamide gel. The transfer is carried out essentially as for the protein blotting on nitrocellulose membranes (ref. 8). This allows the determination of amino-acid composition and partial sequence of the immobilized proteins. The portion of the sheet carrying the immobilized 22 kd protein produced by *S. hygroscopicus* pBG20 was cut out and the disc was mounted in the reaction chamber of a gas-phase sequenator to subject the glass-fibre bound 22 Kd protein to the Edman degradation procedure. The following amino-acid sequence was obtained SEQ ID NO:4):

Pro-Glu-Arg-Arg-Pro-Ala-Asp-Ile-Arg-Arg

This sequence matched an amino-acid sequence which was deduced from the open reading frame of the 625 bp Sau3A fragment. It corresponded to the stretch from codon 3 to codon 12.

Thus, the NH₂-terminus of the 22 Kd protein was upstream of this sequence. It was determined that translation of the actual protein was likely to be initiated 2 amino-acids earlier at a GTG initiation codon. GTG is often used as initiator codon in *Streptomyces* and translated as methionine. The protein translated from the GTG initiation codon would be 183 amino-acids long and would have a molecular weight of 20 550. This was in good agreement with the observed approximate molecular weight of 22 000.

Furthermore, the termination codon, TGA, was located just downstream of the Sau3A site. Cloning of the 625 bp Sau3A fragment in a BamHI site digested pUC19 did not result in the reconstruction of the termination codon. This explained the fusion proteins which were observed in the in vitro transcription-translation analysis.

Mechanism of PPT-Resistance

Having defined a first phenotype and some of the physical characteristics of the resistance gene and its gene product, a series of experiments was then carried out to understand the mechanism by which it confers resistance. As described hereabove, PPT is the portion of Bialaphos which inhibits glutamine synthetase (GS) and that N-acetyl PPT is not an inhibitor. Using a standard assay (ref. 9), *S. hygroscopicus* ATCC 21 705 derivatives were shown to contain a PPT acetyl transferase which was not found in *S. lividans*. The activity does not acetylate the Bialaphos tripeptide. *S. lividans* carrying the resistance gene cloned in pBG20 or pBG16 (a plasmid containing the 625 bp Sau3A fragment cloned into another streptomyces vector, pIJ680) also contained the activity which could acetylate PPT but not Bialaphos. The PPT derived reaction product produced by extracts of pBG20/*S. lividans* was isolated in order to confirm that it was indeed acetyl-PPT. Analysis by mass spectroscopy showed that the molecular weight had increased relative to PPT by the equivalent of one acetyl group. It was thus concluded that the 625 bp Sau3A fragment contained sequences which code for PPT acetyl transferase.

13

The experimental conditions and reagents used in the techniques disclosed hereabove were as follows:

Preparation and Composition of the Mediums and Buffers Above Used

1° P medium: 10.3 g of sucrose, 0.025 g of K_2SO_4 , 0.203 g of $MgCl_2 \cdot 6H_2O$ and 0.2 ml of a trace element solution are dissolved in 80 ml of distilled water and autoclaved. Then in order, 1 ml of KH_2PO_4 (0.5%), 10 ml of $CaCl_2 \cdot 2H_2O$ (3.68%), and 10 ml of TES buffer (0.25 M), pH: 7.2) are added. Trace element solution (per liter): $ZnCl_2$, 40 mg; $FeCl_3 \cdot 6H_2O$, 200 mg; $CuCl_2 \cdot 2H_2O$, 10 mg; $MnCl_2 \cdot 4H_2O$, 10 mg; $Na_2B_4O_7 \cdot 10H_2O$, 10 mg; $(NH_4)_6Mo_7O_{24} \cdot 4H_2O$, 10 mg.

2° R2YE: 10.3 g of sucrose, 0.025 g of K_2SO_4 , 1.012 g of $MgCl_2 \cdot 6H_2O$, 1 g of glucose, 0.01 g of Difco 25 casamino acids, and 2.2 g of Difco agar are dissolved in 80 ml distilled water and autoclaved. 0.2 ml of trace element solution, 1 ml of KH_2PO_4 (0.5%), 8.02 ml of $CaCl_2 \cdot 2H_2O$ (3.68%), 1.5 ml of L-proline (20%), 10 ml of TES buffer (0.25 M) (pH: 7.2), 0.5 ml of (1 M) NaOH, 5 ml of yeast extract (10%) are sequentially added.

3° TE: 10 mM TRIS HCl, 1 mM EDTA, pH 8.0.

4° YEME: Difco yeast extract (0.3%), Difco peptone (0.5%), oxioid malt extract (0.3%), glucose (1%).

Transformation of *S. lividans* Protoplasts

1. A culture composed of 25 ml YEME, 34% sucrose, 0.005 M $MgCl_2$, 0.5% glycine, in a 250 ml baffled flask, is centrifuged during 30 to 36 hours.
2. The pellet is suspended in 10.3% sucrose and centrifuged. This washing is repeated once.
3. The mycelium is suspended in 4 ml lysozyme solution (1 mg/ml in P medium with $CaCl_2$ and $MgCl_2$ concentrations reduced to 0.0025 M) and incubated at 30° C. for 15 to 60 minutes.
4. The solution is mixed by pipetting three times in a 5 ml pipette and incubated for further 15 minutes.
5. P medium (5 ml) is added and mixed by pipetting as in step 4.
6. The solution is filtered through cotton wool and protoplasts are gently sedimented in a bench centrifuge at 800×G during 7 minutes.
7. Protoplasts are suspended in 4 ml P medium and centrifuged again.
8. Step 7 is repeated and protoplasts are suspended in the drop of P medium left after pouring off the supernatant (for transformation).
9. DNA is added in less than 20 µl TE.
10. 0.5 ml PEG 1 000 solution (2.5 g PEG dissolved in 7.5 ml of 2.5% sucrose, 0.0014 K_2SO_4 , 0.1 M $CaCl_2$, 0.05 M TRIS-maleic acid, pH 8.0, plus trace elements) is immediately added and pipetted once to mix the components.
11. After 60 seconds, 5 ml of P medium are added and the protoplasts are sedimented by gentle centrifugation.
12. The pellet is suspended in P medium (1 ml).
13. 0.1 ml is plated out on R2YE plates (for transformation dry plates to 85% of their fresh weigh e.g. in a laminar flow cabinet).
14. Incubation at 30° C.

A—Construction of a “sfr” Gene Cassette

A “sfr” gene cassette was constructed to allow subsequent cloning in plant expression vectors.

Isolation of a FokI-BglII fragment from the plasmid pBG39 containing a “sfr” gene fragment led to the loss of the first codons, including the initiation codon, and of the last codons, including the stop codon.

This fragment of the “sfr” gene could be reconstructed in vitro with synthetic oligonucleotides which encode appropriate amino-acids.

14

The complementary synthetic oligonucleotides were (SEQ ID NOS:5-6) 5'-CATGAGCCCAGAAC and 3'-TCGGGTCTTGCTGC.

By using such synthetic oligonucleotides, the 5' end of the “sfr” gene could be reformed and the GTG initiation codon substituted for a codon well translated by plant cells, particularly an ATG codon.

The DNA fragment containing the oligonucleotides linked to the “sfr” gene was then inserted into an appropriate plasmid, which contained a determined nucleotide sequence thereafter designated by an “adapter” fragment.

This adapter fragment comprised:

a TGA termination codon which enabled the last codons of the “sfr” gene to be reformed;

appropriate restriction sites which enabled the insertion of the fragment of the nucleotide sequence comprising the “sfr” gene partially reformed with the synthetic oligonucleotides; this insertion resulted in the reconstruction of an intact “sfr” gene;

appropriate restriction sites for the isolation of the entire “sfr” gene.

The “sfr” gene was then inserted into another plasmid, which contained a suitable plant promoter sequence. The plant promoter sequence consisted of the cauliflower mosaic virus promoter sequence (p35S). Of course the invention is not limited to the use of this particular promoter. Other sequences could be chosen as promoters suitable in plants, for example the TR 1'-2' promoter region and the promoter fragment of a Rubisco small subunit gene from *Argemone* hereafter described.

1° Construction of the Plasmid pLK56.2 (FIG. 3)

The construction of plasmid pLK56.2 aimed at obtaining a suitable adaptor including the following sequence of restriction sites: SmaI, BamHI, NcoI, KpnI, BglII, MluI, BamHI, HindIII and XbaI.

The starting plasmids used for this construction, pLK56, pJB64 and pLK33 were those disclosed by BOTTERMAN (ref. 11).

The DNA fragments hereafter described were isolated and separated from low melting point agarose (LGA).

The plasmid pLK56 was cleaved by the enzymes BamHI and NdeI. A NcoI-NdeI fragment (referred to in the drawings by arc “a” in broken line) obtained from plasmid pJB64 was substituted in pLK56 for the BamHI-NdeI fragment shown at “b”. Ligation was possible after filling in the BamHI and NcoI protruding ends with the DNA polymerase I of *E. coli* (Klenow's fragment).

Particularly recircularization took place by means of a T4 DNA ligase. A new plasmid pLK56.3 was obtained.

This plasmid was cleaved by the enzymes XbaI and PstI.

The BamHI-PstI fragment of pLK33 (c) (on FIG. 3) was substituted for the XbaI-PstI fragment (d) of pLK56.3, after repairing of their respective ends by Klenow's fragment.

After recircularization by means of the T4 DNA ligase, the obtained plasmid pLK56.2 contained a nucleotide sequence which comprised the necessary restriction sites for the subsequent insertion of the “sfr” gene.

2° Construction of the Plasmid pGSH150 (FIG. 4A)

Parallel with the last discussed construction, there was produced a plasmid containing a promoter sequence recognized by the polymerases of plant cells and including suitable restriction sites, downstream of said promoter sequence in the direction of transcription, which restriction sites are then intended to enable the accommodation of the “sfr” gene then obtainable from pLK56.2, under the control of said plant promoter.

15

Plasmid pGV825 is described in DEBLAERE et al. (ref. 10). Plasmid pJB63 is from BOTTERMAN (ref. 11).

pGV825 was linearized with PvuII and recircularized by the T4 DNA ligase, resulting in the deletion of an internal PvuII fragment shown at (e), (plasmid pGV956).

pGV956 was then cleaved by BamHI and BglII.

The BamHI-HindIII fragment (f) obtained from pJB63 was dephosphorylated with calf intestine phosphatase (CIP) and substituted for the BamHI-BglII fragment of pGV956.

Plasmid pGV1500 was obtained after recircularization by means of T4 DNA ligase.

An EcoRI-HindIII fragment obtained from plasmid pGSH50 was purified. The latter plasmid carried the dual TR 1'-2' promoter fragment described in VELTEN et al., (ref. 13). This fragment was inserted in pGV1500, digested with HpaI and HindIII and yielded pGSH150.

This plasmid contains the promoter fragment in front of the 3' end of the T-DNA transcript 7 and a BamHI and ClaI sites for cloning.

3° Construction of the Plasmid pGSJ260 (FIG. 4B)

CP3 is a plasmid derived from pBR322 and which contains the 35S promoter region of cauliflower mosaic virus within a BamHI fragment.

pGSH150 was cut by BamHI and BglII.

The BamHI-BglII fragment (h) of CP3, which contained the nucleotide sequence of p35S promoter, was substituted for the BamHI-BglII fragment (i) in pGSH150 to form plasmid pGSJ250. pGSJ250 was then opened at its BglII restriction site.

A BamHI fragment obtained from mGV2 (ref. 12) was inserted in pGSJ250 at the BglII site to form plasmid pGSJ260.

However prior to inserting the "sfr" gene obtainable from pLK56.2 into plasmid pGSJ260, it was still desirable to further modify the first in order to permit insertion in a more practical manner. Thus pLK56.2 was further modified as discussed below to yield pGSR1.

Starting from plasmid pGSJ260, two plasmid constructions for subsequent transformations of plant cells were made:

a first plasmid permitting the expression of the "sfr" gene in the cytoplasm of plant cells, and

a second plasmid so modified as to achieve transport of the Bialaphos-resistance enzymes to the chloroplasts of plant cells.

First Case: Plasmid Enabling the Expression of the "sfr" Gene in the Cytoplasm of Plant Cells

Cloning of the sfr Gene Cassette in a Plant Expression Vector (pGSR2) (FIG. 5)

On FIG. 5A (SEQ ID NO:13), the nucleotide sequence of the adapter of pLK56.2 is shown. In particular, the locations of BamHI, NcoI, BglII restriction sites are shown.

This adapter fragment was cleaved by the enzymes NcoI and BglII.

FIG. 5B (SEQ ID NO:14) shows the FokI-BglII fragment (j) obtained from pBG39. The locations of these two restriction sites are shown on FIG. 2.

Using synthetic oligonucleotides, the first codons of the "sfr" gene were reformed, particularly the 5' end of the gene in which a ATG initiation codon was substituted for the initial GTG codon.

This FokI-BglII fragment completed with the synthetic oligonucleotides was then substituted in pLK56.2 for the NcoI-BglII fragment of the adapter. The 3' end of the gene was thus reformed too, after recircularization with T4 DNA ligase. The plasmid obtained, pGSR1, thus contained the entire "sfr" gene inserted in its adapter.

16

The plasmid pGSJ260 was then opened by BamHI (FIG. 5C) and the BamHI fragment obtained from pGSR1, which contained the entire "sfr" gene, was inserted into pGSJ260.

The obtained plasmid, pGSR2 (see FIG. 6A) contained a pBR322 replicon, a bacterial streptomycin resistance gene (SDM-SP-AD-transferase) and an engineered T-DNA consisting of:

the border fragments of the T-DNA;

a chimeric kanamycin gene which provided a dominant selectable marker in plant cells; and

a chimeric "sfr" gene.

The chimeric "sfr" gene consisting of:

the promoter region of the cauliflower mosaic virus (p35S);

the "sfr" gene cassette as described in FIG. 5;

the 3' untranslated region, including the polyadenylation signal of T-DNA transcript 7.

pGSR2 was introduced into *Agrobacterium tumefaciens* recipient C58ClRif® (pGV2260) according to the procedure described by DEBLAERE et al. (ref. 10).

This strain was used to introduce the chimeric "sfr" gene in *N. tabacum* SR₁ plants.

Two variant plasmids deriving from pGSR2, namely pGSFR280 and pGSFR281, have been constructed. They differ in the untranslated sequence following the transcription initiation site. In pGSR2, this fragment consists of the following sequence (SEQ ID NO:7):

GAGGACACGCTGAAATCACCAGTCTCGGATCC
ATG;

while it consists of (SEQ ID NO:8):

GAGGACACGCTGAAATCACCAGTCTCTC-
TACAAATCGATCCATG

in pGSR280 and of (SEQ ID NO:9)

GAGGACACGCTGAAATCACCAGTCTCTC-
TACAAATCGATG

in pGSR281, with an ATG codon being the initiation codon of the "sfr" gene. The "sfr" gene is also fused to the TR1'-2' promoter in the plasmid pGSH150 (FIG. 4A) yielding pGSFR160 and pGSFR161 (FIG. 6B). These plasmids contain slight differences in the pTR2 "sfr" gene configuration: the "sfr" gene is correctly fused to the endogenous gene 2' ATG in pGSFR161 (for sequences see ref. 13), whereas 4 extra base pairs (ATCC) are present just ahead of the ATG codon in pGSFR160. Otherwise, plasmids p65FR161 and p65FR160 are completely identical.

All plasmids are introduced in *Agrobacterium* by cointegration in the acceptor plasmids pGV2260 yielding the respective plasmids pGSFR1280, pGSFR1281, pGSFR1160 and pGSFR1161.

Second Case: Construction of a Plasmid Containing the "sfr" Gene Downstream of a DNA Sequence Encoding a Transit Peptide and Suitable for Achieving Subsequent Translocation of the "sfr" Gene Expression Product into Plant-Cell-Chloroplasts

In another set of experiments, the nucleotide sequence which contained the "sfr" gene was fused to a DNA sequence encoding a transit peptide so as to enable its transport into chloroplasts.

A fragment of the "sfr" gene was isolated from the adapter fragment above described and fused to a transit peptide. With synthetic oligonucleotides, the entire "sfr" gene was reconstructed and fused to a transit peptide.

The plasmid (plasmid pATS3 mentioned below) which contained the nucleotide sequence encoding the transit peptide comprised also the promoter sequence thereof.

Construction of the Plasmid pGSR4 which Contains the "sfr" Gene Fused to a DNA Sequence Encoding Transit Peptide (FIG. 7)

Plasmid pLK57 is from BOTTERMAN, (ref. 11). Plasmid pATS3 is a pUC19 clone which contains a 2 Kb EcoRI genomic DNA fragment from Arabidopsis thaliana comprising the promoter region and the transit peptide nucleotide sequence of the gene, the expression thereof is the small subunit of ribulose biphosphate carboxylase (ssu). The A. thaliana small subunit was isolated as a 1 500 bp EcoRI-SphI fragment. The SphI cleavage site exactly occurs at the site where the coding region of the mature ssu protein starts.

Plasmids pLK57 and pATS3 were opened with EcoRI and SphI. After recircularization by means of the T4 DNA ligase, a recombinant plasmid pLKAB1 containing the sequence encoding the transit peptide (Tp) and its promoter region (Pssu) was obtained.

In order to correctly fuse the “sfr” gene at the cleavage site of the signal peptide, the N-terminal gene sequence was first modified. Since it was observed that N-terminal gene fusions with the “sfr” gene retain their enzymatic activity, the second codon (AGC) was modified to a GAC, yielding an NcoI site overlapping with the ATG initiator site. A new plasmid, pGSSFR2 was obtained. It only differs from pGSR1 (FIG. 5B), by that mutation. The NcoI-BamHI fragment obtained from pGSR2 was fused at the SphI end of the transit peptide sequence. In parallel, the “sfr” gene fragment was fused correctly to the ATG initiator of the ssu gene (not shown in figures).

Introduction of the “sfr” Gene into a Different Plant Species

The Bialaphos-resistance induced in plants by the expression of chimeric genes, when the latter have been transformed with appropriate vectors containing said chimeric genes, has been demonstrated as follows. The recombinant plasmids containing the “sfr” gene were introduced separately by mobilization into Agrobacterium strain C58C₁ Rif® (pGV2260) according to the procedure described by DEBLAERE and al., Nucl. Acid. Res., 13, p. 1 477, 1985. Recombinant strains containing hybrid Ti plasmides were formed. These strains were used to infect and transform leaf discs of different plant species, according to a method essentially as described by HORSH and al., 1985, Science, vol. 227. Transformation procedure of these different plant species given by way of example, is described thereafter.

1. Leaf Disc Transformation of Nicotiana tabacum

Used Media are described thereafter:

A ₁ MS salt/2	+1% sucrose 0.8% agar pH 5.7	
A ₁₀ B5-medium	+250 mg/l NH ₄ NO ₃ 750 mg/l CaCl ₂ 2H ₂ O 0.5 g/l 2-(N-Morpholino)ethane-sulfonic acid (MES) pH 5.7	
A ₁₁ B5-medium	30 g/l sucrose +250 mg/l NH ₄ NO ₃ 0.5 g/l MES pH 5.7 2% glucose 0.8% agar 40 mg/l adenine +1 mg/l 6-Benzylaminopurine (BAP) 0.1 mg/l Indole-3-acetic acid (IAA)	
A ₁₂ B5-medium	500 mg/l Claforan +250 mg/l NH ₄ NO ₃ 0.5 g/l MES pH 5.7 2% glucose 0.8% agar 40 mg/l adenine	

-continued

A ₁₃ MS-salt/2	+1 mg/l BAP 200 mg/l claforan +3% sucrose 0.5 MES g/l pH 5.7 0.7% agar 200 mg/l claforan
Bacterial medium = min A:	(Miller 1972) 60 mM K ₂ HPO ₄ , 3H ₂ O, 33 mM KH ₂ PO ₄ ; 7.5 mM (NH ₄) ₂ SO ₄ 1.7M trinatricitrat; 1 mM MgSO ₄ ; 2 g/l glucose; 50 mg/l vitamine B1

Plant Material:

Nicotiana tabacum cv. Petit Havana SR1

Plants are used 6 to 8 weeks after subculture on medium A₁

Infection:

midribs and edges are removed from leaves.

Remaining parts are cut into segments of about 0.25 cm² and are placed in the infection medium A₁₀ (about 12 segments in a 9 cm Petri dish containing 10 ml A₁₀).

Segments are then infected with 25 µl per Petri dish of a late log culture of the Agrobacterium strain grown in min A medium.

Petri dish are incubated for 2 to 3 days at low light intensity.

After 2 to 3 days medium is removed and replaced by 20 ml of medium A₁₀ containing 500 mg/l clarofan.

Selection and Shoot Induction

The leaf discs are placed on medium A₁₁ containing a selective agent:

100 mg/l kanamycin and

10 to 100 mg/l phosphinotricin.

Leaf discs are transferred to fresh medium weekly.

After 3 to 4 weeks regenerating calli arise. They are separated and placed on medium A₁₂ with the same concentration of selective agent as used for the selection.

Rooting

After 2 to 3 weeks the calli are covered with shoots, which can be isolated and transferred to rooting medium A₁₃ (without selection).

Rooting takes 1 to 2 weeks.

After a few more weeks, these plants are propagated on medium A₁.

2. Tuber Disc Infection of Solanum tuberosum (Potato)

Used media are described thereafter:

C ₁ B5-medium	+250 mg/l NH ₄ NO ₃ 300 mg/l (CaCH ₂ PO ₄) ₂ 0.5 g/l MES pH 5.7 0.5 g/l polyvinylpyrrolidone (PVP) 40 g/l mannitol (=0.22M) 0.8% agar 40 mg/l adenine
C ₂ B5-medium	+250 mg/l NH ₄ NO ₃ 400 mg/l glutamine 0.5 g/l MES pH 5.7 0.5 g/l PVP 40 g/l mannitol 40 mg/l adenine
C ₅ MS salt/2	0.8% agar +0.5 mg/l transzeatine 0.1 mg/l IAA 500 mg/l claforan
C ₇ B5-medium	+3% sucrose 0.7% agar pH 5.7 +250 mg/l NH ₄ NO ₃ 400 mg/l glutamine

19		20	
-continued		3. Leaf Disc Infection of Lycopersicum esculentum (Tomato)	
		Used media are described thereafter	
C ₈ B5-medium	0.5 g/l MES pH 5.7	5	<hr/>
	0.5 g/l PVP		
	20 g/l mannitol		
	20 g/l glucose		
	40 mg/l adenine		
	0.6% agarose		
	+0.5 mg/l transzeatine		
	0.1 mg/l IAA		
	500 mg/l clarofan		
	+250 mg/l NH ₄ NO ₃		
C ₉ B5-medium	400 mg/l glutamine	10	<hr/>
	0.5 g/l MES pH 5.7		
	0.5 g/l PVP		
	20 g/l mannitol		
	20 g/l glucose		
	40 mg/l adenine		
	0.6% agarose		
	+200 mg/l clarofan		
	1 mg/l transzeatine		
	+250 mg/l NH ₄ NO ₃		
C ₁₁ MS salt/2	400 mg/l glutamine	15	<hr/>
	0.5 g/l MES pH 5.7		
	0.5 g/l PVP		
	20 g/l mannitol		
	20 g/l glucose		
	40 mg/l adenine		
	0.6% agarose		
	+1 mg/l transzeatine		
	0.01 mg/l Gibberellic acid A ₃ (GA ₃)		
	100 mg/l clarofan		
Bacterial medium = min A:	+6% sucrose	20	<hr/>
	0.7% agar		
	(Miller 1972 60 mM K ₂ HPO ₄ •3H ₂ O;		
	33 mM KH ₂ PO ₄ ; 7.5 mM (NH ₄) ₂ SO ₄ ;		
	1.7 trisodiumcitrat; 1 mM		
	MgSO ₄ ;		
	2 g/l glucose; 50 mg/l vitamine B1.		
Plant Material		25	<hr/>
	Tubers of Solanum tuberosum c.v Berolina		
	c.v Désirée		
	Infection		
	Potatoes are peeled and washed with water.		
	Then they are washed with concentrated commercial		
	bleach for 20 minutes, and		
	rinsed 3 to 5 times with sterile water.		
	The outer layer is removed (1 to 1.5 cm)		
	The central part is cut into discs of about 1 cm ² and 2 to 3		
mm thick.		30	<hr/>
	Discs are placed on medium C ₁ (4 pieces in a 9 cm Petri		
	dish).		
	10 µl of a late log culture of an Agrobacterium strain grown		
	in min A medium is applied on each disc.		
	Discs are incubated for 2 days at low light intensity.		
	Selection and Shoot Induction		
	Discs are dried on a filter paper and transferred to medium		
	C ₂ with 100 mg/l kanamycin.		
	After one month small calli are removed from the discs and		
transferred to medium C ₇ containing 50 mg/l kanamycin.			
After a few more weeks, the calli are transferred to medium		35	<hr/>
	C ₈ containing 50 mg/l kanamycin.		
	If little shoots start to develop, the calli are transferred to		
	elongation medium C ₉ containing 50 mg/l Kanamycin.		
	Rooting		
	Elongated shoots are separated and transferred to rooting		
	medium C ₁₁ .		
	Rooted shoots are propagated on medium C ₅ .		
		40	<hr/>
		45	<hr/>
		50	<hr/>
		55	<hr/>
		60	<hr/>
		65	<hr/>

-continued

B ₉ B5-medium	+250 mg/l NH ₄ NO ₃ 0.5 g/l MES pH 5.7 0.5 g/l PVP 2% glucose 0.6% agarose 40 mg/l adenine +1 mg/l transzeatine 0.01 mg/l GA3
Bacterial medium = min A:	(Miller 1972) 60 mM K ₂ HPO ₄ •3H ₂ O; 33 mM KH ₂ PO ₄ ; 7.5 mM (NH ₄) ₂ SO ₄ ; 1.7M trisodiumcitrat; 1 mM MgSO ₄ ; 2 g/l glucose; 50 mg/l vitamin B1

Plant Material

Lycopersicum esculentum cv. Lucullus.

Plants are used 6 weeks after subculture on medium A₁.

Infection

Midrib is removed from the leaves.

Leaves are cut in segments of about 0.25 to 1 cm² (the edges of the leaves are not wounded, so that only maximum 3 sides of the leaf pieces is wounded).

Segments are placed in infection medium B₁ (upside down), about 10 segments in a 9 cm Petri dish.

Segments are then infected with 20 µl per Petri dish of a late log culture of the *Agrobacterium* strain grown in min A medium.

Petri dishes incubate for 2 days at low light intensity.

Medium is removed after 2 days and replaced by 20 ml of medium B₁ containing 500 mg/l clarofan.

Selection and Shoot Induction

The leaf discs are placed in medium B₂+50 or 100 mg/l kanamycin.

Each 5 days the osmotic pressure of the medium is lowered by decreasing the mannitol concentration, transfers are done consecutively in medium B₃, B₄, B₅, and B₆.

After one month calli with meristems are separated from the leaf discs and placed on medium B₇ with 50 or 100 mg/l kanamycin.

Once little shoots have formed, calli are transferred to elongation medium B₉ with 50 or 100 mg/l kanamycin.

Rooting

Elongated shoots are separated and transferred to medium B₈ for rooting.

Plants are propagated on medium A₁.

Greenhouse Tests for Herbicide Resistance

Material and Method

In this experiment, two herbicides comprising phosphinotricin as active ingredient, are used.

These compounds are those commercially available under the registered trademarks BASTA® and MEIJI HERBI-ACE®.

These products are diluted to 2% with tap water. Spraying is carried out on a square meter area from the four corners. Temperature of the greenhouse is about 22° C. for tobaccos and tomato, and above 10° C. to 15° C. for potato.

Results

Tobacco Spraytest

a) *Nicotiana tabacum* cv. Petit Havana SR1 plants transformed with the chimeric "sfr" genes as present in pGSFR1161 or pGSFR1281, as well as untransformed control plants (from 10 cm to 50 cm high) are treated with 20 l BASTA®/ha. Control SR1 plants die after 6 days, while transformed plants are fully resistant to 20 l BASTA®/ha and continue growing undistinguishable from untreated plants. No visible damage is detected, also the treatment is repeated

every two weeks. The treatment has no effect in subsequent flowering. The recommended dose of BASTA® herbicide in agriculture is 2.5-7.5 l/ha.

b) A similar experiment is performed using 8 l/ha MEIJI HERBIACE®. The transformed plants (the same as above) are fully resistant and continue growing undistinguishable from untreated plants. No visible damage is detectable.

Potato Spraytest

Untransformed and transformed potato plants (*Solanum tuberosum* cv. Berolina) (20 cm high) with the chimeric "sfr" gene as present in pGSFR1161 or pGSFR1281 are treated with 20 l BASTA®/ha. Control plants die after 6 days while the transformed plants do not show any visible damage. They grow undistinguishable from untreated plants.

Tomato Spraytest

Untransformed and transformed tomato plants (*Lycopersium esculentum* c.v. luculus) (25 cm high) with the chimeric "sfr" gene as present in pGSFR1161 and pGSFR1281 are treated with 20 l BASTA®/ha. control plants die after six days while transformed plants are fully resistant. They do not show any visible damage and grow undistinguishable from untreated plants.

Growth Control of Phytopathogenic Fungi with Transformed Plants

In another set of experiments, potato plants expressing chimeric "sfr" genes as present in pGSFR1161 or pGSFR1281 are grown in a greenhouse compartment at 20° C. under high humidity. Plants are inoculated by spraying 1 ml of a suspension of 10⁶ *Phytophthora infestans* spores per ml. Plants grow in growth chambers (20° C., 95% humidity, 4 000 lux) until fungal disease symptoms are visible (one week). One set of the plants are at that moment sprayed with Bialaphos at a dose of 8 l/ha. Two weeks later, untreated plants are completely ingested by the fungus. The growth of the fungus is stopped on the Bialaphos treated plants and no further disease symptoms evolve. The plants are effectively protected by the fungicide action of Bialaphos.

Transmission of the PPT Resistance through Seeds

Transformed tobacco plants expressing the chimeric "sfr" gene present in pGSFR1161 and pGSFR1281 are brought to flowering in the greenhouse. They show a normal fertility.

About 500 F1 seeds of each plant are sown in soil, F1 designating seeds of the first generation, i.e directly issued from the originally transformed plants. When seedlings are 2-3 cm high, they are sprayed with 8 l BASTA®/ha. 7 days later, healthy and damaged plants can be distinguished in a ratio of approximately 3 to 1. this shows that PPT resistance is inherited as a dominant marker encoded by a single locus.

10 resistant F1 seedlings are grown to maturity and seeds are harvested. F2 seedlings are grown as described above and tested for PPT-resistance by spraying BASTA® at a dose of 8 l/ha. Some of the F1 plants produce F2 seedlings which are all PPT-resistant showing that these plants are homozygous for the resistance gene. The 5 invention also concerns plant cells and plants non-essentially-biologically-transformed with a GS inhibitor-inactivating-gene according to the invention.

In a preferred embodiment of the invention, plant cells and plants are non-biologically-transformed with the "sfr" gene hereabove described.

Such plant cells and plants possess, stably integrated in their genome, a non-variety-specific character which render them able to produce detectable amounts of phosphinotricinacetyl transferase.

This character confers to the transformed plant cells and plants a non-variety-specific enzymatic activity capable of inactivating or neutralizing GS inhibitors like Bialaphos and PPT.

Accordingly, plant cells and plants transformed according to the invention are rendered resistant against the herbicidal effects of Bialaphos and related compounds.

Since Bialaphos was first described as a fungicide, transformed plants can also be protected against fungal diseases by spraying with the compound several times.

In a preferred embodiment, Bialaphos or related compounds is applied several times, particularly at time intervals of about 20 to 100 days.

The invention also concerns a new process for selectively protecting a plant species against fungal diseases and selectively destroying weeds in a field comprising the steps of treating a field with an herbicide, wherein the plant species contain in their genome a DNA fragment encoding a protein having an enzymatic activity capable of neutralizing or inactivating GS inhibitors and wherein the used herbicide comprises as active ingredient a GS inhibitor.

It comes without saying that the process according to the invention can be employed with the same efficiency, either to only destroy weeds in a field, if plants are not infected with fungi, either to only stop the development of fungi if the latter appears after destruction of weeds.

In a preferred embodiment of the process according to the invention, plant species are transformed with a DNA fragment comprising the "sfr" gene as described hereabove, and the used herbicide is PPT or a related compound.

Accordingly, a solution of PPT or related compound is applied over the field, for example by spraying, several times after emergence of the plant species to be cultivated, until early and late germinating weeds are destroyed.

It is quite evident that before emergence of plant species to be cultivated, the field can be treated with an herbicidal composition to destroy weeds.

On the same hand, fields can be treated even before the plant species to be cultivated are sowed.

Before emergence of the desired plant species, fields can be treated with any available herbicide, including Bialaphos-type herbicides.

After emergence of the desired plant species, Bialaphos or related compound is applied several times.

In a preferred embodiment, the herbicide is applied at time intervals of about from 20 to 100 days.

Since plants to be cultivated are transformed in such a way as to resist to the herbicidal effects of Bialaphos-type herbicides, fields can be treated even after emergence of the cultivated plants.

This is particularly useful to totally destroy early and late germinating weeds, without any effect on the plants to be produced.

Preferably, Bialaphos or related compound is applied at a dose ranging from about 0.4 to about 1.6 kg/ha, and diluted in a liquid carrier at a concentration such as to enable its application to the field at a rate ranging from about 2 to about 8 l/ha.

There follows examples, given by way of illustration, of some embodiments of the process with different plant species.

Sugarbeets

The North European sugarbeet is planted from March 15 up to April 15, depending upon the weather condition and more precisely on the precipitation and average temperature. the weeds problems are more or less the same in each country and can cause difficulties until the crop closes its canopy around mid-July.

Weed problems can be separated in three situations:

early germination of the grassy weeds,
early germinating broadleaved weeds,
late germinating broadleaved weeds.

Up to now, pre-emergence herbicides have been successfully used. Such compounds are for example those commercially available under the registered trademarks: PYRAMIN®, GOLTIX® and VENZAR®. However, the susceptibility to dry weather conditions of these products as well as the lack of residual activity to control late germinating weeds have led the farmer to use post-emergence products in addition to pre-emergence ones.

Table (I) thereafter indicates the active ingredients contained in the herbicidal compositions cited in the following examples.

TABLE I

Commercial Name	Active Ingredient	Formulation
AVADEX ^R	Diallate	EC 400 g/l
AVADEX BW ^R	Triallate	EC 400 g/l
GOLTIX ^R	Metamitron	WP 70%
RONEET ^R	Cycloate	EC 718 g/l
TRAMAT ^R	Ethofumate	EC 200 g/l
FERVINAL ^R	Alloxydime-sodium	SP 75%
BASTA ^R	Phosphinotricin	200 g/l
PYRAMIN FL ^R	Chloridazon	SC 430 g/l

According to the invention, post-emergence herbicides consist of Bialaphos or related compounds, which offer a good level of growth control of annual grasses (Bromus, Avena spp., Alopecurus, POA) and broadleaves (Galium, Polygonum, Senecio, Solanum, Mercurialis).

Post-emergence herbicides can be applied at different moments of the growth of sugarbeet; at a cotyledon level, two-leave level or at a four-leave level.

Table (II) thereafter represents possible systems of field-treatment, given by way of example.

In those examples, the post-emergence herbicide of the class of Bialaphos used is BASTA®, in combination with different pre-emergence herbicides. Concentrations are indicated in l/ha or kg/ha.

TABLE II

POSSIBLE WEEDCONTROL SYSTEMS IN SUGARBEETS, BASED ON THE USE OF BASTA ^R , PROVIDING BEETS ARE MADE RESISTANT AGAINST THE LATTER CHEMICAL (in lt or kg/ha).				
Pre-sowing	Pre-emergence	Cotyledons	Two-leaves	Four leaves
1. AVADEX ^R 3.5 lt	—	BASTA ^R 3 lt	BASTA ^R /tramat 3 lt 1.5 lt	—
2. AVADEX ^R 3.5 lt	GOLTIX ^R 4 kg	—	—	—
3. RONEET ^R 4 lt	GOLTIX ^R 5 kg	—	—	—

TABLE II-continued

POSSIBLE WEEDCONTROL SYSTEMS IN SUGARBEETS, BASED ON THE USE OF BASTA ^R , PROVIDING BEETS ARE MADE RESISTANT AGAINST THE LATTER CHEMICAL (in lt or kg/ha).				
Pre-sowing	Pre-emergence	Cotyledons	Two-leaves	Four leaves
4. RONEET ^R 4 lt	GOLTIX ^R 2.5 kg	—	BASTA ^R 3 lt	—
5. TRAMAT ^R 5 lt	—	—	BASTA ^R 3 lt	BASTA ^R /GOLTIX ^R 2 lt 2 kg
6. —	GOLTIX ^R 2.5 kg	—	BASTA ^R 3 lt	—
7. —	—	BASTA ^R /tramat 3 lt 1.7 lt	—	BASTA ^R /GOLTIX ^R 3 lt 2 kg
8. PYRAMIN ^R 6 lt	—	BASTA ^R 3 lt	Venzar 1 kg	—
9. —	—	BASTA ^R 3 lt	BASTA ^R /GOLTIX ^R 3 lt 2 kg	—
10. DIALLATE ^R PYRAMIN ^R 3.5 lt 6 lt	—	—	BASTA ^R /Metamitron 3 lt 1 kg	—

Potatoes

Potatoes are grown in Europe on about 8.10⁶ Ha. The major products used for weed control are Linuron/mono-linuron or the compound commercially available under the denomination METRABUZIN

These products perform well against most weedspecies.

However, weeds such as Galium and Solanum plus late germinating Chenopodium and Polygonum are not always effectively controlled, while control of the annual grasses is also sometime erratic.

Once again, late germinating broadleaved weeds are only controllable by post-emergence applications of herbicides such as BASTA®.

Table (III) thereafter represents some examples given by way of example of field-treatment in the case of potatoes.

TABLE III

Weeds control systems in potatoes, based on the use of BAST ^R , providing potatoes are rendered resistant to BASTA ^R .	
Linuron + monolinuron	(375 g + 375 g/ha) prior to emergence
BASTA ^R	3-4 lt/ha after emergence (5-15 cm)
BASTA ^R /fluazifop-butyl	3-4 lt/ha + 2 lt/ha after emergence (5-15 cm)
Linuron	WP 50% (AFALON ^R)
Monolinuron	WP 47.5% (ARESSIN ^R)
fluazifop-butyl	EL 250 g/l (FUSILADE ^R)

The strains PGSJ260 and pBG39 used hereabove have been deposited on Dec. 12, 1985, at the “German Collection of Micro-organisms” (DEUTSCHE SAMMLUNG VON MIKROORGANISMEN) at Gottingen, Germany. They received the deposition numbers DSM 3 606 and DSM 3 607 respectively.

Further embodiments of the invention are described hereafter with reference to the figures in which:

FIG. 8 shows the restriction map of a plasmid pJS1 containing another Bialaphos-resistance-gene;

FIG. 9 (SEQ ID NO:10) shows the nucleotide sequence of the “sfrsv” gene containing the resistance gene;

FIG. 10 (SEQ ID NO:18) shows the amino acid homology of “sfrsv” gene and “sfr” gene,

FIG. 11 (SEQ ID NOS:19-21) shows the construction of a plasmid, given by way of example, which contains the “sfrsv” gene and suitable for the transformation of plant cells.

Another Bialaphos-resistance-gene has been isolated form another Bialaphos-producing-strains, i.e. streptomyces viridochromogenes. This second resistance-gene is thereafter designated by “sfrsv” gene.

This second preferred DNA fragment according to the invention, for the subsequent transformation of plant cells, consists of a nucleotide sequence (SEQ ID NO:11) coding for at least part of a polypeptide having the following sequence:

V S P E R R P V E I R P A T A A D M
A A V C D I V N H Y I E T S T V N P
R T E P Q T P Q E W I D D L E R L Q
D R Y P W L V A E V E G V V A G I A
Y A G P W K A R N A Y D W T V E S T
V Y V S H R H Q R L G L G S T L Y T
H L L K S M E A Q G F K S V V A V I
G L P N D P S V R L H E A L G Y T A
R G T L R A A G Y K H G G W H D V G
F W Q R D F E L P A P P R P V R P V
T Q I *

which part of said polypeptide is of sufficient length to confer protection against Bialaphos-“plant-protecting-capability”-, to plant cells, when incorporated genetically and expressed therein. Reference will also be made hereafter to the “plant-protecting-capability”against Bialaphos of the abovesaid nucleotide sequence.

Meaning of the designation of amino acids by a single letter is given thereafter.

Alanine	A	Leucine	L
Arginine	R	Lysine	K
Asparagine	N	Methionine	M
Aspartic Acid	D	Phenylalanine	F
Cysteine	C	Proline	P
Cystine	C	Serine	S
Glycine	G	Threoriine	T
Glutamic Acid	E	Tryptophan	W
Glutamine	Q	Tyrosine	Y
Histidine	H	Valine	V
Isoleucine	I		

This second preferred DNA fragment consists of the following nucleotide sequence (SEQ ID NO:12):

TAAAGAGGTGCCCGCCACCCGCTTTCGAGAACACCGAAGGAGACCACAC
↓
GTGAGCCCAGAACGACGCCCGGTGAGATCCGTCCCGCCACCGCCGCCGA
CATGGCGGGCTGCGACATCGTCAATCACTACATCGAGACGAGCACGG
TCAACTTCCGTACGGAGCCGAGACTCCGCAGGAGTGGATCGACGACCTG
GAGCGCTCCAGGACCGCTACCCCTGGCTCGTCGCCGAGGTGGAGGGCGT
CGTCGCCGGCATCGCCTACGCCGGCCCCCTGGAAGGCCCGCAACGCCTACG
ACTGGACCGTCGAGTCGACGGTGTACGTCTCCACCGGCACCGCGGCTC
GGACTGGGCTCCACCCTCTACACCCACCTGCTGAAGTCCATGGAGGCCCA
GGGCTTCAAGAGCGTGGTGCCTCATCGGACTGCCCAACGACCCGAGCG
TGCGCCTGCACGAGGCGCTCGGATACACCGCGCGGGACGCTGCGGGCA
GCCGGCTACAAGCACGGGGCTGGCACGACGTGGGGTTCTGGCAGCGCGA
CTTCGAGCTGCCGGCCCCCGCCCCGCTCCGGCCCGTACACAGATCT
↑
GAGCGGAGAGCGCATGGC

or of a part thereof expressing a polypeptide having plant-protecting capability against Bialaphos;

There follows hereafter the description of experiments carried out for the isolation of the "sfrsv" resistance gene, the construction of expression vectors which contain the resistance gene and which allow the subsequent transformation of plant cells, in order to render them resistant to GS inhibitors.

Cloning of the Bialaphos-Resistance-"sfrsv" Gene from *Streptomyces viridochromogenes*

The strain *Streptomyces viridochromogenes* DSM 40736 (ref 1) was grown and total DNA of this strain was prepared according to standard techniques. DNA samples were digested respectively with PstI, SmaI and Sau3AI in three different reactions and separated on an agarose gel, together with plasmid DNA from pGSR1 (FIG. 5B) digested with BamHI. In a Southern analysis the DNA was blotted on a nitrocellulose filter and hybridized with the labeled BamHI fragment from pGSR1 containing the "sfr" gene. In all four lanes of the gel, a restriction fragment was showing strong homology with the probe: a PstI fragment of about 3 kb, a SmaI fragment of about 1.2 kb and Sau3AI fragment of 0.5 kb. In order to clone this gene, PstI restriction fragments were directly cloned in the *Escherichia coli* vector pUC8. 3000 colonies obtained after transformation were transferred to nitrocellulose filters, and hybridized with the "sfr" probe. Positive candidates were further tested for their growth on minimal medium plates containing 300 µg/ml PPT. One transformant that grew on PPT-containing-medium was further analysed. The plasmid map and relevant restriction sites of this plasmid pJS1 are represented in FIG. 8. The strain MC1061 (pJS1) has been deposited on Mar. 6, 1987 at the DEUTSCHE SAMMLUNG VON MIKROORGANISMEN (DSM) under deposition number DSM 4023. The clone restriction fragment has been sequenced according to the Maxam and Gilbert method and the coding region of the gene could be identified through homology. The sequence of the "sfrsv" gene is represented in FIG. 9 and the homology on the nucleotide and amino acid sequence level with "sfr" gene is shown in FIG. 10.

Expression of the "sfrsv" Gene

A "sfrsv gene cassette" was also constructed to allow subsequent cloning in plant expression vectors. A BanII-BglII

fragment containing the "sfrsv" coding region without the initiation codon GTG was isolated from pJS1. This fragment was ligated in the vector pLK56-2 digested with NcoI and BglII, together with a synthetic oligonucleotide 5'-CAT-GAGCC-3', similar with the one described for "sfr" gene and shown in FIG. 5. The construction of pGSR1SV is schematically shown in FIG. 11. Since similar cassettes of both genes are present in respectively pGSR1 and pGSR1SV, previous described constructions for the expression of the "sfr" gene in plants can be repeated.

Enzymatic analysis of crude extracts from *E. coli* strains carrying plasmid pGSR1SV demonstrated the synthesis of an acetylase which could acetylate PPT. This was shown by thin layer chromatography of the reaction products.

The "sfrsv" gene was then inserted into the plasmid vector pGSJ260 (FIG. 4B) under the control of the CaMV 35s promoter, to yield a plasmid pGS2SV, similar to pGSR2 (FIG. 6A) except that the "sfrsv" gene is substituted for the "sfr" gene.

It is clear that herbicide resistance genes of the above type may be obtained from many other microorganisms that produce PPT or PPT derivatives. Herbicide resistance gene can then be incorporated in plant cells with a view of protecting them against the action of such Glutamine Synthetase inhibitors. For instance, a Bialaphos-resistance-gene is obtained from *Kitasotosporia* (ref. 15).

Transformed plant cells and plants which contain the "sfrsv" resistance gene can be obtained with plasmid pGSR2SV, using the same *Agrobacterium*-mediated-transformation system as hereabove described for the transformation of different plant species with the "sfr" gene.

Plants are regenerated and tested for their resistance with similar spraying tests as described hereabove. All plants behaved similarly and show resistance against herbicides consisting of glutamine synthetase inhibitors.

Finally, the inventors also pertains to the combination of the plants resistant to an inhibitor of glutamine synthetase as defined above with the corresponding inhibitor of glutamine synthetase for use in the production of cultures of said plants free from weeds.

REFERENCES

1. BAYER et al., *HELVETICA CHIMICA ACTA*, 1972
2. WAKABAYASHI K. and MATSUNAKA S., *Proc. 1982, British Crop Protection Conference*, 439-450
3. M. MASON et al., *PHYTOCHEMISTRY*, 1982, vol. 21, No. 4, p. 855-857.
4. C. J. THOMPSON et al., *NATURE*, Jul. 31, 1980, vol. 286, No. 5 772, p. 525-527
5. C. J. THOMPSON et al., *JOURNAL OF BACTERIOLOGY*, August 1982, p. 678-685
6. C. J. THOMPSON et al., *GENE* 20, 1982, p. 51-62
7. C. J. THOMPSON et al., *MOL. GEN. GENET.*, 1984, 195, p. 39-43
8. TOWBIN et al., *PROC. NATL. ACAD. SCI. USA*, 1979, 76, p. 4 350-4 354
9. *METHODS OF ENZYMOLOGY*, V.XLIII, p. 737-755
10. DEBLAERE H. et al., 1985, *Nucl. Acid. Res.*, 13, 1 477
11. BOTTERMAN J., February 1986, Ph. D. Thesis, State University of Ghent
12. DEBLAERE R., February 1986, Ph. D Thesis, Free University of Brussel, Belgium
13. VELTEN et al, *EMBO J.* 1984, vol. 3, No. 12, p. 2 7232-2 730
14. CHATER et al, *Gene cloning in Streptomyces. Curr. Top. Microbiol. Immunol.*, 1982, 96, p. 69-75
15. OMURA et al, *J. of Antibiotics*, Vol. 37, 8, 939-940, 1984
16. MURAKAMI et al, *Mol. Gen. Genet.*, 205, 42-50, 1986
17. MANDERSCHIED and WILD, *J. Plant Phys.*, 123, 135-142, 1986

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 21

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 183 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: Not Relevant
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (iii) HYPOTHETICAL: NO
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Streptomyces hygroscopicus
 - (B) STRAIN: ATCC 21705
- (ix) FEATURE:
 - (A) NAME/KEY: Region
 - (B) LOCATION: 1..2
 - (D) OTHER INFORMATION: /label= Xaa
/note= "Xaa is Val or Met"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Xaa Ser Pro Glu Arg Arg Pro Ala Asp Ile Arg Arg Ala Thr Glu Ala
1 5 10 15
Asp Met Pro Ala Val Cys Thr Ile Val Asn His Tyr Ile Glu Thr Ser
 20 25 30
Thr Val Asn Phe Arg Thr Glu Pro Gln Glu Pro Gln Glu Trp Thr Asp
 35 40 45
Asp Leu Val Arg Leu Arg Glu Arg Tyr Pro Trp Leu Val Ala Glu Val
50 55 60
Asp Gly Glu Val Ala Gly Ile Ala Tyr Ala Gly Pro Trp Lys Ala Arg
65 70 75 80
Asn Ala Tyr Asp Trp Thr Ala Glu Ser Thr Val Tyr Val Ser Pro Arg
 85 90 95
His Gln Arg Thr Gly Leu Gly Ser Thr Leu Tyr Thr His Leu Leu Lys
 100 105 110
Ser Leu Glu Ala Gln Gly Phe Lys Ser Val Val Ala Val Ile Gly Leu
 115 120 125
Pro Asn Asp Pro Ser Val Arg Met His Glu Ala Leu Gly Tyr Ala Pro
 130 135 140
Arg Gly Met Leu Arg Ala Ala Gly Phe Lys His Gly Asn Trp His Asp
145 150 155 160
Val Gly Phe Trp Gln Leu Asp Phe Ser Leu Pro Val Pro Pro Arg Pro
 165 170 175
Val Leu Pro Val Thr Glu Ile
 180

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 549 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO

-continued

20					25					30						
Thr	Val	Asn	Phe	Arg	Thr	Glu	Pro	Gln	Glu	Pro	Gln	Glu	Trp	Thr	Asp	
35					40					45						
Asp	Leu	Val	Arg	Leu	Arg	Glu	Arg	Tyr	Pro	Trp	Leu	Val	Ala	Glu	Val	
50					55					60						
Asp	Gly	Glu	Val	Ala	Gly	Ile	Ala	Tyr	Ala	Gly	Pro	Trp	Lys	Ala	Arg	
65					70					75					80	
Asn	Ala	Tyr	Asp	Trp	Thr	Ala	Glu	Ser	Thr	Val	Tyr	Val	Ser	Pro	Arg	
85					90					95						
His	Gln	Arg	Thr	Gly	Leu	Gly	Ser	Thr	Leu	Tyr	Thr	His	Leu	Leu	Lys	
100					105					110						
Ser	Leu	Glu	Ala	Gln	Gly	Phe	Lys	Ser	Val	Val	Ala	Val	Ile	Gly	Leu	
115					120					125						
Pro	Asn	Asp	Pro	Ser	Val	Arg	Met	His	Glu	Ala	Leu	Gly	Tyr	Ala	Pro	
130					135					140						
Arg	Gly	Met	Leu	Arg	Ala	Ala	Gly	Phe	Lys	His	Gly	Asn	Trp	His	Asp	
145					150					155					160	
Val	Gly	Phe	Trp	Gln	Leu	Asp	Phe	Ser	Leu	Pro	Val	Pro	Pro	Arg	Pro	
165					170					175						
Val	Leu	Pro	Val	Thr	Glu	Ile										
180																

- (2) INFORMATION FOR SEQ ID NO:4:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: Not Relevant
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: peptide
 - (iii) HYPOTHETICAL: NO
 - (v) FRAGMENT TYPE: N-terminal
 - (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Streptomyces hygroscopicus
 - (B) STRAIN: ATCC 21705
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Pro Glu Arg Arg Pro Ala Asp Ile Arg Arg
1 5 10

- (2) INFORMATION FOR SEQ ID NO:5:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "oligonucleotide 1"
 - (iii) HYPOTHETICAL: NO
 - (iv) ANTI-SENSE: NO
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CATGAGCCCA GAAC

- (2) INFORMATION FOR SEQ ID NO:6:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "oligonucleotide 2"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CGTCGTTCTG GGCT 14

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "DNA"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Cauliflower mosaic virus

(ix) FEATURE:
 (A) NAME/KEY: 5'UTR
 (B) LOCATION: 1..32

(ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION: 33..35
 (D) OTHER INFORMATION: /label= ATG
 /note= "translation initiation codon"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GAGGACACGC TGAAATCACC AGTCTCGGAT CCATG 35

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 44 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "DNA"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Cauliflower mosaic virus

(ix) FEATURE:
 (A) NAME/KEY: 5'UTR
 (B) LOCATION: 1..41

(ix) FEATURE:
 (A) NAME/KEY: -
 (B) LOCATION: 42..44
 (D) OTHER INFORMATION: /label= ATG
 /note= "translation initiation codon"

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GAGGACACGC TGAAATCACC AGTCTCTCTA CAAATCGATC CATG 44

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 40 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Cauliflower mosaic virus

(ix) FEATURE:

(A) NAME/KEY: 5'UTR

(B) LOCATION: 1..37

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION: 38..40

(D) OTHER INFORMATION: /label= ATG

/note= "translation initiation codon"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GAGGACACGC TGAAATCACC AGTCTCTCTA CAAATCGATG 40

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 618 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Streptomyces viridochromogenes

(B) STRAIN: DSM 40736

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 51..599

(D) OTHER INFORMATION: /product= "Phosphinothricin Acetyl Transferase"

/label= sfrv

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION: 51..53

(D) OTHER INFORMATION: /label= GTG

/note= "translation initiation codon in Streptomyces viridochromogenes"

(ix) FEATURE:

(A) NAME/KEY: -

(B) LOCATION: 600..602

(D) OTHER INFORMATION: /label= TGA

/note= "stop codon"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

TAAAGAGGTG CCCGCCACCC GCTTTCGCAG AACACCGAAG GAGACCACAC GTG AGC 56

-continued

															Val Ser			
															1			
CCA	GAA	CGA	CGC	CCG	GTC	GAG	ATC	CGT	CCC	GCC	ACC	GCC	GCC	GAC	ATG	104		
Pro	Glu	Arg	Arg	Pro	Val	Glu	Ile	Arg	Pro	Ala	Thr	Ala	Ala	Asp	Met			
5			10					15										
GCG	GCG	GTC	TGC	GAC	ATC	GTC	AAT	CAC	TAC	ATC	GAG	ACG	AGC	ACG	GTC	152		
Ala	Ala	Val	Cys	Asp	Ile	Val	Asn	His	Tyr	Ile	Glu	Thr	Ser	Thr	Val			
20			25					30										
AAC	TTC	CGT	ACG	GAG	CCG	CAG	ACT	CCG	CAG	GAG	TGG	ATC	GAC	GAC	CTG	200		
Asn	Phe	Arg	Thr	Glu	Pro	Gln	Thr	Pro	Gln	Glu	Trp	Ile	Asp	Asp	Leu			
35			40					45					50					
GAG	CGC	CTC	CAG	GAC	CGC	TAC	CCC	TGG	CTC	GTC	GCC	GAG	GTG	GAG	GGC	248		
Glu	Arg	Leu	Gln	Asp	Arg	Tyr	Pro	Trp	Leu	Val	Ala	Glu	Val	Glu	Gly			
				55					60					65				
GTC	GTC	GCC	GGC	ATC	GCC	TAC	GCC	GGC	CCC	TGG	AAG	GCC	CGC	AAC	GCC	296		
Val	Val	Ala	Gly	Ile	Ala	Tyr	Ala	Gly	Pro	Trp	Lys	Ala	Arg	Asn	Ala			
70			75					80										
TAC	GAC	TGG	ACC	GTC	GAG	TCG	ACG	GTG	TAC	GTC	TCC	CAC	CGG	CAC	CAG	344		
Tyr	Asp	Trp	Thr	Val	Glu	Ser	Thr	Val	Tyr	Val	Ser	His	Arg	His	Gln			
85			90					95										
CGG	CTC	GGA	CTG	GGC	TCC	ACC	CTC	TAC	ACC	CAC	CTG	CTG	AAG	TCC	ATG	392		
Arg	Leu	Gly	Leu	Gly	Ser	Thr	Leu	Tyr	Thr	His	Leu	Leu	Lys	Ser	Met			
100			105					110										
GAG	GCC	CAG	GGC	TTC	AAG	AGC	GTG	GTC	GCC	GTC	ATC	GGA	CTG	CCC	AAC	440		
Glu	Ala	Gln	Gly	Phe	Lys	Ser	Val	Val	Ala	Val	Ile	Gly	Leu	Pro	Asn			
115			120					125					130					
GAC	CCG	AGC	GTG	CGC	CTG	CAC	GAG	GCG	CTC	GGA	TAC	ACC	GCG	CGC	GGG	488		
Asp	Pro	Ser	Val	Arg	Leu	His	Glu	Ala	Leu	Gly	Tyr	Thr	Ala	Arg	Gly			
				135					140					145				
ACG	CTG	CGG	GCA	GCC	GGC	TAC	AAG	CAC	GGG	GGC	TGG	CAC	GAC	GTG	GGG	536		
Thr	Leu	Arg	Ala	Ala	Gly	Tyr	Lys	His	Gly	Gly	Trp	His	Asp	Val	Gly			
150			155					160										
TTC	TGG	CAG	CGC	GAC	TTC	GAG	CTG	CCG	GCC	CCG	CCC	CGC	CCC	GTC	CGG	584		
Phe	Trp	Gln	Arg	Asp	Phe	Glu	Leu	Pro	Ala	Pro	Pro	Arg	Pro	Val	Arg			
165			170					175										
CCC	GTC	ACA	CAG	ATC	TGAGCGGAGA GCGCATGGC										618			
Pro	Val	Thr	Gln	Ile														
180																		

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 183 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Val	Ser	Pro	Glu	Arg	Arg	Pro	Val	Glu	Ile	Arg	Pro	Ala	Thr	Ala	Ala	
1				5					10					15		
Asp	Met	Ala	Ala	Val	Cys	Asp	Ile	Val	Asn	His	Tyr	Ile	Glu	Thr	Ser	
20				25				30								
Thr	Val	Asn	Phe	Arg	Thr	Glu	Pro	Gln	Thr	Pro	Gln	Glu	Trp	Ile	Asp	
35				40				45								
Asp	Leu	Glu	Arg	Leu	Gln	Asp	Arg	Tyr	Pro	Trp	Leu	Val	Ala	Glu	Val	
50				55				60								
Glu	Gly	Val	Val	Ala	Gly	Ile	Ala	Tyr	Ala	Gly	Pro	Trp	Lys	Ala	Arg	
65				70				75				80				
Asn	Ala	Tyr	Asp	Trp	Thr	Val	Glu	Ser	Thr	Val	Tyr	Val	Ser	His	Arg	

-continued

85						90						95							
His	Gln	Arg	Leu	Gly	Leu	Gly	Ser	Thr	Leu	Tyr	Thr	His	Leu	Leu	Lys				
100						105						110							
Ser	Met	Glu	Ala	Gln	Gly	Phe	Lys	Ser	Val	Val	Ala	Val	Ile	Gly	Leu				
115						120						125							
Pro	Asn	Asp	Pro	Ser	Val	Arg	Leu	His	Glu	Ala	Leu	Gly	Tyr	Thr	Ala				
130						135						140							
Arg	Gly	Thr	Leu	Arg	Ala	Ala	Gly	Tyr	Lys	His	Gly	Gly	Trp	His	Asp				
145						150						155						160	
Val	Gly	Phe	Trp	Gln	Arg	Asp	Phe	Glu	Leu	Pro	Ala	Pro	Pro	Arg	Pro				
165						170						175							
Val	Arg	Pro	Val	Thr	Gln	Ile													
180																			

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 838 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Streptomyces hygroscopicus
 - (B) STRAIN: ATTC 21705

- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 182..199
 - (D) OTHER INFORMATION: /label= FokI
- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 713..718
 - (D) OTHER INFORMATION: /label= BglII

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CCCGCTCAAG CTCGCTGTCA TTTTCGAGAC GCCATCTTTG GAAGCGGTGG CCGAATCCGT	60
ACTGCGCGGA CTCGACGACG CGTAAAACGA TCGACCACGT ACACGAGTCC GGACACGGGG	120
CGAGGAGGCC CGGTTCCGGC ACCGAGGAAG ACCGAAGGAA GACCACACGT GAGCCCAGAA	180
CGACGCCCCG CCGACATCCG CCGTGCCACC GAGGCGGACA TGCCGGCGGT CTGCACCATC	240
GTCAACCACT ACATCGAGAC AAGCACGGTC AACTTCCGTA CCGAGCCGCA GGAACCGCAG	300
GAGTGGACGG ACGACCTCGT CCGTCTGCGG GAGCGCTATC CCTGGCTCGT CGCCGAGGTG	360
GACGGCGAGG TCGCCGGCAT CGCCTACGCG GGCCCCTGGA AGGCACGCAA CGCCTACGAC	420
TGGACGGCCG AGTCGACCGT GTACGTCTCC CCCCGCCACC AGCGGACGGG ACTGGGCTCC	480
ACGCTCTACA CCCACCTGCT GAAGTCCCTG GAGGCACAGG GCTTCAAGAG CGTGGTCGCT	540
GTCATCGGGC TGCCCAACGA CCCGAGCGTG CGCATGCACG AGGCGCTCGG ATATGCCCCC	600
CGCGGCATGC TGCGGGCGGC CGGCTTCAAG CACGGGAACT GGCATGACGT GGGTTTCTGG	660
CAGCTGGACT TCAGCCTGCC GGTACCGCCC CGTCCGGTCC TGCCCGTCAC CGAGATCTGA	720
ACGGAGTGCG CGTGGGCATC GCCCAGATTG GAGCTGGTAC GGGAATCAT CGAACTCAAC	780
TGGCATACCC GCAATGGTGA GGTGGAACCG CGGCGGATCG CGTACGACCG TGCCCAGG	838

-continued

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 56 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "adapter from plasmid pLK56.2"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 10..15
 - (D) OTHER INFORMATION: /label= BamHI

- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 14..19
 - (D) OTHER INFORMATION: /label= NcoI

- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 24..29
 - (D) OTHER INFORMATION: /label= BglII

- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 45..50
 - (D) OTHER INFORMATION: /label= BamHI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

AATTCCTGGG GATCCATGGT ACCAGATCTG ATCTCACGCG TCTAGGATCC GTCGAC 56

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "region around ATG translation initiation codon of sfr gene"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Streptomyces hygroscopicus
 - (B) STRAIN: ATCC 21705

- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 2..4
 - (D) OTHER INFORMATION: /label= ATG
/note= "translation initiation codon"

- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 5..14
 - (D) OTHER INFORMATION: /label= o11
/note= "region complementary to part of oligonucleotide 2"

- (ix) FEATURE:
 - (A) NAME/KEY: -
 - (B) LOCATION: 15..16
 - (D) OTHER INFORMATION: /label= FokI
/note= "splicing site of FokI"

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CATGAGCCCA GAACCGACGC CCG 23

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "DNA fragment in plasmid pGSR1"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 2..7
- (D) OTHER INFORMATION: /label= BglII

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 7..9
- (D) OTHER INFORMATION: /label= TGA
/note= "stop codon"

(ix) FEATURE:

- (A) NAME/KEY: -
- (B) LOCATION: 23..28
- (D) OTHER INFORMATION: /label= BamHI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GAGATCTGAT CTCACGCGTC TAGGATCC 28

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "oligonucleotide 3"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CATGGACCCA GAAC 14

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "oligonucleotide 4"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

-continued

CGTCGTTCTG GGTC

14

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 183 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: Not Relevant

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Streptomyces hygroscopicus

(B) STRAIN: ATCC 21705

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Met Ser Pro Glu Arg Arg Pro Ala Asp Ile Arg Arg Ala Thr Glu Ala

1 5 10 15

Asp Met Pro Ala Val Cys Thr Ile Val Asn His Tyr Ile Glu Thr Ser

20 25 30

Thr Val Asn Phe Arg Thr Glu Pro Gln Glu Pro Gln Glu Trp Thr Asp

35 40 45

Asp Leu Val Arg Leu Arg Glu Arg Tyr Pro Trp Leu Val Ala Glu Val

50 55 60

Asp Gly Glu Val Ala Gly Ile Ala Tyr Ala Gly Pro Trp Lys Ala Arg

65 70 75 80

Asn Ala Tyr Asp Trp Thr Ala Glu Ser Thr Val Tyr Val Ser Pro Arg

85 90 95

His Gln Arg Thr Gly Leu Gly Ser Thr Leu Tyr Thr His Leu Leu Lys

100 105 110

Ser Leu Glu Ala Gln Gly Phe Lys Ser Val Val Ala Val Ile Gly Leu

115 120 125

Pro Asn Asp Pro Ser Val Arg Met His Glu Ala Leu Gly Tyr Ala Pro

130 135 140

Arg Gly Met Leu Arg Ala Ala Gly Phe Lys His Gly Asn Trp His Asp

145 150 155 160

Val Gly Phe Trp Gln Leu Asp Phe Ser Leu Pro Val Pro Pro Arg Pro

165 170 175

Val Leu Pro Val Thr Glu Ile

180

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "region around GTG translation

initiation codon of sfrv gene in plasmid pJS1"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Streptomyces viridochromogenes

(B) STRAIN: DSM 40736

(ix) FEATURE:

-continued

(A) NAME/KEY: -
(B) LOCATION: 1..3
(D) OTHER INFORMATION: /label= GTG
/note= "translation initiation codon"

(ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION: 3..8
(D) OTHER INFORMATION: /label= BanII

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GTGAGCCCAG AAG 13

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "region around translation initiation
codon of sfrv gene in plasmid pGSR1SV"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION: 1..6
(D) OTHER INFORMATION: /label= BamHI

(ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION: 7..9
(D) OTHER INFORMATION: /label= ATG
/note= "translation initiation codon"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GGATCCATGA GC 12

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "region around stop codon of sfrv
gene in plasmid pGSR1SV"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION: 7..9
(D) OTHER INFORMATION: /label= TGA
/note= "stop codon"

(ix) FEATURE:
(A) NAME/KEY: -
(B) LOCATION: 23..28
(D) OTHER INFORMATION: /label= BamHI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GAGATCTGAT CTCACGCGTC TAGGATCC 28

51

The invention claimed is:

1. An isolated DNA encoding a protein having phosphino-thricin acetyltransferase activity, or a variant thereof retaining said activity, said protein comprising the amino acid sequence (SEQ ID No. 1):

X Ser Pro Glu Arg Arg Pro Ala Asp
Ile Arg Arg Ala Thr Glu Ala Asp MET
Pro Ala Val Cys Thr Ile Val Asn His
Tyr Ile Glu Thr Ser Thr Val Asn Phe
Arg Thr Glu Pro Gln Glu Pro Gln Glu
Trp Thr Asp Asp Leu Val Arg Leu Arg
Glu Arg Tyr Pro Trp Leu Val Ala Glu
Val Asp Gly Glu Val Ala Gly Ile Ala
Tyr Ala Gly Pro Trp Lys Ala Arg Asn
Ala Tyr Asp Trp Thr Ala Glu Ser Thr
Val Tyr Val Ser Pro Arg His Gln Arg
Thr Gly Leu Gly Ser Thr Leu Tyr Thr
His Leu Leu Lys Ser Leu Glu Ala Gln
Gly Phe Lys Ser Val Val Ala Val Ile
Gly Leu Pro Asn Asp Pro Ser Val Arg
Met His Glu Ala Leu Gly Tyr Ala Pro
Arg Gly Met Leu Arg Ala Ala Gly Phe
Lys His Gly Asn Trp His Asp Val Gly
Phe Trp Gln Leu Asp Phe Ser Leu Pro
Val Pro Pro Arg Pro Val Leu Pro Val
Thr Glu Ile

in which X is Met [or Val], said DNA consisting of between 549 and 625 nucleotides wherein X is encoded by ATG.

52

2. The isolated DNA of claim 1, consisting of the nucleotide sequence (SEQ ID No. 2 from nucleotide 2 to nucleotide 549):

5 NTG AGC CCA GAA CGA CGC CCG GCC GAC
ATC CGC CGT GCC ACC GAG GCG GAC ATG
10 CCG GCG GTC TGC ACC ATC GTC AAC CAC
TAC ATC GAG ACA AGC ACG GTC AAC TTC
CGT ACC GAG CCG CAG GAA CCG CAG GAG
15 TGG ACG GAC GAC CTC GTC CGT CTG CGG
GAG CGC TAT CCC TGG CTC GTC GCC GAG
GTG GAC GGC GAG GTC GCC GGC ATC GCC
TAC GCG GGC CCC TGG AAG GCA CGC AAC
20 GCC TAC GAC TGG ACG GCC GAG TCG ACC
GTG TAC GTC TCC CCC CGC CAC CAG CGG
ACG GGA CTG GGC TCC ACG CTC TAC ACC
25 CAC CTG CTG AAG TCC CTG GAG GCA CAG
GGC TTC AAG AGC GTG GTC GCT GTC ATC
GGG CTG CCC AAC GAC CCG AGC GTG CGC
30 ATG CAC GAG GCG CTC GGA TAT GCC CCC
CGC GGC ATG CTG CGG GCG GCC GGC TTC
AAG CAC GGG AAC TGG CAT GAC GTG GGT
35 TTC TGG CAG CTG GAC TTC AGC CTG CCG
GTA CCG CCC CGT CCG GTC CTG CCC GTC
ACC GAG ATC

in which N is A [or G].

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : RE44,962 C1
APPLICATION NO. : 90/013394
DATED : February 12, 2021
INVENTOR(S) : Leemans et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page

Item (73) Assignees:

After, BAYERCROPSCIENCE N.V., DIEGEM (BE)

Add --BIOGEN IDEC MA, INC., CAMBRIDGE, MA (US)--

Signed and Sealed this
Twenty-third Day of March, 2021



Drew Hirshfeld
*Performing the Functions and Duties of the
Under Secretary of Commerce for Intellectual Property and
Director of the United States Patent and Trademark Office*

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : RE44,962 C1
APPLICATION NO. : 90/013515
DATED : February 12, 2021
INVENTOR(S) : Leemans et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page

(73) Assignees:

After:

BAYERCROPSCIENCE N.V., DIEGEM (BE)

Add:

--BIOGEN IDEC MA, INC., CAMBRIDGE, MA (US)--

Signed and Sealed this
Sixth Day of April, 2021



Drew Hirshfeld
*Performing the Functions and Duties of the
Under Secretary of Commerce for Intellectual Property and
Director of the United States Patent and Trademark Office*

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : RE44,962 C1
APPLICATION NO. : 90/013563
DATED : February 12, 2021
INVENTOR(S) : Leemans et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page

(73) Assignees:

After:

BAYERCROPSCIENCE N.V., DIEGEM (BE)

Add:

--BIOGEN IDEC MA, INC., CAMBRIDGE, MA (US)--

Signed and Sealed this
Thirteenth Day of April, 2021



Drew Hirshfeld
*Performing the Functions and Duties of the
Under Secretary of Commerce for Intellectual Property and
Director of the United States Patent and Trademark Office*



US00RE44962C1

(12) **EX PARTE REEXAMINATION CERTIFICATE** (11803rd)
United States Patent
Leemans et al.

(10) **Number:** **US RE44,962 C1**(45) **Certificate Issued:** **Feb. 12, 2021**

(54) **GENETICALLY ENGINEERED PLANT CELLS AND PLANTS EXHIBITING RESISTANCE TO GLUTAMINE SYNTHETASE INHIBITORS, DNA FRAGMENTS AND RECOMBINANTS FOR USE IN THE PRODUCTION OF SAID CELLS AND PLANTS**

Nov. 5, 1987, now abandoned, which is a continuation of application No. PCT/EP87/00141, filed on Mar. 11, 1987.

(30) **Foreign Application Priority Data**

Mar. 11, 1986 (GB) 86400521.0
Jan. 21, 1987 (GB) 87400141.5

(71) Applicants: **Bayer Cropscience N.V.**, Diegem (BE); **Biogen Idec MA, Inc.**, Cambridge, MA (US)

(51) **Int. Cl.**
C12N 15/52 (2006.01)
C12N 15/31 (2006.01)
A01H 1/00 (2006.01)
C12N 9/10 (2006.01)
C12N 15/63 (2006.01)
C12N 15/70 (2006.01)
C12N 15/76 (2006.01)
C12N 15/82 (2006.01)

(72) Inventors: **Jan Leemans**, Heusden (BE); **Johan Botterman**, Zwijnaarde (BE); **Marc De Block**, Gentbrugge (BE); **Charles Thompson**, Grand Lancy/Geneva (CH); **Rao Mouva**, Geneva (CH)

(52) **U.S. Cl.**
CPC **A01H 1/00** (2013.01); **C12N 9/1029** (2013.01); **C12N 15/63** (2013.01); **C12N 15/70** (2013.01); **C12N 15/76** (2013.01); **C12N 15/8277** (2013.01); **C07K 2319/00** (2013.01); **Y10S 47/01** (2013.01)

(73) Assignee: **BAYER CROPSCIENCE N.V.**, Diegem (BE)

(58) **Field of Classification Search**

None
See application file for complete search history.

Reexamination Request:

No. 90/013,394, Nov. 7, 2014
No. 90/013,515, Jun. 1, 2015
No. 90/013,563, Aug. 7, 2015

Reexamination Certificate for:

Patent No.: **Re. 44,962**
Issued: **Jun. 24, 2014**
Appl. No.: **14/017,108**
Filed: **Sep. 3, 2013**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **7,112,665**
Issued: **Sep. 26, 2006**
Appl. No.: **08/465,219**
Filed: **Jun. 5, 1995**

Related U.S. Application Data

(60) Division of application No. 07/525,300, filed on May 17, 1990, now Pat. No. 5,561,236, which is a continuation of application No. 07/131,140, filed on

(56) **References Cited**

To view the complete listing of prior art documents cited during the proceedings for Reexamination Control Numbers 90/013,394, 90/013,515 and 90/013,563, please refer to the USPTO's public Patent Application Information Retrieval (PAIR) system under the Display References tab.

Primary Examiner — Padmashri Ponnaluri

(57) **ABSTRACT**

The invention relates to a DNA fragment containing a determined gene, the expression of which inhibits the anti-biotic and herbicidal effects of Bialaphos and related products. It also relates to recombinant vectors, containing such DNA fragment, which enable this protective gene to be introduced and expressed into cells and plant cells.

1
EX PARTE
REEXAMINATION CERTIFICATE

2

NO AMENDMENTS HAVE BEEN MADE TO
THE PATENT 5

AS A RESULT OF REEXAMINATION, IT HAS BEEN
DETERMINED THAT:

The patentability of claims **1** and **2** is confirmed. 10

* * * * *