

US00RE46042E

(19) **United States**
(12) **Reissued Patent**
Whitehead et al.

(10) **Patent Number: US RE46,042 E**
(45) **Date of Reissued Patent: Jun. 28, 2016**

- (54) **DEVELOPMENT OF DENGUE VIRUS VACCINE COMPONENTS**
- (71) Applicant: **The Government of The United States of America, as represented by the Secretary, Department of Health, Rockville, MD (US)**
- (72) Inventors: **Stephen S. Whitehead**, Montgomery Village, MD (US); **Joseph E. Blaney**, Gettysburg, PA (US); **Brian R. Murphy**, Bethesda, MD (US); **Ching-Juh Lai**, Bethesda, MD (US)
- (73) Assignee: **The United States of America, as represented by the Secretary, Department of Health & Human Services, Rockville, MD (US)**
- (21) Appl. No.: **13/895,908**
- (22) Filed: **May 16, 2013**

Related U.S. Patent Documents

Reissue of:

- (64) Patent No.: **8,337,860**
- Issued: **Dec. 25, 2012**
- Appl. No.: **12/376,756**
- PCT Filed: **Aug. 15, 2007**
- PCT No.: **PCT/US2007/076004**
- § 371 (c)(1),
- (2) Date: **Dec. 17, 2009**
- PCT Pub. No.: **WO2008/022196**
- PCT Pub. Date: **Feb. 21, 2008**

U.S. Applications:

- (60) Provisional application No. 60/837,723, filed on Aug. 15, 2006.

- (51) **Int. Cl.**
A61K 39/12 (2006.01)
C12N 15/40 (2006.01)
C12N 7/04 (2006.01)
C12N 7/01 (2006.01)
A61K 48/00 (2006.01)
A61K 39/295 (2006.01)
C12N 15/83 (2006.01)
C07K 14/005 (2006.01)
C12N 7/00 (2006.01)
C07K 14/18 (2006.01)
A61K 39/00 (2006.01)

- (52) **U.S. Cl.**
CPC *C12N 7/045* (2013.01); *C07K 14/005* (2013.01); *C07K 14/1825* (2013.01); *C12N 7/00* (2013.01); *A61K 2039/5254* (2013.01); *C12N 2770/24122* (2013.01); *C12N 2770/24162* (2013.01)

- (58) **Field of Classification Search**
CPC *C12N 2770/24134*; *C12N 2770/24161*; *C12N 7/00*; *C12N 2770/24122*; *C12N 2770/24162*; *C12N 7/045*; *A61K 2039/5254*; *A61K 2039/5256*; *A61K 39/12*; *C07K 14/005*; *C07K 14/1825*
USPC 424/202.1
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,184,024	B1 *	2/2001	Lai et al.	435/235.1
7,517,531	B2 *	4/2009	Whitehead et al.	424/202.1
8,075,903	B2 *	12/2011	Whitehead et al.	424/202.1

FOREIGN PATENT DOCUMENTS

WO		03/092592	A2	11/2003
WO		WO 03/092592	A2	11/2003

OTHER PUBLICATIONS

Zhou et al., "Comparative analysis reveals no consistent association between the secondary structure of the 3'-untranslated region of dengue viruses and disease syndrome," *Journal of General Virology*, 87:2595-2603 (2006).

Proutski et al., "Secondary structure of the 3' untranslated region of flaviviruses: similarities and differences," *Nucleic Acids Research*, 25(6):1194-1202 (1997).

Blaney et al., "Dengue virus type 3 vaccine candidates generated by introduction of deletions in the 3' untranslated region (3'-UTR) or by exchange of the DENV-3 3'-UTR with that of DENV-4," *Vaccine*, 26:817-828 (2008).

Alvarez et al., "Role of RNA structures present at the 3'UTR of dengue virus on translation, RNA synthesis, and viral replication," *Virology*, 339:200-212 (2005).

Blaney Jr., et al. "Genetically Modified, Live Attenuated Dengue Virus Type 3 Vaccine Candidates" *Am. J. Trop. Med. Hyg.*, 71:811-821 (2004).

Blaney Jr., et al. "Development of a Live Attenuated Dengue Virus Vaccine Using Reverse Genetics" *Viral Immunology*, 19(1):10-32 (2006).

Durbin et al., "rDEN2/4E30(ME), A Live Attenuated Chimeric Dengue Serotype 2 Vaccine Is Safe and Highly Immunogenic in Health Dengue-Naive Adults" *Human Vaccines*, 2(6):255-260 (2006).

Men et al., "Dengue Type 4 Virus Mutants Containing Deletions in the 3' Noncoding Region of the RNA Genome: Analysis of Growth Restriction in Cell Culture and Altered Viremia Pattern and Immunogenicity in Rhesus Monkeys" *Journal of Virology*, 70(5):3930-3937 (1996).

Whitehead et al. "Substitution of the structural genes of dengue virus type 4 with those of type 2 results in chimeric vaccine candidates which are attenuated for mosquitoes, mice, and rhesus monkeys" *Vaccine*, 21:4307-4316 (2003).

International Search Report dated Jan. 13, 2009 for PCT/US20071076004.

* cited by examiner


Primary Examiner — Shri Ponnaluri
(74) *Attorney, Agent, or Firm* — Locke Lorde LLP; Bryan D. Zerhusen; Gabriel J. McCool

(57) **ABSTRACT**

The invention is related to a dengue virus or chimeric dengue virus that contains a mutation in the 3' untranslated region (3'-UTR) comprising a Δ30 mutation that removes the TL-2 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4, and nucleotides additional to the Δ30 mutation deleted from the 3'-UTR that removes sequence in the 5' direction as far as the 5' boundary of the TL-3 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4, or a replacement of the 3'-UTR of a dengue virus of a first serotype with the 3'-UTR of a dengue virus of a second serotype, optionally containing the Δ30 mutation and nucleotides additional to the Δ30 mutation deleted from the 3'-UTR; and immunogenic compositions, methods of inducing an immune response, and methods of producing a dengue virus or chimeric dengue virus.

14 Claims, 22 Drawing Sheets

A. Delete additional nucleotides from the 3' UTR:



B. Replace the 3' UTR of DEN3 with the UTR from DEN4 $\Delta 30$:




Figure 1




Figure 1a




Figure 1b




Figure 1c





Figure 2



DEN2 wt (Tonga/74)
GenBank: AY744147
 $\Delta G = -88.3$
(mfold v3.2: 1 0 10, P 79 0 14, P 178 0 10)

Figure 3





Figure 4



DEN4 wt (814669)
GenBank: AF326573
 $\Delta G = -98.8$
(mfold v3.2: 1 0 4, P 82 0 15, P 168 0 20)

Figure 5





Figure 6



DEN2 Δ 30
 $\Delta G = -76.0$
(mfold v3.2: 1 0 10, P 79 0 14, P 148 0 10)

Figure 7





Figure 8



DEN4 Δ 30
 $\Delta G = -80.4$
(mfold v3.2: 1 0 4, P 82 0 15, P 137 0 20)

Figure 9





Figure 10



DEN2Δ30/31
 $\Delta G = -65.7$
(mfold v3.2: 1 0 10, P 48 0 14, P 117 0 10)

Figure 11




Figure 12




Figure 13





Figure 14



DEN2Δ86

$\Delta G = -58.6$

(mfold v3.2: 1 0 10, P 93 0 10)

Figure 15





Figure 16



DEN4Δ86
 $\Delta G = -61.8$
(mfold v3.2: 1 0 4, P 82 0 20)

Figure 17




Figure 18




Figure 19

DEVELOPMENT OF DENGUE VIRUS VACCINE COMPONENTS

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue; a claim printed with strikethrough indicates that the claim was canceled, disclaimed, or held invalid by a prior post-patent action or proceeding.

CROSS REFERENCE TO RELATED APPLICATION

This application claims the benefit and priority to and is a U.S. National Phase of International Application Number PCT/US2007/076004, filed on Aug. 15, 2007, designating the United States of America and published in the English language, which is an International Application of and claims the benefit of priority to U.S. Provisional Patent Application No. 60/837,723, filed Aug. 15, 2006. The disclosures of the above-referenced applications are hereby expressly incorporated by reference in their entireties.

FIELD OF THE INVENTION

The invention relates to mutations in the 3' untranslated region of the genome of dengue virus serotypes 1, 2, 3, and 4 that are useful in attenuating the growth characteristics of dengue virus vaccines.

DESCRIPTION OF THE RELATED ART

There are four serotypes of dengue virus (dengue virus type 1 [DEN1], DEN2, DEN3, and DEN4) that annually cause an estimated 50 to 100 million cases of dengue fever and 500,000 cases of the more severe form of dengue virus infection known as dengue hemorrhagic fever/dengue shock syndrome (Gubler, D. J. and M. Meltzer 1999 *Adv Virus Res* 53:35-70). Dengue virus is widely distributed throughout the tropical and semitropical regions of the world, and the number of dengue virus infections continues to increase due to the expanding range of its *Aedes aegypti* mosquito vector. A vaccine is not available for the control of dengue disease despite its importance as a reemerging disease. The goal of immunization is to protect against dengue virus disease by the induction of a long-lived neutralizing antibody response against each of the four serotypes. Simultaneous protection against all four serotypes is required, since an increase in disease severity can occur in persons with preexisting antibodies to a heterotypic dengue virus. Such immunization can be achieved economically with a live, attenuated virus vaccine.

Dengue viruses are positive-sense RNA viruses belonging to the Flavivirus genus. The approximately 11,000-base genome contains a single open reading frame encoding a polyprotein which is processed by proteases of both viral and cellular origin into three structural proteins (C, prM, and E) and at least seven nonstructural (NS) proteins. Both ends of the dengue virus genome contain an untranslated region (UTR), and the overall genome organization is 5'-UTR-CprM-E-NS1-NS2A-NS2B-NS3-NS4A-NS4B-NS5-UTR-3'. The 3' UTR is nearly 400 bases in length and is predicted to contain several stem-loop structures conserved among dengue virus serotypes (Brinton, M. A. et al. 1986 *Virology* 153:113-121, Hahn, C. S. et al. 1987 *J Mol Biol* 198:33-41, Proutski, V. et al. 1997 *Nucleic Acids Res* 25:1194-1202,

Rauscher, S. et al. 1997 *RNA* 3:779-791, Shurtleff, A. et al. 2001 *Virology* 281:75-87). One such stem-loop structure, identified as TL-2 in the proposed secondary structure of the 3' UTR (Proutski, V. et al. 1997 *Nucleic Acids Res* 25:1194-1202), was previously removed by deletion of 30 nucleotides from the DEN4 genome (3' nucleotides 172 to 143) (Men, R. et al. 1996 *J Virol* 70:3930-3937) and has subsequently been designated as the $\Delta 30$ mutation (Durbin, A. P. et al. 2001 *Am J Trop Med Hyg* 65:405-413). The resulting virus, rDEN4 $\Delta 30$, was shown to be attenuated in rhesus monkeys compared to parental viruses containing an intact TL-2 sequence and is attenuated in humans (Durbin, A. P. et al. 2001 *Am J Trop Med Hyg* 65:405-413).

SUMMARY OF THE INVENTION

The invention is related to a dengue virus or chimeric dengue virus comprising a mutation in the 3' untranslated region (3'-UTR) selected from the group consisting of:

(a) a $\Delta 30$ mutation that removes the TL-2 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4, and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR that removes sequence in the 5' direction as far as the 5' boundary of the TL-3 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4; and

(b) a replacement of the 3'-UTR of a dengue virus of a first serotype with the 3'-UTR of a dengue virus of a second serotype, optionally containing the $\Delta 30$ mutation and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR;

and immunogenic compositions, methods of inducing an immune response, and methods of producing a dengue virus or chimeric dengue virus.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Two approaches to attenuate dengue viruses. A) (a-c) Deletion of additional nucleotides from the 3'-UTR (DEN3 wt Sleman/78, SEQ ID NO: 1). B) Replacement of the 3'-UTR of a dengue virus of a first serotype with the 3'-UTR of a dengue virus of a second serotype.

FIG. 2. Predicted secondary structure of the TL-1, TL-2 and TL-3 region of the 3'-UTR of DEN1 serotype virus. The GenBank accession number of the sequence used for construction of the secondary structure model is indicated. Only the last 278 nucleotides which comprise TL-1, TL-2 and TL-3, are used to avoid circularization of the structure and subsequent misfolding of known and experimentally-verified structural elements. The mfold program constraints specific for each structure model are indicated. Nucleotides that border the principle deletions are circled and numbered, with nucleotide numbering beginning at the 3' genome end (reverse-direction numbering system). SEQ ID NO: 2.

FIG. 3. Predicted secondary structure of the TL-1, TL-2 and TL-3 region of the 3'-UTR of DEN2 serotype virus. The GenBank accession number of the sequence used for construction of the secondary structure model is indicated. Only the last 281 nucleotides which comprise TL-1, TL-2 and TL-3, are used to avoid circularization of the structure and subsequent misfolding of known and experimentally-verified structural elements. The mfold program constraints specific for each structure model are indicated. Nucleotides that border the principle deletions are circled and numbered, with nucleotide numbering beginning at the 3' genome end (reverse-direction numbering system). SEQ ID NO: 3.

FIG. 4. Predicted secondary structure of the TL-1, TL-2 and TL-3 region of the 3'-UTR of DEN3 serotype virus. The GenBank accession number of the sequence used for construction of the secondary structure model is indicated. Only the last 276 nucleotides which comprise TL-1, TL-2 and TL-3, are used to avoid circularization of the structure and subsequent misfolding of known and experimentally-verified structural elements. The mfold program constraints specific for each structure model are indicated. Nucleotides that border the principle deletions are circled and numbered, with nucleotide numbering beginning at the 3' genome end (reverse-direction numbering system). SEQ ID NO: 4.

FIG. 5. Predicted secondary structure of the TL-1, TL-2 and TL-3 region of the 3'-UTR of DEN4 serotype virus. The GenBank accession number of the sequence used for construction of the secondary structure model is indicated. Only the last 281 nucleotides which comprise TL-1, TL-2 and TL-3, are used to avoid circularization of the structure and subsequent misfolding of known and experimentally-verified structural elements. The mfold program constraints specific for each structure model are indicated. Nucleotides that border the principle deletions are circled and numbered, with nucleotide numbering beginning at the 3' genome end (reverse-direction numbering system). SEQ ID NO: 5.

FIG. 6. $\Delta 30$ deletion mutation depicted for DEN1. The $\Delta 30$ mutation deletes nt 174 to 145 of DEN1, with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 6.

FIG. 7. $\Delta 30$ deletion mutation depicted for DEN2. The $\Delta 30$ mutation deletes nt 173 to 144 of DEN2, with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 7.

FIG. 8. $\Delta 30$ deletion mutation depicted for DEN3. The $\Delta 30$ mutation deletes nt 173 to 143 of DEN3, with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 8.

FIG. 9. $\Delta 30$ deletion mutation depicted for DEN4. The $\Delta 30$ mutation deletes nt 172 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 9.

FIG. 10. $\Delta 30/31$ deletion mutation depicted for DEN1. In addition to the deletion of the nucleotides comprising the $\Delta 30$ mutation, the $\Delta 31$ mutation deletes nt 258 to 228 of DEN1 with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 10.

FIG. 11. $\Delta 30/31$ deletion mutation depicted for DEN2. In addition to the deletion of the nucleotides comprising the $\Delta 30$ mutation, the $\Delta 31$ mutation deletes nt 258 to 228 of DEN2 with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 11.

FIG. 12. $\Delta 30/31$ deletion mutation depicted for DEN3. In addition to the deletion of the nucleotides comprising the $\Delta 30$ mutation, the $\Delta 31$ mutation deletes nt 258 to 228 of DEN3 with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 12.

FIG. 13. $\Delta 30/31$ deletion mutation depicted for DEN4. In addition to the deletion of the nucleotides comprising the $\Delta 30$ mutation, the $\Delta 31$ mutation deletes nt 258 to 228 of DEN4 with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 13.

FIG. 14. $\Delta 86$ deletion mutation depicted for DEN1. The $\Delta 86$ mutation deletes nt 228 to 145 of DEN1 with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 14.

FIG. 15. $\Delta 86$ deletion mutation depicted for DEN2. The $\Delta 86$ mutation deletes nt 228 to 144 of DEN2 with reverse-

direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 15.

FIG. 16. $\Delta 86$ deletion mutation depicted for DEN3. The $\Delta 86$ mutation deletes nt 228 to 143 of DEN3 with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 16.

FIG. 17. $\Delta 86$ deletion mutation depicted for DEN4. The $\Delta 86$ mutation deletes nt 228 to 143 of DEN4 with reverse-direction numbering system. The deleted region is indicated by the Δ symbol. SEQ ID NO: 17.

FIG. 18. Chimerization of rDEN3 with the rDEN4 or rDEN4 $\Delta 30$ 3'-UTR. A) Recombinant 3'-UTR chimeric dengue viruses were constructed by replacing the 3'-UTR of rDEN3 with regions derived from either rDEN4 or rDEN4 $\Delta 30$. The relative location of the $\Delta 30$ mutation in the 3'-UTR is indicated by an arrow. The junctions between the ORF and UTR for rDEN3 and rDEN4 are indicated as junctions 1 and 2, respectively. Intertypic junction 3 is also indicated for the resulting chimeric viruses. B) Nucleotide and amino acid sequence of the junction regions are shown. For junction 3, nucleotide substitutions used to introduce a unique HpaI restriction enzyme recognition site are shown in lower case. Junction 1—SEQ ID NOs: 18 (nucleotide) and 19 (amino acid); Junction 2—SEQ ID NOs: 20 (nucleotide) and 21 (amino acid); Junction 3—SEQ ID NOs: 22 (nucleotide) and 23 (amino acid).

FIG. 19. Replication in Vero cells and C6/36 cells. Four mutant viruses were compared to wild type rDEN3 for replication in Vero cells and C6/36 cells. 75 cm² flasks of confluent cells were infected at a multiplicity of infection of 0.01. Aliquots of 0.5 ml were removed from flasks daily for seven days. After addition of SPG to a concentration of 1 \times , samples were frozen on dry ice and stored at -80 $^{\circ}$ C. Virus titer was determined by plaque assay on Vero cells for all samples. The limit of detection is 1.0 log₁₀ PFU/ml.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

Definitions

Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. See, e.g., Singleton P and Sainsbury D., Dictionary of Microbiology and Molecular Biology 3rd ed., J. Wiley & Sons, Chichester, N.Y., 2001, and Fields Virology 4th ed., Knipe D. M. and Howley P. M. eds, Lippincott Williams & Wilkins, Philadelphia 2001.

The term "about" means within 1, 2, or 3 nucleotides. Mutant Dengue Viruses and Chimeric Dengue Viruses

A goal of the invention is to develop a set of type-specific, live attenuated dengue vaccine components that can be formulated into a safe, effective, and economical tetravalent dengue vaccine. The $\Delta 30$ mutation attenuates DEN4 in rhesus monkeys (Men, R. et al. 1996 J Virol 70:3930-3937)). The $\Delta 30$ mutation removes a homologous structure (TL-2) in each of the dengue virus serotypes 1, 2, 3, and 4 (FIGS. 2-5). However, the $\Delta 30$ mutation was found to not attenuate DEN3 in rhesus monkeys.

An embodiment of the invention provides dengue viruses and chimeric dengue viruses having one or more mutations that result in attenuation, methods of making such dengue viruses, and methods for using these dengue viruses to prevent or treat dengue virus infection. The mutation (or mutations) in the dengue virus of the invention is present in the 3' untranslated region (3'-UTR) formed by the most down-

stream approximately 384 nucleotides of the viral RNA, which have been shown to play a role in determining attenuation. The viruses and methods of the invention are described further, as follows.

One example of a dengue virus that can be used in the invention is the serotype 3, Sleman/78 strain. The applicability of the invention to all members of the dengue virus taxonomic group is inferred by the observation that the properties of other dengue virus strains are similar to that of any one dengue virus strain. Dengue viruses have been grouped into four serotypes (DEN1, DEN2, DEN3 and DEN4). Numerous strains have been identified for each of the four serotypes. The complete genomic sequences of various dengue virus strains are provided as Genbank accession numbers in Table A.

TABLE A

Examples of Dengue Virus Strains		
Serotype	Strain	Accession No.
1	02-20	AB178040
1	16007	AF180817
1	16007 PDK-13	AF180818
1	259par00	AF514883
1	280par00	AF514878
1	293arg00	AY206457
1	295arg00	AF514885
1	297arg00	AF514889
1	301arg00	AF514876
1	98901518	AB189120
1	98901530	AB189121
1	A88	AB074761
1	Abidjan	AF298807
1	ARG0028	AY277665
1	ARG0048	AY277666
1	ARG9920	AY277664
1	BR-90	AF226685
1	BR-01-MR	AF513110
1	BR-97-111	AF311956
1	BR-97-233	AF311958
1	BR-97-409	AF311957
1	Cambodia	AF309641
1	FGA-89	AF226687
1	FGA-NA d1d	AF226686
1	Fj231 -04	DQ193572
1	GD05-99	AY376738
1	GD23-95	AY373427
1	GZ-80	AF350498
1	D1-hu-Yap-NIID27-2004	AB204803
1	D1-H-IMTSSA-98-606	AF298808
1	Mochizuki	AB074760
1	D1.Myanmar.059-01	AY708047
1	D1.Myanmar.194-01	AY713474
1	D1.Myanmar.206-01	AY713475
1	D1.Myanmar.23819-96	AY722802
1	D1.Myanmar.305-01	AY713476
1	D1.Myanmar.31459-98	AY726555
1	D1.Myanmar.31987-98	AY726554
1	D1.Myanmar.32514-98	AY722803
1	D1.Myanmar.37726-01	AY726549
1	D1.Myanmar.38862-01	AY726550
1	D1.Myanmar.40553-71	AY713473
1	D1.Myanmar.40568-76	AY722801
1	D1.Myanmar.44168-01	AY726551
1	D1.Myanmar.44988-02	AY726552
1	D1.Myanmar.49440-02	AY726553
1	rWestern Pacific-delta30	AY145123
1	Western Pacific rDEN1mutF	AY145122
1	S275-90	A75711
1	D1-hu-Seychelles-NIID41-2003	AB195673
1	Singapore 8114-93	AY762084
1	Singapore S275-90	M87512
1	ThD1_0008_81	AY732483
1	ThD1_0049_01	AY732482
1	ThD1_0081_82	AY732481
1	ThD1_0097_94	AY732480
1	ThD1_0102_01	AY732479

TABLE A-continued

Examples of Dengue Virus Strains			
Serotype	Strain	Accession No.	
1	ThD1_0323_91	AY732478	
1	ThD1_0336_91	AY732477	
1	ThD1_0442_80	AY732476	
1	ThD1_0488_94	AY732475	
1	ThD1_0673_80	AY732474	
10	Recombinant Western Pacific	AY145121	
1	Nauru Island Western Pacific 45AZ5	NC_001477	
1	Nauru Island, Western Pacific Bethesda	U88535	
1	Nauru Island Western Pacific 45AZ5-PDK27	U88537	
2	131	AF100469	
2	16681-PDK53	M84728	
2	16681 Blok	M84727	
15	16681 Kinney	U87411	
2	43	AF204178	
2	44	AF204177	
2	98900663	AB189122	
2	98900665	AB189123	
2	98900666	AB189124	
20	2	BA05i	AY858035
2	Bangkok 1974	AJ487271	
2	BR64022	AF489932	
2	C0166	AF100463	
2	C0167	AF100464	
2	C0371	AF100461	
25	2	C0390	AF100462
2	China 04	AF119661	
2	Cuba115-97	AY702036	
2	Cuba13-97	AY702034	
2	Cuba165-97	AY702038	
2	Cuba205-97	AY702039	
2	Cuba58-97	AY702035	
30	2	Cuba89-97	AY702037
2	DR23-01	AB122020	
2	DR31-01	AB122021	
2	DR59-01	AB122022	
2	FJ-10	AF276619	
35	2	FJ11-99	AF359579
2	I348600	AY702040	
2	IQT1797	AF100467	
2	IQT2913	AF100468	
2	Jamaica-N 1409	M20558	
2	K0008	AF100459	
2	K0010	AF100460	
40	2	Mara4	AF100466
2	DEN2-H-IMTSSA-MART-98-703	AF208496	
2	New Guinea C	AF038403	
2	New Guinea C-PUO-218 hybrid	AF038402	
2	New Guinea-C	M29095	
2	PDK-53	U87412	
45	2	S1 vaccine	NC_001474
2	TB16i	AY858036	
2	ThD2_0017_98	DQ181799	
2	ThD2_0026_88	DQ181802	
2	ThD2_0038_74	DQ181806	
2	ThD2_0055_99	DQ181798	
50	2	ThD2_0078_01	DQ181797
2	ThD2_0168_79	DQ181805	
2	ThD2_0263_95	DQ181800	
2	ThD2_0284_90	DQ181801	
2	ThD2_0433_85	DQ181803	
2	ThD2_0498_84	DQ181804	
55	2	ThNH-28-93	AF022435
2	ThNH29-93	AF169678	
2	ThNH36-93	AF169679	
2	ThNH45-93	AF169680	
2	ThNH-52-93	AF022436	
2	ThNH54-93	AF169682	
2	ThNH55-93	AF169681	
60	2	ThNH62-93	AF169683
2	ThNH63-93	AF169684	
2	ThNH69-93	AF169685	
2	ThNH73-93	AF169686	
2	ThNH76-93	AF169687	
2	ThNH81-93	AF169688	
65	2	ThNH-p36-93	AF022441
2	ThNH-7-93	AF022434	

TABLE A-continued

Examples of Dengue Virus Strains		
Serotype	Strain	Accession No.
2	ThNH-p11-93	AF022437
2	ThNH-p12-93	AF022438
2	ThNH-p14-93	AF022439
2	ThNH-p16-93	AF022440
2	Tonga-74	AY744147
2	TSV01	AY037116
2	Taiwan-1008DHF	AY776328
2	Ven2	AF100465
3	D3-H-IMTSSA-MART-1999-1243	AY099337
3	D3-H-IMTSSA-SRI-2000-1266	AY099336
3	80-2	AF317645
3	98901403	AB189125
3	98901437	AB189126
3	98901517	A6189127
3	98902890	AB189128
3	BA51	AY858037
3	BDH02-1	AY496871
3	BDH02-3	AY496873
3	BDH02-4	AY496874
3	BDH02-7	AY496877
3	BR74886-02	AY679147
3	C0331-94	AY876494
3	C0360-94	AY923865
3	den3_88	AY858038
3	den3_98	AY858039
3	FW01	AY858040
3	FW06	AY858041
3	H87	NC_001475
3	D3-Hu-TL018NIID-2005	AB214879
3	D3-Hu-TL029NIID-2005	AB214880
3	D3-Hu-TL109NIID-2005	AB214881
3	D3-Hu-TL129NIID-2005	AB214882
3	InJ_16_82	DQ401690
3	KJ30i	AY858042
3	kJ46	AY858043
3	kJ71	AY858044
3	mutant BDH02_01	DQ401689
3	mutant BDH02_03	DQ401691
3	mutant BDH02_04	DQ401692
3	mutant BDH02_07	DQ401693
3	mutant InJ_16_82	DQ401694
3	mutant PhMH_J1_97	DQ401695
3	PF89-27643	AY744677
3	PF89-320219	AY744678
3	PF90-3050	AY744679
3	PF90-3056	AY744680
3	PF90-6056	AY744681
3	PF92-2956	AY744682
3	PF92-2986	AY744683
3	PH86	AY858045
3	PhMH-J1-97	AY496879
3	PI64	AY858046
3	Singapore	AY662691
3	Singapore 8120-95	AY766104
3	Sleman-78	AY648961
3	TB16	AY858047
3	TB55i	AY858048
3	ThD3_0007_87	AY676353
3	ThD3_0010_87	AY676353
3	ThD3_0055_93	AY676351
3	ThD3_0104_93	AY676350
3	ThD3_1283_98	AY676349
3	ThD3_1687_98	AY676348
3	PF92-4190	AY744684
3	PF94-136116	AY744685
3	Taiwan-739079A	AY776329
4	2A	AF375822
4	Recombinant clone rDEN4	AF326825
4	2AdeI30	AF326826
4	814669	AF326573
4	B5	AF289029
4	rDEN4del30	AF326827
4	H241	AY947539
4	rDEN4	NC_002640
4	Singapore 8976-95	AY762085
4	SW38i	AY858050

TABLE A-continued

Examples of Dengue Virus Strains		
Serotype	Strain	Accession No.
4	ThD4_0017_97	AY618989
4	ThD4_0087_77	AY618991
4	ThD4_0348_91	AY618990
4	ThD4_0476_97	AY618988
4	ThD4_0485_01	AY618992
4	ThD4_0734_00	AY618993
4	Taiwan-2K0713	AY776330
4	Unknown	M14931

Mutations can be made in the 3'-UTR of a wild type infectious clone, e.g., dengue virus serotype 3, strain Sleman/78 or an infectious clone of another wild type, virulent dengue virus, and the mutants can then be tested in an animal model system (e.g., in mouse and/or monkey model systems) to identify sites that cause attenuation. Attenuation is judged by, for example, detection of decreased viremia. One or more additional mutations found to attenuate the wild-type virus are optionally introduced into a wild type dengue virus, and these mutants are tested in an animal model system (e.g., in a mouse and/or a monkey model system) to determine whether the resulting mutants are attenuated. Mutants that are found to be attenuated can then be used as new vaccine strains that have increased safety, due to attenuation.

In addition to the viruses listed above, dengue viruses including chimeric dengue viruses that include one or more attenuating mutations are included in the invention. These chimeras can consist of a dengue virus of a first serotype (i.e., a background dengue virus) in which a structural protein (or proteins) has been replaced with a corresponding structural protein (or proteins) of a dengue virus of a second serotype. For example, the chimeras can consist of a background dengue virus in which the prM and E proteins of the dengue virus of the first serotype have been replaced with the prM and E proteins of the dengue virus of the second serotype. The chimeric viruses can be made from any combination of dengue viruses of different serotypes. The dengue virus against which immunity is sought is the source of the inserted structural protein(s).

As is noted above, mutations that are included in the viruses of the present invention are attenuating. These mutations are present in the dengue virus 3'-UTR structure to attenuate the virus. Mutations can be made in the 3'-UTR using standard methods, such as site-directed mutagenesis. One example of the type of mutation present in the viruses of the invention is substitutions, but other types of mutations, such as deletions and insertions, can be used as well. In addition, as is noted above, the mutations can be present singly or in the context of one or more additional mutations.

Referring to FIG. 1, two approaches were taken to attenuate dengue virus. In one aspect, nucleotides additional to the $\Delta 30$ mutation were deleted from the 3'-UTR. In another aspect, the 3'-UTR of a dengue virus of a first serotype was replaced with the 3'-UTR from a dengue virus of a second serotype (optionally containing the $\Delta 30$ mutation and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR).

Deletion of Nucleotides Additional to the $\Delta 30$ Mutation from the 3'-UTR

Referring to FIGS. 2-5, using the first approach, the 3'-UTR of dengue viruses contain various conserved sequence motifs. The sequence of the DEN4 3'-UTR is illustrated in FIG. 5. The genome of DEN4 strain 814669 contains 10,646 nucleotides, of which the last 384 nt at the 3' terminus

are untranslated (non-coding). The locations of various sequence components in this region are designated with the reverse-direction numbering system. These sequences include the 3' distal secondary structure (nt 1 to 93), predicted to form stem-loop 1 (SL-1), which contains terminal loop 1 (TL-1). Nucleotides 117-183 form stem-loop 2 (SL-2) which contains TL-2. Nucleotides 201-277 form a pair of stem-loops (SL-3) which in part contains TL-3. Although the primary sequence of stem-loop 1 differs slightly among the dengue serotypes, the secondary structure is strictly conserved (compare FIGS. 2-5). Although the nucleotide spacing between SL-2 and neighboring SL-1 and SL-3 differ among the dengue virus serotypes, the overall structure of SL-2 is well-conserved. In addition, the exposed 9 nucleotides that comprise TL-2 are identical within all 4 dengue serotypes. It is TL-2 and its supporting stem structure that are removed by the $\Delta 30$ mutation (about nt 143-172). Removal of these 30 nucleotides results in formation of a new predicted structural element (SL-2 $\Delta 30$) which has a primary sequence and secondary structure which is identical for each of the dengue virus serotypes (compare FIGS. 6-9).

FIGS. 10-13 illustrate the approach where nucleotides additional to the $\Delta 30$ mutation are deleted from the 3'-UTR. The $\Delta 30$ mutation removes the TL-2 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4. The approach where nucleotides additional to the $\Delta 30$ mutation are deleted from the 3'-UTR removes the TL-2 homologous structure and sequence up to and optionally including the TL-3 homologous structure so that the deletion extends as far as the 5' boundary of the TL-3 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4. In the approach illustrated in FIGS. 10-14, an additional deletion of about 31 nucleotides from TL-3 results in formation of a new predicted structural element (SL-3 $\Delta 31$).

Referring to FIGS. 14-17, the $\Delta 86$ mutation removes the TL-2 homologous structure and removes sequence up to the TL-3 homologous structure in each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. This deletion results in the formation of a new predicted structural element (SL-2 $\Delta 86$).

In some embodiments that involve deletion of nucleotides additional to the $\Delta 30$ mutation, nucleic acid deletions are made to the 3'-UTR structure of the dengue virus genome to attenuate the virus while maintaining its immunogenicity. The deletions include the $\Delta 30$ deletion (nt 173-143 of the serotype 3 Sleman/78 strain in an exemplary manner or corresponding thereto in other strains of DEN1, DEN2, DEN3, or DEN4; numbering is from the 3' end of the viral genome) in addition to deletion of additional 3'-UTR sequence that is contiguous or non-contiguous to the $\Delta 30$ deletion. The $\Delta 30$ deletion corresponds to the TL-2 structure of the 3'-UTR. One type of embodiment, termed rDEN1 $\Delta 30/31$, rDEN2 $\Delta 30/31$, rDEN3 $\Delta 30/31$, or rDEN4 $\Delta 30/31$ includes the original $\Delta 30$ deletion and a non-contiguous 31 nt deletion that removes both the original TL-2 and TL-3 structures. Another type of embodiment, termed rDEN1 $\Delta 61$, rDEN2 $\Delta 61$, rDEN3 $\Delta 61$, or rDEN4 $\Delta 61$ includes the $\Delta 30$ deletion and deletion of 31 contiguous nucleotides extending 3' from the $\Delta 30$ deletion. Another type of embodiment, termed rDEN1 $\Delta 86$, rDEN2 $\Delta 86$, rDEN3 $\Delta 86$, or rDEN4 $\Delta 86$, includes the $\Delta 30$ deletion and deletion of 56 contiguous nucleotides extending 5' from the $\Delta 30$ deletion. For DEN3, a complete list of mutant viruses constructed to contain 3'-UTR deletion mutations is presented below in Table 2.

Replacement of the 3'-UTR of a Dengue Virus of a First Serotype with the 3'-UTR from a Dengue Virus of a Second Serotype

Using the second approach, the 3'-UTR of rDEN3 may be replaced with the 3'-UTR of rDEN4, optionally containing the $\Delta 30$ mutation and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR. Other examples include replacement of the 3'-UTR of rDEN3 with the 3'-UTR of dengue virus serotypes 1 and 2, optionally containing the $\Delta 30$ mutation and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR. Other examples include: replacement of the 3'-UTR of rDEN1 with the 3'-UTR of dengue virus serotypes 2, 3, and 4, optionally containing the $\Delta 30$ mutation and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR; replacement of the 3'-UTR of rDEN2 with the 3'-UTR of dengue virus serotypes 1, 3, and 4, optionally containing the $\Delta 30$ mutation and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR; and, replacement of the 3'-UTR of rDEN4 with the 3'-UTR of dengue virus serotypes 1, 2, and 3, optionally containing the $\Delta 30$ mutation and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR.

Embodiments that involve replacement of the 3'-UTR of a dengue virus of a first serotype with the 3'-UTR of dengue virus of a second serotype include:

- a) rDEN1-3'D2, rDEN1-3'D2x, rDEN1-3'D3, rDEN1-3'D3x, rDEN1-3'D4, rDEN1-3'D4x; rDEN1/2-3'D1, rDEN1/2-3'D1x, rDEN1/2-3'D3, rDEN1/2-3'D3x, rDEN1/2-3'D4, rDEN1/2-3'D4x; rDEN1/3-3'D1, rDEN1/3-3'D1x, rDEN1/3-3'D2, rDEN1/3-3'D2x, rDEN1/3-3'D4, rDEN1/3-3'D4x; rDEN1/4-3'D1, rDEN1/4-3'D1x, rDEN1/4-3'D2, rDEN1/4-3'D2x, rDEN1/4-3'D3, rDEN1/4-3'D3x;
 - b) rDEN2-3'D1, rDEN2-3'D1x, rDEN2-3'D3, rDEN2-3'D3x, rDEN2-3'D4, rDEN2-3'D4x; rDEN2/1-3'D2, rDEN2/1-3'D2x, rDEN2/1-3'D3, rDEN2/1-3'D3x, rDEN2/1-3'D4, rDEN2/1-3'D4x; rDEN2/3-3'D1, rDEN2/3-3'D1x, rDEN2/3-3'D2, rDEN2/3-3'D2x, rDEN2/3-3'D4, rDEN2/3-3'D4x; rDEN2/4-3'D1, rDEN2/4-3'D1x, rDEN2/4-3'D2, rDEN2/4-3'D2x, rDEN2/4-3'D3, rDEN2/4-3'D3x;
 - c) rDEN3-3'D1, rDEN3-3'D1x, rDEN3-3'D2, rDEN3-3'D2x, rDEN3-3'D4, rDEN3-3'D4x; rDEN3/1-3'D1, rDEN3/1-3'D2x, rDEN3/1-3'D3, rDEN3/1-3'D3x, rDEN3/1-3'D4, rDEN3/1-3'D4x; rDEN3/2-3'D1, rDEN3/2-3'D1x, rDEN3/2-3'D3, rDEN3/2-3'D3x, rDEN3/2-3'D4, rDEN3/2-3'D4x; rDEN3/4-3'D1, rDEN3/4-3'D1x, rDEN3/4-3'D2, rDEN3/4-3'D2x, rDEN3/4-3'D3, rDEN3/4-3'D3x; and
 - d) rDEN4-3'D1, rDEN4-3'D1x, rDEN4-3'D2, rDEN4-3'D2x, rDEN4-3'D3, rDEN4-3'D3x; rDEN4/1-3'D2, rDEN4/1-3'D2x, rDEN4/1-3'D3, rDEN4/1-3'D3x, rDEN4/1-3'D4, rDEN4/1-3'D4x; rDEN4/2-3'D1, rDEN4/2-3'D1x, rDEN4/2-3'D3, rDEN4/2-3'D3x, rDEN4/2-3'D4, rDEN4/2-3'D4x; rDEN4/3-3'D1, rDEN4/3-3'D1x, rDEN4/3-3'D2, rDEN4/3-3'D2x, rDEN4/3-3'D4, rDEN4/3-3'D4x;
- where x is a mutation listed in Table 2.

Method of Making and Using Dengue or Chimeric Dengue Viruses

The viruses (including chimeric viruses) of the present invention can be made using standard methods in the art. For example, an RNA molecule corresponding to the genome of a virus can be introduced into host cells, e.g., Vero cells, from which (or the supernatants of which) progeny virus can then be purified. In this method, a nucleic acid molecule (e.g., an RNA molecule) corresponding to the genome of a virus is

introduced into the host cells, virus is harvested from the medium in which the cells have been cultured, and the virus is formulated for the purposes of vaccination.

The viruses of the invention can be administered as primary prophylactic agents in adults or children at risk of infection, or can be used as secondary agents for treating infected patients. For example, in the case of DEN virus and chimeric DEN viruses, the vaccines can be used in adults or children at risk of DEN virus infection, or can be used as secondary agents for treating DEN virus-infected patients. Examples of patients who can be treated using the DEN virus-related vaccines and methods of the invention include (i) children in areas in which DEN virus is endemic, (ii) foreign travelers, (iii) military personnel, and (iv) patients in areas of a DEN virus epidemic. Moreover, inhabitants of regions into which the disease has been observed to be expanding (e.g., beyond Sri Lanka, East Africa and Latin America), or regions in which it may be observed to expand in the future can be treated according to the invention.

Formulation of the viruses of the invention can be carried out using methods that are standard in the art. Numerous pharmaceutically acceptable solutions for use in vaccine preparation are well known and can readily be adapted for use in the present invention by those of skill in this art (see, e.g., Remington's Pharmaceutical Sciences (18th edition), ed. A. Gennaro, 1990, Mack Publishing Co., Easton, Pa.). The viruses can be diluted in a physiologically acceptable solution, such as sterile saline, sterile buffered saline, or L-15 medium. In another example, the viruses can be administered and formulated, for example, as a fluid harvested from cell cultures infected with dengue virus or chimeric dengue virus.

The vaccines of the invention can be administered using methods that are well known in the art, and appropriate amounts of the vaccines administered can readily be determined by those of skill in the art. For example, the viruses of the invention can be formulated as sterile aqueous solutions containing between 10^2 and 10^7 infectious units (e.g., plaque-forming units or tissue culture infectious doses) in a dose volume of 0.1 to 1.0 ml, to be administered by, for example, intramuscular, subcutaneous, or intradermal routes. Further, the vaccines of the invention can be administered in a single dose or, optionally, administration can involve the use of a priming dose followed by a booster dose that is administered, e.g., 2-6 months later, as determined to be appropriate by those of skill in the art.

Optionally, adjuvants that are known to those skilled in the art can be used in the administration of the viruses of the invention. Adjuvants that can be used to enhance the immunogenicity of the viruses include, for example, liposomal formulations, synthetic adjuvants, such as (e.g., QS21), muramyl dipeptide, monophosphoryl lipid A, or polyphosphazene. Although these adjuvants are typically used to enhance immune responses to inactivated vaccines, they can also be used with live vaccines.

Nucleic Acid Sequences

Nucleic acid sequences of DEN viruses are useful for designing nucleic acid probes and primers for the detection of deletion or chimeric 3'-UTRs in a sample or specimen with high sensitivity and specificity. Probes or primers corresponding to deletion or chimeric 3'-UTRs can be used to detect the presence of deletion or chimeric 3'-UTRs in general in the sample, to quantify the amount of deletion or chimeric 3'-UTRs in the sample, or to monitor the progress of therapies used to treat DEN virus infection. The nucleic acid and corresponding amino acid sequences are useful as laboratory tools to study the organisms and diseases and to develop therapies and treatments for the diseases.

Nucleic acid probes and primers selectively hybridize with nucleic acid molecules encoding deletion or chimeric 3'-UTRs or complementary sequences thereof. By "selective" or "selectively" is meant a sequence which does not hybridize with other nucleic acids to prevent adequate detection of the deletion or chimeric 3'-UTRs. Therefore, in the design of hybridizing nucleic acids, selectivity will depend upon the other components present in the sample. The hybridizing nucleic acid should have at least 70% complementarity with the segment of the nucleic acid to which it hybridizes. As used herein to describe nucleic acids, the term "selectively hybridizes" excludes the occasional randomly hybridizing nucleic acids, and thus has the same meaning as "specifically hybridizing." The selectively hybridizing nucleic acid probes and primers of this invention can have at least 70%, 80%, 85%, 90%, 95%, 97%, 98% and 99% complementarity with the segment of the sequence to which it hybridizes, preferably 85% or more.

The present invention also contemplates sequences, probes and primers that selectively hybridize to the encoding nucleic acid or the complementary, or opposite, strand of the nucleic acid. Specific hybridization with nucleic acid can occur with minor modifications or substitutions in the nucleic acid, so long as functional species-species hybridization capability is maintained. By "probe" or "primer" is meant nucleic acid sequences that can be used as probes or primers for selective hybridization with complementary nucleic acid sequences for their detection or amplification, which probes or primers can vary in length from about 5 to 100 nucleotides, or preferably from about 10 to 50 nucleotides, or most preferably about 18-24 nucleotides. Isolated nucleic acids are provided herein that selectively hybridize with the species-specific nucleic acids under stringent conditions and should have at least five nucleotides complementary to the sequence of interest as described in *Molecular Cloning: A Laboratory Manual*, 2nd ed., Sambrook, Fritsch and Maniatis, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989.

If used as primers, the composition preferably includes at least two nucleic acid molecules which hybridize to different regions of the target molecule so as to amplify a desired region. Depending on the length of the probe or primer, the target region can range between 70% complementary bases and full complementarity and still hybridize under stringent conditions. For example, for the purpose of detecting the presence of deletion or chimeric 3'-UTRs, the degree of complementarity between the hybridizing nucleic acid (probe or primer) and the sequence to which it hybridizes is at least enough to distinguish hybridization with a nucleic acid from other organisms.

The nucleic acid sequences of the invention include a diagnostic probe that serves to report the detection of a cDNA amplicon amplified from the viral genomic RNA template by using a reverse-transcription/polymerase chain reaction (RT-PCR), as well as forward and reverse amplimers that are designed to amplify the cDNA amplicon. In certain instances, one of the amplimers is designed to contain a vaccine virus-specific mutation at the 3'-terminal end of the amplimer, which effectively makes the test even more specific for the vaccine strain because extension of the primer at the target site, and consequently amplification, will occur only if the viral RNA template contains that specific mutation.

Automated PCR-based nucleic acid sequence detection systems have been recently developed. TaqMan assay (Applied Biosystems) is widely used. A more recently developed strategy for diagnostic genetic testing makes use of molecular beacons (Tyagi and Kramer 1996 *Nature Biotechnology* 14:303-308). Molecular beacon assays employ quencher and

reporter dyes that differ from those used in the TaqMan assay. These and other detection systems may be used by one skilled in the art.

Dengue Virus Type 3 (DEN3) Vaccine Components Generated by Introduction of Deletions in the 3' Untranslated Region (UTR) or Exchange of the DEN3 3'-UTR with that of DEN4

There are four dengue virus serotypes (DEN1, DEN2, DEN3, and DEN4) which circulate in tropical and subtropical regions of the world inhabited by more than 2.5 billion people (Gubler D J 1998 Clin Microbiol Rev 11:480-496). DEN viruses are endemic in at least 100 countries and cause more human disease than any other arbovirus. Annually, there are an estimated 50-100 million dengue infections and hundreds of thousands of cases of dengue hemorrhagic fever/shock syndrome (DHF/DSS), with children bearing much of the disease burden (Gubler D J and Meltzer M 1999 Adv Virus Res 53:35-70). DHF/DSS remains a leading cause of hospitalization and death of children in at least eight southeast Asian countries (World Health Organization 1997 Dengue Haemorrhagic Fever: Diagnosis, Treatment, Prevention and Control, 2nd edition, WHO, Geneva). The dramatic increase in both the incidence and severity of disease caused by the four DEN serotypes over the past two decades is due in large part to the geographic expansion of the mosquito vectors, *Aedes aegypti* and *Aedes albopictus*, and the increased prevalence of the four DEN serotypes (Gubler D J 1998 Clin Microbiol Rev 11:480-496). The dengue viruses are maintained in a life cycle of transmission from mosquito to human to mosquito with no other apparent viral reservoir participating in this life cycle in urban settings (Rice CM, 1996 in Flaviviridae: The viruses and their replication, Fields B N, Knipe D M, Howley P M, Chanock R M, Melnick J L, Monath T P, Roizman B, Straus S E, eds. Fields Virology. Philadelphia: Lippincott-Raven Publishers, pp. 931-959).

The DEN viruses, members of the Flaviviridae family, have spherical virions of approximately 40 to 60 nm which contain a single-stranded positive-sense RNA genome. A single polypeptide is co-translationally processed by viral and cellular proteases generating three structural proteins (capsid C, membrane M, and envelope E) and at least seven non-structural (NS) proteins. The genome organization of the DEN viruses is 5'-UTR-C-prM-E-NS1-NS2A-NS2B-NS3-NS4A-NS4B-NS5-UTR-3' (UTR—untranslated region, prM—membrane precursor) (Rice C M, 1996 in Flaviviridae: The viruses and their replication, Fields B N, Knipe D M, Howley P M, Chanock R M, Melnick J L, Monath T P, Roizman B, Straus S E, eds. Fields Virology. Philadelphia: Lippincott-Raven Publishers, pp. 931-959).

In response to the increasing incidence and severity of DEN infection, development of vaccines is being pursued to prevent DEN virus disease. An economical vaccine that prevents disease caused by the DEN viruses has become a global public health priority. The cost-effectiveness, safety, and

long-term efficacy associated with the live attenuated vaccine against yellow fever (YF) virus, another mosquito-borne flavivirus, serves as a model for the feasibility of developing of a live attenuated DEN virus vaccine (Monath T P, 1999 in Yellow fever, Plotkin S A, Orenstein W A, eds. Vaccines, Philadelphia: W.B. Saunders Co., 815-879). Additionally, an effective live attenuated Japanese encephalitis (JE) virus vaccine is used in Asia, and inactivated virus vaccines are available for JE and tick-borne encephalitis virus. The need for a vaccine against the DEN viruses is mounting, and, despite much effort, the goal of developing a safe and efficacious DEN virus vaccine has yet to be attained. An effective DEN virus vaccine must confer protection from each serotype because all four serotypes commonly circulate in endemic regions and secondary infection with a heterologous serotype is associated with increased disease severity.

We have employed two strategies for generating live attenuated vaccine components against each serotype that can then be combined into tetravalent formulations (Blaney J E et al. 2006 Viral Immunol. 19:10-32). First, reverse genetics has been used to introduce an attenuating 30 nucleotide deletion ($\Delta 30$) mutation into the 3'-UTR of cDNA clones of each DEN serotype (Durbin, A P et al. 2001 Am J Trop Med Hyg 65:405-413; Whitehead S S et al. 2003 J Virol 77:1653-1657; Blaney J E et al. 2004 Am J Trop Med Hyg 71:811-821; Blaney J E et al. 2004 BMC Inf Dis 4:39). Initially, the rDEN4 $\Delta 30$ vaccine component was found to be attenuated in rhesus monkeys (Table 1) and phase I/II clinical trials in humans have demonstrated that virus infection results in low viremia, is strongly immunogenic, and exhibits minimal reactogenicity with no observation of serious adverse events (Durbin, A. P. et al. 2001 Am J Trop Med Hyg 65:405-413; Durbin et al. 2005 J Inf Dis 191:710-718). Recently, the rDEN1 $\Delta 30$ vaccine component, which was also attenuated in rhesus monkeys (Table 1), has been found to share a similar phenotype in clinical trials as that observed for rDEN4 $\Delta 30$; low viremia, strong immunogenicity, and minimal reactogenicity in 20 volunteers (Whitehead S S et al. 2003 J Virol 77:1653-1657; Blaney J E et al. 2006 Viral Immunol. 19:10-32). Unfortunately, the rDEN2 $\Delta 30$ and rDEN3 $\Delta 30$ vaccine components did not appear to be satisfactorily attenuated in rhesus monkeys during pre-clinical testing and there is no plan to test these in humans (Table 1) (Blaney J E et al. 2004 Am J Trop Med Hyg 71:811-821; Blaney J E et al. 2004 BMC Inf Dis 4:39). Consequently, an alternative strategy for vaccine development has been generation of antigenic chimeric viruses by replacement of structural proteins of the attenuated rDEN4 $\Delta 30$ vaccine component with those from DEN2 or DEN3 yielding the rDEN2/4 $\Delta 30$ and rDEN3/4 $\Delta 30$ vaccine components, respectively (Whitehead S S et al. 2003 Vaccine 21:4307-4316; Blaney J E et al. 2004 Am J Trop Med Hyg 71:811-821). The rDEN2/4 $\Delta 30$ vaccine virus has been tested in humans and appears safe and strongly immunogenic, while clinical evaluation of the rDEN3/4 $\Delta 30$ virus is currently planned.

TABLE 1

Effects of the $\Delta 30$ mutation on the four DEN serotypes in rhesus monkeys					
Virus	% of viremic monkeys	Mean no. of viremic days per monkey	Viremia ^a		
			Mean peak virus titer (\log_{10} PFU/ml \pm SE)	Geometric mean neutralizing antibody titers ^b	Reference
rDEN1	100	2.8	2.1 \pm 0.1	1,230	Whitehead et al.
rDEN1 $\Delta 30$	50	0.5	0.8 \pm 0.1	780	J. Virol, 2003, 77: 1653
rDEN2	100	4.0	1.9 \pm 0.1	173	Blaney et al. BMC Inf

TABLE 1-continued

Effects of the $\Delta 30$ mutation on the four DEN serotypes in rhesus monkeys					
Virus	Viremia ^a				Reference
	% of viremic monkeys	Mean no. of viremic days per monkey	Mean peak virus titer (log ₁₀ PFU/ml \pm SE)	Geometric mean neutralizing antibody titers ^b	
rDEN2 $\Delta 30$	100	2.8	1.7 \pm 0.2	91	Dis., 2004, 4: 39
rDEN3	100	2.3	1.4 \pm 0.2	363	Blaney et al. Am. J. Trop. Med. Hyg., 2004, 71: 811
rDEN3 $\Delta 30$	100	2.0	1.5 \pm 0.2	265	Hanley et al. Vaccine, 2004, 22: 3440
rDEN4	100	3.0	2.2 \pm 0.2	322	
rDEN4 $\Delta 30$	100	2.0	1.4 \pm 0.2	154	

^aGroups of rhesus monkeys were inoculated subcutaneously with 5.0 log₁₀ PFU of the indicated virus in a 1 ml dose. Serum was collected daily for 10 days. Virus titer in serum was determined by plaque assay in Vero cells.

^bPlaque reduction (60%) neutralizing antibody titers were determined on day 28 serum using indicated wild type virus. Reciprocal dilution of geometric mean is indicated.

Here, we describe novel vaccine components for the DEN3 serotype generated by genetic modification of the 3'-UTR of the DEN3 cDNA clone (Blaney J E et al. 2004 Am J Trop Med Hyg 71:811-821). Development of these DEN3 vaccine components, which possess the full complement of wild type DEN3 proteins, is important for two reasons. First, the present vaccine component for DEN3, rDEN3/4 $\Delta 30$, may be found to be under- or over-attenuated in clinical trials. Second, an optimal vaccine for conferring protection from disease caused by DEN3 may require induction of T cell responses against the entire set of DEN3 proteins, rather than just the M and E which are the only DEN3 sequences present in the rDEN3/4 $\Delta 30$ chimeric virus. To generate additional DEN3 vaccine components, novel deletions which encompass or border the $\Delta 30$ deletion in the 3'-UTR were introduced into the rDEN3 cDNA clone. Alternatively, the 3'-UTR of the rDEN3 cDNA clone was replaced with that of rDEN4 or rDEN4 $\Delta 30$. Viable viruses were analyzed for attenuation phenotypes in tissue culture, SCID mice transplanted with HuH-7 cells, and rhesus monkeys. Three mutant viruses (rDEN3 $\Delta 30/31$, rDEN3 $\Delta 86$, and rDEN3-3'D4 $\Delta 30$) have pre-clinical phenotypes which suggest they may be safe and immunogenic in humans.

Generation of rDEN3 Deletion Mutants

We sought to generate expanded deletion mutations which include the original $\Delta 30$ (nt 173-143) mutation. Table 2 lists seven deletion mutations which encompass the original $\Delta 30$ mutation including $\Delta 50$, $\Delta 61$, $\Delta 80$, $\Delta 86$, $\Delta 116A$, $\Delta 116B$, and $\Delta 146$. In addition, the $\Delta 30/31$ mutation includes the original $\Delta 30$ mutation and a non-contiguous 31 nt deletion. The $\Delta 31$ mutation was also generated alone to discern the contribution of either $\Delta 30$ or $\Delta 31$ in the combined $\Delta 30/31$ deletion mutation. The location of bordering nucleotides of deletions in the predicted secondary structure of the DEN3 3'-UTR are indicated in FIG. 4. In addition, the predicted secondary structure of the DEN3 3'-UTR for rDEN3 $\Delta 30$, rDEN3 $\Delta 30/31$, and rDEN3 $\Delta 86$ are indicated in FIGS. 8, 12, and 16, respectively.

TABLE 2

Deletion mutations created in the 3'-UTR of DEN3 Sleman/78		
Mutation	Deleted nucleotides ^a	Deletion junction
$\Delta 30$	173-143	-CCAA Δ GACU-
$\Delta 31$	258-228	-CUGC Δ GACU-
$\Delta 50$	192-143	-CACA Δ GACU-
$\Delta 61$	173-113	-CCGA Δ UAAA-
$\Delta 80$	192-113	-CACA Δ UAAA-

TABLE 2-continued

Deletion mutations created in the 3'-UTR of DEN3 Sleman/78		
Mutation	Deleted nucleotides ^a	Deletion junction
$\Delta 86$	228-143	-UAGCAGACU-
$\Delta 116$ (A)	228-113	-UAGCAUAAA-
$\Delta 116$ (B)	258-143	-CUGCAGACU-
$\Delta 146$	258-113	-CUGC Δ UAAA-
$\Delta 30/31$	173-143	-CCAA Δ GACU-
	258-228	-CUGCAGACU-

^aNumbering is from the 3'-end of viral genome

PCR mutagenesis was used to introduce the nine new deletion mutations into the DEN3 Sleman/78 cDNA plasmid, p3, which was previously used to generate the rDEN3 $\Delta 30$ vaccine component (Blaney J E et al. 2004 Am J Trop Med Hyg 71:811-821). The p3-frag.4 cDNA subclone was used as the template for PCR reactions with indicated pairs of mutagenic oligonucleotides listed in Table 3, except for the $\Delta 30/31$ deletion mutation which used p3-frag.4 $\Delta 30$ cDNA subclone as a template. PCR products were ligated and used to transform competent bacterial cells. Plasmid DNA was isolated from bacterial clones and the presence of the appropriate deletion mutation was confirmed by sequence analysis. To generate intact DEN3 cDNA plasmids containing the deletion mutations, the KpnI-PstI fragment (963 nt) from the mutated p3-frag.4 cDNA subclones were introduced into the p3-7164 cDNA plasmid. The p3-7164 plasmid encodes the 7164 Vero cell adaptation mutation which had previously been shown to enhance growth and transfection efficiency in Vero cells (Blaney J E et al. 2004 Am J Trop Med Hyg 71:811-821). Full length p3 plasmids containing the deletion mutations were confirmed to contain the correct 3'-UTR sequence. Mutations in addition to the engineered deletions were identified in the rDEN3 $\Delta 30/31$ and rDEN3 $\Delta 86$ viruses when compared to the DEN3 p3 plasmid cDNA (Genbank # AY656169) (Table 4).

TABLE 3

Mutagenic primer sequences for construction of 3'-UTR deletions	
Primer name	Sequence (5'→3')
113F	TAAAAACAGCATATTGACGCTGGGAG (SEQ ID NO: 24)
143F	GACTAGAGGTTAGAGGAGAC (SEQ ID NO: 25)

TABLE 3-continued

Mutagenic primer sequences for construction of 3'-UTR deletions	
Primer name	Sequence (5'→3')
228F	GACTAGCGGTTAGAGGAGACCCC (SEQ ID NO: 26)
173R	TCGGGCCCGCTGCTGCGTTG (SEQ ID NO: 27)
173R (for Δ30)	TTGGGCCCGCTGCTGCGTTG (SEQ ID NO: 28)
192R	TGTGTCATGGGAGGGGTCTC (SEQ ID NO: 29)
228R	GCTACACCGTTCGTACAGCTTCC (SEQ ID NO: 30)
258R	GCAGCCTCCAGGTTTACGTCC (SEQ ID NO: 31)

TABLE 4

Mutations in addition to the engineered deletions that were identified in the rDEN3Δ30/31 and rDEN3Δ86 viruses when compared to the DEN3 p3 plasmid cDNA (Genbank # AY656169)

Virus	Gene	Nucleotide position	Nucleotide substitution	Amino acid position	Amino acid change
rDEN3Δ30/31	NS4B	7164a	U → C	115	Val → Ala
	NS4B	7398	C → U	193	Ala → Val
rDEN3Δ86	M	512	A → G	26	Lys → Glu
	NS3	6076	C → U	521	silent
	NS4B	7164a	U → C	115	Val → Ala
	NS5	8623	U → C	353	silent
	NS5	10267b	A → U	END	end → Tyr
	3'-UTR	10455	G → C	—	—

^aThe 7164 mutation is a Vero cell adaptation mutation which was engineered into the cDNA construct.

^bThere is a mixed population at this nt position (A → A/U) that changes the stop codon (UAA) at the end of NS5 to UAU which encodes Tyr. This would serve to extend NS5 by 2 amino acids (Tyr-Thr-End) because a stop codon remains at nts 10271-10273.

For recovery of viruses, 5'-capped RNA transcripts were synthesized in vitro from cDNA plasmids and transfected into either Vero cells or C6/36 cells. Prior to transcription and generation of infectious virus, the linker sequences were removed from cDNA plasmids by digestion with SpeI. Plasmids were then recircularized by ligation, linearized with Acc65I (isoschizomer of KpnI which cleaves leaving only a single 3' nucleotide), and transcribed in vitro using SP6 polymerase. Purified transcripts were then transfected into Vero or C6/36 cells.

Recombinant viruses encoding each of the nine mutations, Δ30/31, Δ31, Δ50, Δ61, Δ80, Δ86, Δ116A, Δ116B, and Δ146, were successfully recovered in C6/36 cells, while only rDENΔ31 was recovered in Vero cells. The rDEN3 deletion mutant viruses were then passaged once in Vero cells followed by biological cloning by two terminal dilutions in Vero cells. Cloned viruses were then passaged two to seven times in Vero cells in an attempt to reach a stock titer of at least 6.0 log₁₀ PFU/ml which is considered sufficient to allow for cost-effective manufacture. Three recombinant viruses (rDEN3Δ50, rDEN3Δ116A, and rDEN3Δ146) were found to be excessively restricted for replication in Vero cells, despite being viable. Therefore, these three viruses were not studied further. The genetic sequence of the 3'-UTR was determined for the six remaining deletion mutant viruses that reached peak virus titers of at least 6.0 log₁₀ PFU/ml. The correct 3'-UTR sequence with the appropriate deletion was found for rDEN3Δ61, rDEN3Δ80, rDEN3Δ86 and rDEN3Δ30/31. However, two mutant viruses were found to contain addi-

tional deletions or mutations and were deemed to potentially have unstable genotypes. First, rDEN3Δ31 had the correct 3'-UTR deletion of nt 258-228 but also contained a 25 nt deletion of nt 222-198. Second, rDEN3Δ116B had the correct 3'-UTR deletion of nt 258-143 but also contained a 8 nt deletion of nt 430-423 and a single A→G substitution at nt 265. The potential of genetic instability with these viruses precludes their use as vaccine components so they were not further studied. Therefore, of the nine original deletions constructed, four mutant viruses were found to replicate efficiently in Vero cells and were studied further; rDEN3Δ61, rDEN3Δ80, rDEN3Δ86 and rDEN3Δ30/31.

Generation of rDEN3 Chimeric Viruses with the 3'-UTR Derived from rDEN4 or rDEN4Δ30

Another strategy was employed to generate novel rDEN3 vaccine components; replacement of the 3'-UTR of the rDEN3 cDNA clone with that of rDEN4 or rDEN4Δ30 (FIG. 18A). The 3'-UTR chimeric virus, rDEN3-3'D4Δ30, was designed to be a vaccine component for inclusion in tetravalent formulations which share the Δ30 deletion mutation among all four serotypes. The rDEN3-3'D4 virus was designed to discern the contribution of the 3'-UTR chimerization and the Δ30 mutation to any observed phenotypes.

The p3-3'D4Δ30 plasmid was generated as follows. First, PCR mutagenesis was used to introduce a HpaI restriction site into the p3-frag.4 cDNA subclone (FIG. 18B). PCR products were ligated and used to transform competent bacterial cells. Plasmid DNA was isolated from bacterial clones and the presence of the appropriate deletion mutation was confirmed by sequence analysis. To introduce the rDEN4Δ30 3'-UTR into the p3-frag.4 (HpaI) cDNA subclone, a 364 nt fragment encompassing the p4Δ30 3'-UTR was amplified by PCR using a forward primer (5'-AACAACAACAACACCAAAGGCTATTG-3', SEQ ID NO: 32) and reverse primer (5'-CCTACCGGTACCAGAACCCTGTTG-3', SEQ ID NO: 33). To generate the p3-frag.4-3'D4Δ30 cDNA subclone, the HpaI-KpnI fragment was removed from p3-frag.4 (HpaI) and replaced with the p4Δ30 3'-UTR PCR fragment which had been cleaved by KpnI. The PstI-KpnI fragment of p3-frag.4-3'D4Δ30 was introduced into the p3 plasmid to make the full length cDNA clone, p3-3'D4Δ30. The sequence of the 3'-UTR and NS5 junction were confirmed to be correct.

To generate p3-3'D4, the 30 deleted nucleotides of the Δ30 deletion mutation were introduced into the p3-frag.4-3'D4Δ30 subclone. Briefly, the MluI-KpnI fragment of p3-frag.4-3'D4Δ30, which encompasses the Δ30 region, was replaced with the corresponding fragment of p4 to make the plasmid, p3-frag.4-3'D4. To generate a full length p3 genome, the PstI-KpnI fragment of p3 was replaced with the corresponding fragment of p3-frag.4-3'D4. The 3'-UTR sequence of the p3-3'D4 plasmid was determined to be correct and contained the missing 30 nt of the Δ30 mutation.

For recovery of viruses, 5'-capped RNA transcripts were synthesized in vitro from cDNA plasmids and transfected into either Vero cells or C6/36 cells. Prior to transcription and generation of infectious virus, the linker sequences were removed from cDNA plasmids by digestion with SpeI. Plasmids were then recircularized by ligation, linearized with Acc65I (isoschizomer of KpnI which cleaves leaving only a single 3' nucleotide), and transcribed in vitro using SP6 polymerase. Purified transcripts were then transfected into Vero or C6/36 cells.

rDEN3-3'D4 was recovered in C6/36 cells and Vero cells, while rDEN3-3'D4Δ30 could only be recovered in Vero cells. Mutant viruses were then passaged once in Vero cells followed by biological cloning by two terminal dilutions in Vero cells. rDEN3-3'D4 and rDEN3-3'D4Δ30 were then passaged four or six times in Vero cells, respectively. The genetic sequence of the NS5-3'-UTR junction and 3'-UTR was found

19

to be correct for rDEN3-3'D4 and rDEN3-3'D4Δ30. Therefore, both viruses were studied further.

Mutations were also identified in the rDEN3-3'D4Δ30 virus compared to the DEN3 p3 plasmid cDNA clone (5'-UTR and genes) and DEN4 p4 cDNA clone (3'-UTR) (Table 5).

TABLE 5

Mutations in the rDEN3-3D4Δ30 virus compared to the DEN3 p3 plasmid cDNA clone (5'-UTR and genes) and DEN4 p4 cDNA clone (3'-UTR)					
Virus	Gene	Nucleotide position	Nucleotide substitution	Amino acid position	Amino acid change
rDEN3-3'D4Δ30	C	250	U → C	52	silent
	NS3	5899	U → C	462	silent
	NS4B ^a	7164	U → C	115	Val → Ala
	3'-UTR	10534	A → G	—	—

^aThe 7164 mutation is a Vero cell adaptation mutation which was engineered into the cDNA construct.

Replication of DEN3 Mutant Viruses in SCID-HuH-7 Mice

The four deletion mutant viruses (rDEN3Δ30/31, rDEN3Δ61, rDEN3Δ80, and rDEN3Δ86) which were found to replicate to high titer in Vero cells and were confirmed to have the correct 3'-UTR sequence and the rDEN3-3'D4 and rDEN3-3'D4Δ30 viruses were first evaluated in SCID-HuH-7 mice. The rDEN3-3'D4 and rDEN3-3'D4Δ30 were compared to determine the effect on replication of the 3'-UTR chimerization and any further attenuation conferred by the Δ30 mutation. SCID-HuH-7 mice contain solid tumors of the HuH-7 human hepatoma cell line, and analysis of virus replication in this mouse model serves as a surrogate for DEN virus replication in the human liver. Numerous DEN virus mutant viruses have been identified by evaluation in SCID-HuH-7 mice (Blaney J E, et al. 2002 Virology 300:125-139; Hanley et al. 2004 Vaccine 22:3440-3448; Blaney J E et al. 2006 Viral Immunol. 19:10-32). This mouse model provided the original evidence that the rDEN3Δ30 virus was not attenuated compared to parent virus rDEN3, while the antigenic chimeric virus, rDEN3/4Δ30, was approximately 100-fold restricted in replication in the SCID-HuH-7 mice when compared to wild type parent viruses (Blaney J E et al. 2004 Am J Trop Med Hyg 71:811-821).

For analysis of virus replication in SCID-HuH-7 mice, four to six week-old SCID mice (Tac:1cr:Ha(ICR)-Prkdc^{scid}) (Taconic Farms) were injected intraperitoneally with 0.1 mL phosphate-buffered saline containing 10⁷ HuH-7 cells which had been propagated in tissue culture. Tumors were detected in the peritoneum five to six weeks after transplantation, and tumor-bearing mice were infected by direct inoculation into the tumor with 10⁴ PFU of virus in 50 μl Opti-MEM I. Serum was collected from infected mice on day 7 post-infection and frozen at -80° C. The virus titer was determined by plaque assay in Vero cells.

As indicated in Table 6, wild type DEN3 Sleman/78 replicated to a mean peak virus titer of nearly 10^{6.9} PFU/ml. Although a decreased level of replication was observed for each of the six mutant viruses, the differences in replication were not statistically significant. However, rDEN3Δ86 and rDEN3-3'D4Δ30 were more than 10-fold restricted in replication compared to wild type DEN3 virus, while the replication of rDEN3Δ30/31 was slightly less than 10-fold restricted. On the basis of this arbitrary cut-off, these three viruses were selected for further evaluation. It is important to note that the rDEN4Δ30 virus which has a well-characterized, attenuation and non-reactogenic phenotype in humans was found to be only 6-fold restricted in replication in SCID-

20

HuH-7 mice compared to wild type rDEN4 virus (Hanley et al. 2004 Vaccine 22:3440-3448).

TABLE 6

Replication of mutant DEN3 viruses in HuH-7-SCID mice.			
Virus ¹	No. of mice	Mean peak virus titer (log ₁₀ pfu/ml ± SE)	Fold-reduction compared to DEN3 (Sleman/78)
10 DEN3 (Sleman/78)	8	6.9 ± 0.1	—
rDEN3Δ30/31	8	6.0 ± 0.3	8
rDEN3Δ61	9	6.3 ± 0.2	4
rDEN3Δ80	9	6.3 ± 0.3	4
rDEN3Δ86	10	5.6 ± 0.4	20
rDEN3-3'D4	11	6.5 ± 0.4	3
15 rDEN3-3'D4Δ30	9	5.7 ± 0.2	16

¹Groups of HuH-7-SCID mice were inoculated into the tumor with 4.0 log₁₀ PFU of the indicated virus. Serum was collected on day 7 and virus titer was determined in Vero cells.

Because the rDEN3-3'D4Δ30 virus and the rDEN3Δ30/31 and rDEN3Δ86 viruses encode the full set of DEN3 structural and non-structural proteins, they would be expected to induce the full complement of humoral and cellular immunity. This more complete immune induction would be advantageous compared to that induced by the chimeric rDEN3/4Δ30, which encodes only the structural proteins of DEN3.

Replication of DEN3 Mutant Viruses in Tissue Culture

The level of virus replication in Vero cells and mosquito C6/36 cells was assessed for the rDEN3Δ30/31 and rDEN3Δ86 deletion mutant viruses and the rDEN3-3'D4 viruses with and without Δ30. Replication in Vero cells was analyzed because these cells are the substrate for manufacture, while growth in C6/36 cells was assessed because attenuation phenotypes in these mosquito cells may be associated with restricted replication in Aedes mosquitoes which serve as the vector for DEN virus transmission (Hanley et al. 2003 Virology 312:222-232).

Growth kinetics were evaluated as follows. Confluent monolayers of Vero cells and C6/36 cells in 75 cm² flasks were infected at a multiplicity of infection of 0.01. Aliquots of 0.5 ml tissue culture supernatant were removed daily for seven days, combined with SPG stabilizer, and frozen at -80° C. Virus titer of all samples was determined by plaque assay in Vero cells. The limit of detection for the plaque assay is 1.0 log₁₀ PFU/ml.

The replication kinetics of each virus in both cell lines is shown in FIG. 19. In Vero cells, rDEN3Δ30/31, rDEN3Δ86, and rDEN3-3'D4Δ30 replicated to a peak level that approximated that of wild type rDEN3 and with similar kinetics to that of wild type rDEN3. These three vaccine components reached peak virus titers of 6.5 to 6.7 log₁₀ PFU/ml which demonstrates the feasibility of manufacture for each of these viruses. In Vero cells, the rDEN3-3'D4 virus replicated to a peak titer of 7.8 log₁₀ PFU/ml which is nearly 100-fold higher than that observed for wild type rDEN3 indicating that inclusion of the DEN4 3'-UTR may augment replication in Vero cells. This could also be attributed to more efficient Vero cell adaptation of the rDEN3-3'D4 virus. rDEN4 replicates to a peak titer of approximately 8.0 log₁₀ PFU/ml which indicates that the chimeric virus achieves a peak titer that does not exceed that of either of its parent viruses (Blaney J E et al. 2006 Viral Immunol. 19:10-32).

Analysis of virus replication in C6/36 cells demonstrated that rDEN3Δ86 and rDEN3-3'D4Δ30 reached peak titers approximately 10-fold lower than the peak virus titer of wild type rDEN3 virus, 6.9 log₁₀ PFU/ml (FIG. 19). The rDEN3-3'D4 virus replicated to a peak titer similar to that observed for wild type rDEN3. The most striking result was the lack of replication observed in C6/36 cells for the rDEN3Δ30/31 virus. After day 1, virus was not detected in culture medium

from C6/36 cells infected with rDEN3Δ30/31 virus despite the efficient replication observed in Vero cells. These results were confirmed in a second independent growth curve experiment and indicate a host range attenuation phenotype in tissue culture which is envisioned as being accompanied by an attenuation phenotype in mosquitoes as well.

Replication and Immunogenicity of DEN3 Mutant Viruses in Rhesus Monkeys

Based on the slight attenuation in SOD-HuH-7 mice and efficient growth in Vero cells, rDEN3Δ30/31, rDEN3Δ86, and rDEN3-3'D4Δ30 were evaluated in rhesus monkeys. The mutant viruses were compared with wild type DEN3 for level and duration of viremia, neutralizing antibody induction, and the ability to confer protection from wild type DEN3 virus challenge. The rDEN3-3'D4 virus was also evaluated to discern the contribution of the 3'-UTR chimerization upon attenuation with and without the Δ30 mutation. An attenuation phenotype in rhesus monkeys has generally been a strong predictor of safety for vaccine components in clinical trials including rDEN4Δ30, rDEN1Δ30, and rDEN2/4Δ30 (Blaney J E et al. 2006 *Viral Immunol.* 19:10-32).

Groups of four rhesus monkeys were inoculated subcutaneously with 10^5 PFU of the indicated viruses (Table 7). Two monkeys were mock infected with virus diluent. For detection of viremia, serum was collected on days 0-8 and on day 10 and frozen at -80° C. Virus titer in serum samples was determined by plaque assay in Vero cells. Serum was collected on day 28 for detection of neutralizing antibodies directed against DEN3. Levels of neutralizing antibodies were determined using a plaque reduction neutralization assay in Vero cells against wild type DEN3 virus. On day 35 post-infection, all monkeys were challenged by subcutaneous infection with 10^5 PFU DEN3 wild type virus. Serum was collected on days 0-8 and on day 10 and frozen at -80° C. Virus titer in serum samples was determined by plaque assay in Vero cells.

TABLE 7

Virus ¹	No. of monkeys	% of monkeys with viremia	Mean no. of viremic days per monkey	Mean peak virus titer ² log ₁₀ pfu/ml ±SE	Geometric mean serum neutralizing antibody titer (reciprocal dilution) ³		Post-challenge ⁴	
					Day 0	Day 28	% of monkeys with viremia	Mean peak virus titer ² (log ₁₀ pfu/ml ± SE)
					DEN3 (Sleman/78)	4	100	3.5
rDEN3Δ30/31	4	0	0	<1.0	<5	304	0	<1.0
rDEN3Δ86	4	0	0	<1.0	<5	224	0	<1.0
rDEN3-3'D4	4	75	1.5	1.3 ± 0.2	<5	229	0	<1.0
rDEN3-3'D4Δ30	4	0	0	<1.0	<5	77	0	<1.0
mock infected	2	0	0	<1.0	<5	<5	100	1.8 ± 0.2

¹Groups of rhesus monkeys were inoculated subcutaneously on day 0 with $5.0 \log_{10}$ PFU of the indicated virus in a 1 ml dose. Serum was collected daily on days 0-8 and 10 and once on day 28.

²Virus titer in serum was determined by plaque assay in Vero cells.

³Plaque reduction (60%) neutralizing antibody titers were determined using DEN3 (Sleman/78).

⁴Monkeys were challenged after 35 days with DEN3 (Sleman/78) administered subcutaneously in a 1 ml dose containing $5.0 \log_{10}$ PFU. Serum was collected daily on days 0-8 and 10.

Wild type DEN3 Sleman/78 virus replicated in rhesus monkeys to a mean peak virus titer of $1.8 \log_{10}$ PFU/ml serum with all monkeys developing viremia (Table 7). These results parallel previous studies of DEN3 in rhesus monkeys (Blaney J E et al. 2004 *Am J Trop Med Hyg* 71:811-821). No viremia was detected in any monkey infected with any of the three vaccine components, rDEN3Δ30/31, rDEN3Δ86, or rDEN3-3'D4Δ30 demonstrating a clear attenuation phenotype for each of these viruses in rhesus monkeys. Interestingly, the rDEN3-3'D4 virus was detected in 75% of monkeys with a mean peak virus titer of $1.3 \log_{10}$ PFU/ml serum suggesting that the presence of the Δ30 mutation is critical for attenuation

of the 3'-UTR chimeric virus. Despite the lack of detectable viremia, mean neutralizing antibody levels in monkeys infected with rDEN3Δ30/31 and rDEN3Δ86 reached levels similar to that of wild type DEN3 virus, 1:253 (Table 7). In contrast, the rDEN3-3'D4Δ30 virus induced mean neutralizing antibody levels approximately three-fold lower than DEN3. However, 100% of monkeys immunized with each vaccine component seroconverted as defined by a four-fold or greater rise in serum neutralizing antibody levels after infection. Thus all monkeys were deemed to be infected by each of the vaccine components despite the lack of detectable viremia. Determination of virus titer in serum after challenge with DEN3 virus indicated that immunization with each of the vaccine components induced complete protection from detectable viremia as would be expected given the observed neutralizing antibody levels.

Replication in Mosquitoes

Replication of rDEN3 and rDEN3Δ30/31 was studied in *Toxorhynchites amboinensis* mosquitoes. Intrathoracic inoculation of serial ten-fold dilutions of test virus was performed as described previously (Troyer J. M. et al. 2001 *Am. J. Trop. Med. Hyg.* 65:414-9). After a 14 day incubation, heads were separated and homogenized in diluent. Virus titer in head homogenates was determined by plaque assay in Vero cells.

Based on the attenuation of rDEN3Δ30/31 in rhesus monkeys and its restricted replication in C6/36 mosquito cells, rDEN3Δ30/31 was compared to wild type rDEN3 for infectivity and level of replication in highly sensitive *Toxorhynchites amboinensis* mosquitoes (Table 8). Ten-fold serial dilutions of virus were inoculated intrathoracically, and the ability to infect head tissues was evaluated by performing a plaque assay on mosquito head homogenates after a 14 day incubation.

The infectivity of rDEN3 and rDEN3Δ30/31 was very similar as the 50% mosquito infectious dose was approximately $10^{1.3}$ PFU for both viruses (Table 8). However, the level of replication of rDEN3Δ30/31 in the heads of infected mosquitoes was about 5- to 50-fold reduced. This reduction was significant at the $10^{2.3}$ and $10^{1.3}$ PFU doses tested. This finding indicates that although rDEN3Δ30/31 has infectivity for *Toxorhynchites* by intrathoracic infection similar to that of wild type rDEN3, there is a statistically significant restriction in the level of replication in mosquitoes afforded by the Δ30/31 mutation.

TABLE 8

Replication of rDEN3 and rDEN3Δ30/31 in <i>Toxorhynchites amboinensis</i>					
Virus	Dose ^a (log ₁₀ PFU)	No tested	% infected ^b	Mean virus titer ^c (log ₁₀ PFU/head)	Reduction (log ₁₀) compared to same dose of wt virus
rDEN3 wt	2.3	20	90	4.2 ± 0.1 ^d	
	1.3	19	53	4.2 ± 0.1 ^e	
	0.3	17	18	4.3 ± 0.3	
rDEN3Δ30/31	2.3	12	83	2.7 ± 0.3 ^d	1.5
	1.3	16	44	3.1 ± 0.3 ^e	1.1
	0.3	8	13	3.6 ± 0.0	0.7

^aVirus titer administered intrathoracically in a 0.2 μl inoculum.

^bPercentage of mosquitoes with detectable virus at day 14 post-inoculation was determined by plaque assay on mosquito head homogenates in Vero cells.

^cCalculated using only values of virus-positive heads.

^dFor 10^{2.3} PFU dose of rDEN3 and rDEN3Δ30/31, mean virus titers were significantly different as determined by a Tukey-Kramer post-hoc test (P < 0.001).

^eFor 10^{1.3} PFU dose of rDEN3 and rDEN3Δ30/31, mean virus titers were significantly different as determined by a Tukey-Kramer post-hoc test (P < 0.005).

While the present invention has been described in some detail for purposes of clarity and understanding, one skilled in the art will appreciate that various changes in form and detail can be made without departing from the true scope of the

invention. All figures, tables, and appendices, as well as patents, applications, and publications, referred to above, are hereby incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 33

<210> SEQ ID NO 1

<211> LENGTH: 276

<212> TYPE: DNA

<213> ORGANISM: Dengue virus

<400> SEQUENCE: 1

```

uaaaaccugg gaggcugcaa acuguggaag cuguacgcac gguguagcag acuagcgguu      60
agaggagacc ccucccauga cacaacgcag cagcggggcc cgagcucuga gggaagcugu      120
accuccuugc aaaggacuag agguuagagg agaccccccg caaaiaaaaa cagcauauug      180
acgcugggag agaccagaga uccugcuguc uccucagcau cauuccaggc acagaacgcc      240
agaaaugga auggugcugu ugaaucaaca gguucu                                     276

```

<210> SEQ ID NO 2

<211> LENGTH: 278

<212> TYPE: DNA

<213> ORGANISM: Dengue virus

<400> SEQUENCE: 2

```

uaaaaacccg ggaggcugca aaccauggaa gcuguacgca uggguagca gacuaguggu      60
uagaggagac ccucccaag acacaacgca gcagcggggc ccaacaccag gggaagcugu      120
accugggugg uaaggacuag agguuagagg agaccccccg cacaacaaca aacagcauau      180
ugacgcuggg agagaccaga gaucugcug ucucuacagc aucauuccag gcacagaacg      240
ccagaaaaug gaauggugcu guugaaucaa cagguucu                                     278

```

<210> SEQ ID NO 3

<211> LENGTH: 281

<212> TYPE: DNA

<213> ORGANISM: Dengue virus

<400> SEQUENCE: 3

```

uaaaaaaucc gggaggccac aaaccaugga agcuguacgc auggcguagu ggacuagcgg      60
uuagaggaga cccucccuu acagaucgca gcaacaauagg gggcccaagg ugagaugaag      120

```

-continued

```

cuguagucuc acuggaagga cuagagguua gaggagaccc ccccaaaaca aaaaacagca 180
uauugacgcu gggaaagacc agagaucug cugucuccuc agcaucauuc caggcacagg 240
acgccagaaa auggaauggu gcuguugaau caacagguuc u 281

```

```

<210> SEQ ID NO 4
<211> LENGTH: 276
<212> TYPE: DNA
<213> ORGANISM: Dengue virus

```

```

<400> SEQUENCE: 4

```

```

uaaaaccugg gaggcugcaa acuguggaag cuguacgcac gguguagcag acuagcgguu 60
agaggagacc ccuccauga cacaacgcag cagcggggcc cgagcucuga gggaagcugu 120
accuccuugc aaaggacuag agguuagagg agaccccccg caaauaaaaa cagcauuug 180
acgcugggag agaccagaga uccugcuguc uccucagcau cauuccaggc acagaacgcc 240
agaaaugga auggugcugu ugaaucaaca gguucu 276

```

```

<210> SEQ ID NO 5
<211> LENGTH: 281
<212> TYPE: DNA
<213> ORGANISM: Dengue virus

```

```

<400> SEQUENCE: 5

```

```

auccccaggg aggccaugcg ccacggaagc uguacgcgug gcuuuuugga cuagcgguaa 60
gaggagaccc cucccaucac ugacaaaacg cagcaaaagg gggcccgaag ccaggaggaa 120
gcuguacucc ugguggaagg acuagagguu agaggagacc ccccaaacac aaaaacagca 180
uauugacgcu gggaaagacc agagaucug cugucucugc aacaucuauc caggcacaga 240
gcgccgaag auggauuggu guuguugauc caacagguuc u 281

```

```

<210> SEQ ID NO 6
<211> LENGTH: 248
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Dengue virus construct

```

```

<400> SEQUENCE: 6

```

```

uaaaaacccg ggaggcugca aaccauggaa gcuguacgca ugggguagca gacuaguggu 60
uagaggagac ccucuccaag acacaacgca gcagcggggc ccaagacuag agguuagagg 120
agaccccccg cacaacaaca aacagcauau ugacgcuggg agagaccaga gaucucugc 180
ucucuacagc aucauuccag gcacagaacg ccagaaaug gauggugcu guugaaucaa 240
cagguucu 248

```

```

<210> SEQ ID NO 7
<211> LENGTH: 251
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Dengue virus construct

```

```

<400> SEQUENCE: 7

```

```

uaaaaaaacc gggaggccac aaaccaugga agcuguacgc auggcguagu ggacuagcgg 60
uuagaggaga cccucuccuu acagaucgca gcaacaauug gggccaaga cuagagguua 120
gaggagaccc ccccaaaaca aaaaacagca uauugacgcu gggaaagacc agagaucug 180
cugucuccuc agcaucauuc caggcacagg acgccagaaa auggaauggu gcuguugaau 240

```

-continued

caacagguuc u 251

<210> SEQ ID NO 8
 <211> LENGTH: 245
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 8

uaaaaccugg gaggcugcaa acuguggaag cuguacgcac gguguagcag acuagcgguu 60
 agaggagacc ccucccauga cacaacgcag cagcggggcc caagacuaga gguuagagga 120
 gacccccgc aaauaaaaac agcauauuga cgcugggaga gaccagagau ccugcugucu 180
 ccucagcauc auuccaggca cagaacgcca gaaauggaa uggugcuguu gaaucaacag 240
 guucu 245

<210> SEQ ID NO 9
 <211> LENGTH: 250
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 9

aucacccagg aggccaugcg ccacggaagc uguacgcgug gcuaauugga cuagcgguaa 60
 gaggagacc cucccaucac ugacaaaacg cagcaaaagg gggccaaga cuagagguua 120
 gaggagacc cccaacaca aaaacagcau auugacgcug ggaaagacca gagauccugc 180
 ugucucugca acaucaaacc aggcacagag cgccgcaaga uggauuggug uuguugaacc 240
 aacagguucu 250

<210> SEQ ID NO 10
 <211> LENGTH: 217
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 10

uaaaaacccg ggaggcugca acuagugguu agaggagacc ccucccaaga cacaacgcag 60
 cagcggggcc caagacuaga gguuagagga gacccccgc acaacaaca acagcauau 120
 gacgcuggga gagaccagag auccugcugu cucuacagca ucauuccagg cacagaacgc 180
 cagaaaugg auggugcug uugaaucaac agguucu 217

<210> SEQ ID NO 11
 <211> LENGTH: 220
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 11

uaaaaaaacc gggaggccac aaauagcgg uagaggagac ccucuccua cagaucgcag 60
 caacaauagg ggccaagac uagagguuag aggagacccc ccaaaaaca aaaacagcau 120
 auugacgcug ggaaagacca gagauccugc ugucuccuca gcaucauucc aggcacagga 180
 cgccagaaaa uggauggug cuguugaauc aacagguucu 220

-continued

<210> SEQ ID NO 12
 <211> LENGTH: 214
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 12

uaaaaccugg gaggcugcga cuagcgguaa gaggagacc cuccaugac acaacgcagc 60
 agcggggccc aagacuagag guuagaggag acccccgcga aaauaaaaca gcgauuugac 120
 gcugggagag accagagauc cugcugucuc cucagcauca uuccaggcac agaacgccag 180
 aaaauggaau ggugcuguug aaucaacagg uucu 214

<210> SEQ ID NO 13
 <211> LENGTH: 219
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 13

aucuccaggg aggccaugcg ccacgguuag aggagacccc uccaucacu gacaaaacgc 60
 agcaaaaggg ggccaagac uagagguuag aggagacccc cccaacacaa aaacagcaua 120
 uugacgcugg gaaagaccag agauccugcu gucucugcaa caucaauca ggcacagagc 180
 gccgaagau ggauuggugu uguugaacca acagguucu 219

<210> SEQ ID NO 14
 <211> LENGTH: 194
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 14

uaaaaaccg ggaggcugca aaccauggaa gcuguacgca uggguagca gacuagaggu 60
 uagaggagac ccccgccaca acaacaaaca gcgauuugac gcugggagag accagagauc 120
 cugcugucuc uacagcauca uuccaggcac agaacgccag aaaauggaau ggugcuguug 180
 aaucaacagg uucu 194

<210> SEQ ID NO 15
 <211> LENGTH: 196
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 15

uaaaaaucc gggaggccac aaaccaugga agcuguacgc auggcguagu ggagacuaga 60
 gguuagagga gacccccca aaacaaaaa cagcauuug acgcugggaa agaccagaga 120
 uccugcuguc uccucagcau cauuccaggc acaggacgcc agaaaugga auggugcugu 180
 ugaaucaaca gguucu 196

<210> SEQ ID NO 16
 <211> LENGTH: 190
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Dengue virus construct

-continued

<400> SEQUENCE: 16

uaaaaccugg gaggcugcaa acuguggaag cuguacgcac gguguagcga cuagagguua 60
 gaggagaccc cccgcaaaua aaaacagcau auugacgcug ggagagacca gagaucugc 120
 ugucuccuca gcaucauucc aggcacagaa cgccagaaaa uggauggug cuguugaauc 180
 aacagguucu 190

<210> SEQ ID NO 17

<211> LENGTH: 195

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Dengue virus construct

<400> SEQUENCE: 17

auccccaggg aggccaugcg ccacggaagc uguacgcgug gcgauuugga cuagacuaga 60
 gguuagagga gacccccca acacaaaaac agcauuuga cgcugggaaa gaccagagau 120
 ccugcugucu cugcaaacuc aauccaggca cagagcgccg caagauggau ugguguuguu 180
 gauccaacag guucu 195

<210> SEQ ID NO 18

<211> LENGTH: 39

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Junction region

<400> SEQUENCE: 18

gagggagcca tttggtaaac gtaggaagtg aaaaagagg 39

<210> SEQ ID NO 19

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Junction region

<400> SEQUENCE: 19

Glu Gly Ala Ile Trp
1 5

<210> SEQ ID NO 20

<211> LENGTH: 39

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Junction region

<400> SEQUENCE: 20

agtgaaggag ttctgtaatt accaacaaca aacaccaaaa 39

<210> SEQ ID NO 21

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Junction region

<400> SEQUENCE: 21

Ser Glu Gly Val Leu
1 5

-continued

<210> SEQ ID NO 22
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Junction region

<400> SEQUENCE: 22

gagggagcca tttggtagtt aacaacaaca aacaccaaaa 39

<210> SEQ ID NO 23
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Junction region

<400> SEQUENCE: 23

Glu Gly Ala Ile Trp
1 5

<210> SEQ ID NO 24
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 24

taaaaacagc atattgacgc tgggag 26

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 25

gactagaggt tagaggagac 20

<210> SEQ ID NO 26
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 26

gactagcggg tagaggagac ccc 23

<210> SEQ ID NO 27
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 27

tcgggccccg ctgctgcgtt g 21

<210> SEQ ID NO 28
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

```

<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 28

ttgggccccg ctgctgcgtt g                               21

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 29

tgtgtcatgg gaggggtctc                               20

<210> SEQ ID NO 30
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 30

gctacaccgt gcgtacagct tcc                            23

<210> SEQ ID NO 31
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 31

gcagcctccc aggttttacg tcc                            23

<210> SEQ ID NO 32
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 32

aacaacaaca aacaccaaag gctattg                       27

<210> SEQ ID NO 33
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 33

cctaccggta ccagaacctg ttg                            23

```

What is claimed is:

1. A nucleic acid encoding a dengue virus or chimeric dengue virus comprising

a Δ 30 mutation that removes the TL-2 homologous structure in DEN1 or DEN3, and nucleotides additional to the Δ 30 mutation deleted from the 3'-UTR that removes one or more nucleotides in the 5' direction up to and including nucleotide 258 with reverse-direction numbering.

2. The nucleic acid encoding a dengue virus or chimeric dengue virus of claim 1 wherein the mutation removes the

TL-2 homologous structure and removes the additional nucleotides in a contiguous manner contiguous to the Δ 30 mutation.

60 3. The nucleic acid encoding a dengue virus or chimeric dengue virus of claim 2 wherein the mutation is the Δ 86 mutation, such that the Δ 86 mutation deletes nucleotides from about 228 to about 145 of DEN1, or nucleotides from about 228 to about 143 of DEN3, designated with the reverse-direction numbering system.

65 4. The nucleic acid encoding a dengue virus or chimeric dengue virus of claim 3 wherein the serotype is DEN3.

37

5. The nucleic acid encoding a dengue virus or chimeric dengue virus of claim 1 wherein the mutation removes both the TL-2 homologous structure and the additional nucleotides in a noncontiguous manner noncontiguous to the $\Delta 30$ mutation.

6. The nucleic acid encoding a dengue virus or chimeric dengue virus of claim 5 wherein the mutation is the $\Delta 30/31$ mutation, such that the $\Delta 30$ mutation deletes nucleotides from about 174 to about 145 of DEN 1, or nucleotides from about 173 to about 143 of DEN3, designated with the reverse-order numbering system, and the $\Delta 31$ mutation deletes nucleotides from about 258 to about 228 of DEN1 or nucleotides from about 258 to about 228 of DEN3, designated with the reverse-order numbering system.

7. The nucleic acid encoding a dengue virus or chimeric dengue virus of claim 6 wherein the serotype is DEN3.

8. An immunogenic composition comprising a nucleic acid encoding a dengue virus or chimeric dengue virus according to claim 1 or a dengue virus or chimeric dengue virus comprising said nucleic acid.

9. A method of inducing an immune response to a dengue virus in a patient comprising administering the immunogenic composition of claim 8 to a patient to induce an immune response to a dengue virus.

10. A method of producing a nucleic acid encoding a dengue virus or chimeric dengue virus comprising introducing a

38

mutation into the 3' untranslated region (3'-UTR) of DEN1 or DEN3, wherein the mutation is

a $\Delta 30$ mutation that removes the TL-2 homologous structure in DEN1 or DEN3, and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR that removes one or more nucleotides in the 5' direction up to and including nucleotide 258 with reverse-direction numbering.

11. A dengue virus or chimeric dengue virus comprising the nucleic acid encoding the dengue virus or chimeric dengue virus of claim 1.

12. A dengue virus or chimeric dengue virus comprising the nucleic acid encoding the dengue virus or chimeric dengue virus of claim 3.

13. A dengue virus or chimeric dengue virus comprising the nucleic acid encoding the dengue virus or chimeric dengue virus of claim 6.

14. A tetravalent immunogenic composition comprising nucleic acids encoding each of DEN1, DEN2, DEN3, and DEN4, wherein each nucleic acid has a $\Delta 30$ mutation that removes the TL-2 homologous structure and nucleotides additional to the $\Delta 30$ mutation deleted from the 3'-UTR that removes one or more nucleotides in the 5' direction up to and including nucleotide 258 with reverse-direction numbering.

* * * * *