

US00RE46292E

(19) **United States**
 (12) **Reissued Patent**
Brinker et al.

(10) **Patent Number: US RE46,292 E**
 (45) **Date of Reissued Patent: Jan. 31, 2017**

(54) **SOYBEAN TRANSGENIC EVENT MON 87708 AND METHODS OF USE THEREOF**

(71) Applicant: **Monsanto Technology LLC**, St. Louis, MO (US)

(72) Inventors: **Ronald J. Brinker**, Ellisville, MO (US); **Wen C. Burns**, Chesterfield, MO (US); **Paul C. C. Feng**, Wildwood, MO (US); **Anju Gupta**, Ankeny, IA (US); **Sio Wai Hoi**, St. Louis, MO (US); **Marianne Malven**, Ellisville, MO (US); **Kunsheng Wu**, Ballwin, MO (US)

(73) Assignee: **Monsanto Technology LLC**, St. Louis, MO (US)

(21) Appl. No.: **14/745,241**

(22) Filed: **Jun. 19, 2015**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **8,501,407**
 Issued: **Aug. 6, 2013**
 Appl. No.: **12/868,989**
 Filed: **Aug. 26, 2010**

U.S. Applications:

(60) Provisional application No. 61/243,227, filed on Sep. 17, 2009.

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
C07H 21/02 (2006.01)
C07H 21/04 (2006.01)
C12P 19/34 (2006.01)
C12N 15/82 (2006.01)
A23D 9/00 (2006.01)

(52) **U.S. Cl.**
 CPC **C12N 15/8274** (2013.01); **A23D 9/00** (2013.01); **A23L 11/03** (2016.08); **C12Q 1/6895** (2013.01)

(58) **Field of Classification Search**
 CPC **A23D 9/00**
 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,850,019 A	12/1998	Maiti et al.	
5,914,451 A	6/1999	Martinell et al.	
7,022,896 B1 *	4/2006	Weeks et al.	800/300
7,241,567 B2 *	7/2007	Weyens et al.	435/6.15
7,838,729 B2	11/2010	Feng et al.	
7,851,670 B2	12/2010	Wan et al.	
7,855,326 B2	12/2010	Feng et al.	
7,884,262 B2	2/2011	Clemente et al.	
7,939,721 B2	5/2011	Arnevik et al.	
8,084,666 B2	12/2011	Feng et al.	
8,207,092 B2	6/2012	Bhatti et al.	
8,420,888 B2	4/2013	Feng et al.	
8,629,328 B2	1/2014	Feng et al.	
RE44,971 E	6/2014	Wan et al.	
8,754,011 B2	6/2014	Bhatti et al.	
RE45,048 E	7/2014	Feng et al.	

8,791,325 B2	7/2014	Feng et al.
8,999,411 B2	4/2015	Froman et al.
2003/0115626 A1	6/2003	Weeks et al.
2004/0117870 A1	6/2004	Weyens et al.
2008/0015110 A1	1/2008	Clemente et al.
2008/0119361 A1	5/2008	Feng et al.
2008/0120739 A1	5/2008	Wan et al.
2008/0166699 A1	7/2008	Baley et al.
2008/0305952 A1	12/2008	Arnevik et al.
2009/0081760 A1	3/2009	D'Ordine et al.
2009/0199308 A1	8/2009	Duff et al.
2011/0067134 A1	3/2011	Brinker et al.
2011/0067141 A1	3/2011	Froman et al.
2011/0245080 A1	10/2011	Arnevik et al.
2011/0302667 A1	12/2011	Brown et al.
2013/0217130 A1	8/2013	Bhatti et al.
2013/0217576 A1	8/2013	Bhatti et al.
2013/0244875 A1	9/2013	Feng et al.
2014/0315717 A1	10/2014	Bhatti et al.
2014/0373190 A1	12/2014	Brown et al.
2015/0119248 A1	4/2015	Brown et al.

FOREIGN PATENT DOCUMENTS

CL	83-08	1/2008
CN	101460626 A	6/2009
CN	101501196 A	8/2009
CN	101636498 A	1/2010
UA	A200810227	12/2008
WO	WO 98/45424	10/1998
WO	WO 2005/061720	7/2005

(Continued)

OTHER PUBLICATIONS

English translation of Office Action regarding Japanese Patent Application No. 2014-235817, dated Dec. 8, 2016.
 Yamauchi et al., Series "Food Science," *Soybean Science* 1(5):188-194, 1997.
 USPTO: Non-Final Office Action regarding U.S. Appl. No. 13/945,741, dated Jan. 12, 2016.
 Hohe et al., "A tool for understanding homologous recombination in plants," *Plant Cell Rep.* 21:1135-1142, 2003.
 Response to Non-Final Office Action regarding U.S. Appl. No. 13/945,741, dated Feb. 18, 2016.
 U.S. Appl. No. 14/179,524, filed Feb. 12, 2014, Froman et al.
 U.S. Appl. No. 14/333,479, filed Jul. 16, 2014, Brown et al.
 Alignment of SEQ ID No. 1 from U.S. Appl. No. 12/865,844 and SEQ ID Nos. 1 and 7 from U.S. Appl. No. 13/945,741, created Aug. 27, 2013.
 Alignment of SEQ ID No. 1 from U.S. Appl. No. 13/151,082 and SEQ ID No. 1 from U.S. Appl. No. 13/945,741, created Aug. 27, 2013.

(Continued)

Primary Examiner — Bruce Campell
 (74) *Attorney, Agent, or Firm* — Dentons US LLP;
 Pamela Sisson

(57) **ABSTRACT**

The invention provides a transgenic soybean event MON 87708 plant and plants, plant cells, seeds, plant parts, and commodity products derived from event MON 87708. The invention also provides polynucleotides specific for event MON 87708 and plants, plant cells, seeds, plant parts, and commodity products comprising polynucleotides specific for event MON 87708. The invention also provides methods related to event MON 87708.

21 Claims, 1 Drawing Sheet

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO 2007/143690	12/2007
WO	WO 2007/146706	12/2007
WO	WO 2008/048964	4/2008
WO	WO 2008/105890	9/2008
WO	WO 2009/102873	8/2009
WO	WO 2009/126462	10/2009

OTHER PUBLICATIONS

Anklam et al., "Analytical methods for detection and determination of genetically modified organisms in agricultural crops and plant-derived food products," *Eur. Food Res Technol.* 214:3-26, 2001.

Astaurov et al., *Great Soviet Encyclopedia* (in Russian), article "Biology," 1970.

Berdal et al., "Roundup Ready® soybean event-specific real-time quantitative PCR assay and estimation of the practical detection and quantification limits in GMO analyses," *Eur. Food Res Technol.* 213:432-438, 2001.

Coruzzi et al., "Tissue-specific and light-regulated expression of a pea nuclear gene encoding the small subunit of ribulose-1,5-bisphosphate carboxylase," *The EMBO Journal* 3(8):1671-1679, 1984.

English translation of Chilean Application No. CL 83-08, filed Jan. 11, 2008.

English translation of p. 2 of Office Action issued in Ukrainian Application No. A 2012 04680, dated Jan. 26, 2015.

English translation of pp. 6-7 of Office Action issued in Russian Application No. 2012115127, dated Aug. 13, 2014.

GenBank Accession No. AY786443, dated Jun. 29, 2005.

Gruzdev, "Chemical protection of plants," *Agropromizdat* (in Russian), 3rd edition, p. 316, 1987.

Herman et al., "A Three-component Dicamba O-Demethylase from *Pseudomonas maltophilia*, Strain DI-6," *The Journal of Biological Chemistry* 280(26):24759-24767, 2005.

International Preliminary Report on Patentability PCT/US2007/070517 (WO 2007/143690), Dec. 10, 2008.

International Preliminary Report on Patentability PCT/US2009/033930 (WO 2009/102873), Aug. 17, 2010.

International Preliminary Report on Patentability PCT/US2010/046759 (WO 2011/034704), Mar. 3, 2012.

International Search Report for PCT/US2010/046759, dated Dec. 9, 2010.

NCBI GenBank Accession No. AC207600.4, "Medicago truncatula chromosome 2 BAC clone mth2-31i16, complete sequence," dated Oct. 27, 2007.

NCBI GenBank Accession No. ET721878.1, "CHO_OF124xh18fl.abl CHO_OF *Nicotiana tabacum* genom 5-, genomic survey sequence," dated Jun. 20, 2008.

Prohorov et al., *Great Soviet Encyclopedia* (in Russian), 16:233, 1974.

Rychlik et al., "A computer program for choosing optimal oligonucleotides for filter hybridization, sequencing and in vitro amplification of DNA," *Nucl. Acids Res* 17(21):8543-51, 1989.

Sequence alignment information as indicated in transmittal letter of Nov. 25, 2013 Information Disclosure Statement for U.S. Appl. No. 13/945,741.

USPTO: Notice of Allowance for U.S. Appl. No. 13/151,082, issued Jun. 17, 2014.

Wang et al., "Characterization of cellular and enzymatic degradation of dicamba by *Pseudomonas maltophilia*, strain DI-6," dissertation presented to the Faculty of the Graduate College at the University of Nebraska, 1996.

Zimmermann et al., "Event Specific Transgene Detection in Bt11 Corn by Quantitative PCR at the Integration Site," *Lebensm.-Wiss. u.-Technol.* 33(3):210-216, 2000.

USPTO: Notice of Allowance and Feed Due regarding U.S. Appl. No. 13/945,741, dated May 12, 2016.

NCBI GenBank Accession No. EI363554 (2007, submitted by Jackson et al. Purdue University).*

New England BioLabs Inc., 1998/1999 Catalog, Nucleic Acids, Linkers and Primers, pp. 121 and 284.

Final Office Action regarding U.S. Appl. No. 11/801,114, dated Aug. 26, 2009.

Behrens et al., "Dicamba resistance: enlarging and preserving biotechnology-based weed management strategies," *Science*, 316:1185-1188, 2007.

Carter et al., "Evaluating weed control options in dicamba tolerant soybean," *North Central Weed Science Society Proc.*, Abstract and Poster, 63:25, 2008.

Carter et al., "Resistant weed management using dicamba tolerant soybean," Weed Science Society of America Annual Meeting, Orlando, FL, Feb. 9-13, 2009.

Carter et al., "Evaluation of dicamba tolerant soybean in Kentucky: a three-year review," Weed Science Society of America Annual Meeting, Denver, CO, Feb. 7-11, 2010.

Clarity herbicide EPA Reg. No. 7969-137, BASF, for use on dicamba-tolerant soybean MON 87708—grown for research, field trials, and seed production only, accepted with comments in EPA letter dated Feb. 13, 2008, Supplemental Label.

Clarity herbicide EPA Reg. No. 7969-137, BASF, For use on dicamba-tolerant soybean MON 87708—grown for seed production only, Feb. 26, 2009, Supplemental Label.

Clarity herbicide, EPA Reg. No. 7969-137, BASF, For use on dicamba-tolerant soybean MON 87708—grown for research, field trials, and seed production only, including USDA regulated plantings or seed production, accepted with comments in EPA letter dated May 28, 2010, Supplemental Label.

GenBank Accession No. CW906303, dated Dec. 8, 2004.


GenBank Accession No. EI363554, dated Feb. 16, 2007.

Hayes et al., "Glyphosate resistant horseweed control in dicamba-glyphosate resistant soybeans," *Southern Weed Science Society Proceedings*, 61:51, 2008, Abstract.

Steckel et al., "Glyphosate-resistant horseweed control in dicamba glyphosate resistant soybeans," *North Central Weed Science Society Proc.*, 63:92, 2008.

Steckel et al., "Glyphosate-resistant horseweed control in soybean tolerant to both dicamba and glyphosate," *North Central Weed Science Society Proceedings*, 62:178, 2007.

* cited by examiner



**SOYBEAN TRANSGENIC EVENT MON 87708
AND METHODS OF USE THEREOF**

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue; a claim printed with strikethrough indicates that the claim was canceled, disclaimed, or held invalid by a prior post-patent action or proceeding.

CROSS REFERENCE TO RELATED
APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 61/243,227 filed Sep. 17, 2009, which is herein incorporated by reference in its entirety.

INCORPORATION OF SEQUENCE LISTING

The sequence listing that is contained in the file named "55544-0001_seqlisting.txt", which is 19.5 kilobytes (size as measured in Microsoft Windows®) and was created on Aug. 13, 2010, is filed herewith by electronic submission and is incorporated by reference herein.

FIELD OF THE INVENTION

The invention relates to transgenic Glycine max event MON 87708. The event exhibits tolerance to dicamba herbicide. The invention also relates to plants, plant parts, plant seeds, plant cells, agricultural products, and methods related to event MON 87708 and provides nucleotide molecules that are unique to the event and were created in connection with the insertion of transgenic DNA into the genome of a Glycine max plant.

BACKGROUND OF THE INVENTION

Soybean (Glycine max) is an important crop in many areas of the world, and the methods of biotechnology have been applied to this crop in order to produce soybean with desirable traits. One such desirable trait is herbicide tolerance. The expression of an herbicide tolerance transgene in a plant can confer the desirable trait of herbicide tolerance on the plant, but expression of the transgene may be influenced by the chromosomal location and the genomic result of the transgene insertion. For example, it has been observed in plants that there often is variation in the level and pattern of transgene expression among individual events that differ in the chromosomal insertion site of the transgene but are otherwise identical. There may also be undesirable and/or desirable phenotypic or agronomic differences between events. Because of this, it is often necessary to produce and analyze a large number of individual plant transformation events in order to select an event having both the desirable trait and the optimal phenotypic and agricultural characteristics necessary to make it suitable for commercial purposes. Such selection often requires greenhouse and field trials with many events over multiple years, in multiple locations, and under a variety of conditions so that a significant amount of agronomic, phenotypic, and molecular data may be collected. The resulting data and observations must then be analyzed by teams of scientists and agronomists with the goal of selecting a commercially suitable event. Such an event, once selected, may then be used for introgressing the desirable trait into other genetic backgrounds using plant

breeding methods, and thus producing a number of different crop varieties that contain the desirable trait and are suitably adapted to specific local growing conditions.

SUMMARY OF THE INVENTION

The invention provides transgenic soybean plants designated event MON 87708, which exhibit commercially acceptable tolerance to applications of dicamba herbicide, having representative seed deposited with American Type Culture Collection (ATCC) with Accession No. PTA-9670. The invention also provides novel DNA molecules related to soybean event MON 87708 and methods of using these molecules. The invention also provides seeds, progeny, plant parts, cells, and commodity products of soybean event MON 87708. The invention also provides methods of using soybean event MON 87708 and methods of producing dicamba tolerant soybean.

The invention provides recombinant DNA molecules related to soybean event MON 87708. These recombinant DNA molecules may comprise nucleotide molecules having a nucleotide sequence representing a region of the genomic DNA flanking the transgene insertion, and/or a region of the transgene insertion, and/or a contiguous sequence of any of these regions such as a region of the junction between the transgene insertion and flanking genomic DNA of soybean event MON 87708. The invention also provides DNA molecules useful as primers and probes diagnostic for soybean event MON 87708 and amplicons diagnostic for the presence of soybean event MON 87708. Soybean plants, plant cells, plant parts, commodity products, progeny, and seeds comprising these molecules are also disclosed.

The invention provides methods, compositions, and kits useful for detecting the presence and/or absence of DNA derived from soybean event MON 87708 and thus the presence and/or absence of the event. The invention provides a method for detection of MON 87708 by contacting a sample comprising DNA with a primer set that when used in a nucleic acid amplification reaction with genomic DNA from soybean event MON 87708 produces an amplified DNA diagnostic for soybean event MON 87708, performing a nucleic acid amplification reaction thereby producing the amplified DNA, and detecting the presence and/or absence of the amplified DNA. The invention also provides a method for detection of MON 87708 by contacting a sample comprising DNA with a probe that when used in a hybridization reaction with DNA from soybean event MON 87708 hybridizes to a DNA molecule specific for soybean event MON 87708, performing a hybridization reaction, and detecting the hybridization of the probe to the DNA molecule. Kits comprising the methods and compositions of the invention useful for detecting the presence of DNA derived from soybean event MON 87708 are also provided.

The invention provides a soybean plant, seed, plant cell, progeny plant, plant part, or commodity product derived from a plant, plant cell, or seed of soybean event MON 87708. The invention also provides a soybean plant, seed, plant cell, progeny plant, plant part, or commodity product comprising a recombinant DNA molecule having a nucleotide sequence selected from the group consisting of SEQ ID NO: 1-8, and complements and fragments thereof. The invention also provides a soybean plant, seed, plant cell, progeny plant, plant part, or commodity product derived from the plant or seed of soybean event MON 87708 and comprising a recombinant DNA molecule that produces an

amplified DNA molecule comprising SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and/or SEQ ID NO: 8 in a DNA amplification method.

The invention provides a method for controlling weeds in a field by planting soybean event MON 87708 and then applying an effective dose of dicamba herbicide capable of controlling the weeds without injuring the soybean event MON 87708 plants. The invention also provides a method for controlling weeds in a field by applying an effective dose of dicamba herbicide to control weeds in a field and then planting soybean event MON 87708 in the field. The invention also provides a method for producing soybean seed essentially free of the seeds of toxic weed species by planting seeds of a dicamba tolerant soybean variety MON 87708 in a field, applying a post-emergence effective dose of dicamba herbicide sufficient to kill the toxic weed species to the field, and harvesting seed from the field.

The invention provides methods of producing a soybean plant and/or seed that tolerates application of dicamba herbicide by sexually crossing a soybean event MON 87708 plant comprising SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and/or SEQ ID NO: 8 with a second soybean plant, thereby producing seed, growing the seed to produce progeny plants, treating the progeny plants with dicamba, and selecting a progeny plant that is tolerant to dicamba. The methods may also include selfing the selected progeny plant to produce a plurality of second generation progeny plants and selecting from these a dicamba tolerant plant. The methods may also include sexually crossing the selected progeny plant with another soybean plant to produce seed, growing the seed to produce a second generation of progeny plants, treating the second generation of progeny plants with dicamba, and selecting a second generation progeny plant that is tolerant to dicamba. The invention provides methods of producing a soybean plant and/or seed that tolerates application of dicamba herbicide by selfing a dicamba tolerant soybean event MON 87708 plant comprising SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and/or SEQ ID NO: 8, thereby producing seed, growing the seed to produce progeny plants, treating the progeny plants with dicamba; and selecting a progeny plant that is tolerant to dicamba.

The invention provides methods of determining the zygosity of a soybean event MON 87708 plant or seed comprising contacting a soybean DNA sample with a primer set comprising SEQ ID NO: 12, SEQ ID NO: 13, and SEQ ID NO: 14 and a probe set comprising SEQ ID NO: 15 and SEQ ID NO: 16; then performing a nucleic acid amplification reaction with the sample, primer set, and probe set; then detecting in then nucleic acid amplification reaction a first fluorescent signal that is diagnostic for event MON 87708 and a second fluorescent signal different from the first fluorescent signal and that is diagnostic for native soybean genomic DNA corresponding to the location of insertion of the event MON 87708 transgene; and analyzing the presence and/or absence of the first fluorescent signal and the second fluorescent signal in the nucleic acid amplification reaction, wherein the presence of both fluorescent signals indicates the sample is heterozygous for event MON 87708 and the presence of only the first fluorescent signal indicates the sample is homozygous for event MON 87708.

The invention also provides a soybean plant, seed, plant cell, or plant part comprising soybean haplotype region on linkage group 9 at approximately map position 143.5 comprising a dicamba tolerance gene and further defined by haplotype window 19743 and 19767, and methods of using

the same. The foregoing and other aspects of the invention will become more apparent from the following detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates the organization of the transgenic insert in the genome of soybean event MON 87708; [A] corresponds to the relative position of SEQ ID NO: 1, which is sixty nucleotides of the junction between the soybean genomic DNA and the 5' portion of the transgene insert DNA; [A'] corresponds to the relative position of SEQ ID NO: 7, which is one hundred nucleotides of the junction between the soybean genomic DNA and the 5' portion of the transgene insert DNA; [B] corresponds to the relative position of SEQ ID NO: 2, which is sixty nucleotides of the junction between the soybean genomic DNA and the 3' portion of the transgene insert DNA; [B'] corresponds to the relative position of SEQ ID NO: 8, which is one hundred nucleotides of the junction between the soybean genomic DNA and the 3' portion of the transgene insert DNA; [C] corresponds to the relative position of SEQ ID NO: 3, which is the soybean genome sequence flanking the arbitrarily assigned/designated 5' end of the expression cassette integrated into the genome in event MON 87708; [D] corresponds to the relative position of SEQ ID NO: 4, which is the soybean genome sequence flanking the arbitrarily assigned/designated 3' end of the expression cassette integrated into the genome in event MON 87708; [E] represents the various elements comprising SEQ ID NO: 5 and is the sequence of the expression cassette inserted into the genome of the event MON 87708; and [F] represents the contiguous sequence (provided as SEQ ID NO: 6) comprising, as represented in the FIGURE from left to right, SEQ ID NO: 3, SEQ ID NO: 5 and SEQ ID NO: 4, in which SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and SEQ ID NO: 8 are included, as these sequences are present in the genome in event MON 87708.

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO: 1 is a sixty nucleotide sequence representing the 5' junction between the soybean genomic DNA and the integrated transgenic expression cassette. SEQ ID NO: 1 is positioned in SEQ ID NO: 6 at nucleotide position 1097-1156.

SEQ ID NO: 2 is a sixty nucleotide sequence representing the 3' junction between the soybean genomic DNA and the integrated transgenic expression cassette. SEQ ID NO: 2 is positioned in SEQ ID NO: 6 at nucleotide position 4100-4159.

SEQ ID NO: 3 is the 5' sequence flanking the inserted DNA of soybean event MON 87708 up to and including a region of transgene DNA insertion.

SEQ ID NO: 4 is the 3' sequence flanking the inserted DNA of soybean event MON 87708 up to and including a region of transgene DNA insertion.

SEQ ID NO: 5 is the sequence of the integrated transgenic expression cassette.

SEQ ID NO: 6 is the nucleotide sequence representing the contig of the 5' sequence flanking the inserted DNA of soybean event MON 87708 (SEQ ID NO: 3), the sequence of the inserted DNA (SEQ ID NO: 5), and the 3' sequence flanking the inserted DNA of soybean event MON 87708 (SEQ ID NO: 4) and includes SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and SEQ ID NO: 8.

SEQ ID NO: 7 is a one hundred nucleotide sequence representing the 5' junction between the soybean genomic DNA and the integrated transgenic expression cassette.

SEQ ID NO: 8 is a one hundred nucleotide sequence representing the 3' junction between the soybean genomic DNA and the integrated transgenic expression cassette.

SEQ ID NO: 9 is the sequence of a primer referred to as Primer SQ13570 and used to identify soybean event MON 87708. It is complimentary to the inserted expression cassette at the region close to the 3' transgene insertion border. A PCR amplicon produced from a TAQMAN® (PE Applied Biosystems, Foster City, Calif.) assay using the combination of primers SQ13570 and SQ13571 (SEQ ID NO: 10) is a positive result for the presence of the event MON 87708.

SEQ ID NO: 10 is the sequence of a primer referred to as Primer SQ13571 and used to identify soybean event MON 87708. It is complimentary to a 3' region flanking the inserted expression cassette and close to the transgene DNA insertion border. A PCR amplicon produced from a TAQMAN® (PE Applied Biosystems, Foster City, Calif.) assay using the combination of primers SQ13570 (SEQ ID NO: 9) and SQ13571 is a positive result for the presence of the event MON 87708.

SEQ ID NO: 11 is the sequence of a probe referred to as Probe PB4655 and used to identify soybean event MON 87708. It is complimentary to a region spanning the 3' junction of the inserted expression cassette and the genomic DNA. This probe is a 6-FAM™-labeled synthetic oligonucleotide. Release of a fluorescent signal in an amplification reaction using primers SQ13570 and SQ13571 (SEQ ID NO: 9-10) in combination with 6-FAM™-labeled probe PB4655 is diagnostic of event MON 87708 in a TAQMAN® assay.

SEQ ID NO: 12 is the sequence of a primer referred to as Primer SQ20632 and used to identify MON 87708 event zygosity.

SEQ ID NO: 13 is the sequence of a primer referred to as Primer SQ20636 and used to identify soybean wild-type and MON 87708 event zygosity.

SEQ ID NO: 14 is the sequence of a primer referred to as Primer SQ20637 and used to identify soybean wild-type zygosity.

SEQ ID NO: 15 is the sequence of a probe referred to as Probe PB10130 and used for a MON 87708 event zygosity assay.

SEQ ID NO: 16 is the sequence of a probe referred to as Probe PB10131 and used for a soybean wild-type zygosity assay.

DETAILED DESCRIPTION

The following definitions and methods are provided to better define the invention and to guide those of ordinary skill in the art in the practice of the invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art.

The invention provides a transgenic soybean event MON 87708 that exhibits commercially acceptable tolerance to applications of dicamba herbicide. The event comprises a single insertion of transgenic DNA into the chromosome/genome of the soybean germplasm. An "event" is produced by: (i) transformation of a plant cell with a nucleic acid construct that includes a transgene of interest, (ii) regeneration of a population of plants resulting from the insertion of the transgene into the genome of the plant, and (iii) selection of a particular plant characterized by insertion of the trans-

gene into a particular location in the plant's genome. The term "event" refers to the original transformant that includes the transgene inserted into the particular location in the plant's genome. The term "event" also refers to progeny of the transformant that include the transgene inserted into the particular location in the plant's genome. Such progeny may be produced by a sexual outcross between the transformant, or its progeny, and another plant. Such other plant may be a transgenic plant comprising the same or different transgene and/or a nontransgenic plant, such as one from a different variety. Even after repeated back-crossing to a recurrent parent, the inserted DNA and flanking DNA from the transformed parent is present in the progeny of the cross at the same genomic location.

As used herein, the term "soybean" means *Glycine max* and includes all plant varieties that can be bred with soybean, including wild soybean species as well as those plants belonging to *Glycine* that permit breeding between species.

The term "event" also refers to a DNA molecule from the original transformant comprising the inserted DNA and the flanking soybean genomic DNA immediately adjacent to either side of the inserted DNA. This DNA molecule is created by the act of inserting the transgenic DNA into the genome of the soybean plant, i.e., by the act of transformation. This DNA molecule therefore comprises a nucleotide sequence that is both specific to the event and that is unique to the genome of the soybean plant into which the transgenic DNA has been inserted, in that this nucleotide sequence contains both the sequence of a particular region of soybean genomic DNA and of the transgenic DNA insert. The arrangement of the inserted DNA in soybean event MON 87708 in relation to the surrounding soybean plant genome DNA is therefore specific and unique for soybean event MON 87708. This DNA molecule is also an integral part of the soybean chromosome of event MON 87708 and as such is static in the plant and may be passed on to progeny of the plant.

Event MON 87708 comprises a transgene that confers tolerance to applications of dicamba herbicide to the soybean plant. "Dicamba" refers to 3,6-dichloro-2-methoxybenzoic acid. Dicamba is a synthetic auxin herbicide useful for controlling broadleaf weeds. Soybean plants were transformed with dicamba mono-oxygenase (DMO), an enzyme cloned from *Stenotrophomonas maltophilia* which is commonly found in soil rhizosphere. Dicamba mono-oxygenase is an enzyme that catalyzes the deactivation of dicamba via an O-demethylation reaction to the nonherbicidal compound 3,5-dichlorosalicylic acid. In some areas of the world toxic weed species seeds may contaminate harvested soybean seeds that can affect the health and nutrition of animals fed the contaminated soybean commodity products. These plants can be eliminated from a soybean field by treatment with a dicamba herbicide. Members of this group of toxic weeds include *Cardaria* spp, *Heliotropium* spp, *Centaurea* spp., *Senecio* spp., *Crotalaria* spp., *Solanum* spp., *Xanthium* spp., *Amsinckia* spp., *Cassia* spp., *Sesbania* spp., *Datura* spp., *Ricinus* spp., *Argemone* spp., *Corchorus* spp., *Ipomoea* spp., and *Echium* spp.

As used herein, the term "recombinant" refers to a form of DNA and/or protein and/or an organism that would not normally be found in nature and as such was created by human intervention. Such human intervention may produce a recombinant DNA molecule and/or a recombinant plant. As used herein, a "recombinant DNA molecule" is a DNA molecule comprising a combination of DNA molecules that would not naturally occur together and is the result of human intervention, e.g., a DNA molecule that is comprised of a

combination of at least two DNA molecules heterologous to each other, and/or a DNA molecule that is artificially synthesized and comprises a polynucleotide sequence that deviates from the polynucleotide sequence that would normally exist in nature, and/or a DNA molecule that comprises a transgene artificially incorporated into a host cell's genomic DNA and the associated flanking DNA of the host cell's genome. An example of a recombinant DNA molecule is a DNA molecule described herein resulting from the insertion of the transgene into the soybean genomic DNA, which may ultimately result in the expression of a recombinant RNA and/or protein molecule in that organism. As used herein, a "recombinant plant" is a plant that would not normally exist in nature, is the result of human intervention, and contains a transgene and/or heterologous DNA molecule incorporated into its genome. As a result of such genomic alteration, the recombinant plant is distinctly different from the related wildtype plant. An example of a recombinant plant is a soybean plant described herein as Event MON 87708.

As used herein, the term "transgene" refers to a nucleotide molecule artificially incorporated into a host cell's genome. Such transgene may be heterologous to the host cell. The term "transgenic plant" refers to a plant comprising such a transgene.

As used herein, the term "heterologous" refers to a first molecule not normally found in combination with a second molecule in nature. For example, a molecule may be derived from a first species and inserted into the genome of a second species. The molecule would thus be heterologous to the host and artificially incorporated into a host cell's genome.

As used herein, the term "chimeric" refers to a single DNA molecule produced by fusing a first DNA molecule to a second DNA molecule, where neither first nor second DNA molecule would normally be found in that configuration, i.e., fused to the other. The chimeric DNA molecule is thus a new DNA molecule not otherwise normally found in nature.

The invention provides DNA molecules and their corresponding nucleotide sequences. As used herein, the term "DNA", "DNA molecule", "nucleotide molecule" refers to a DNA molecule of genomic or synthetic origin, i.e., a polymer of deoxyribonucleotide bases or a polynucleotide molecule, read from the 5' (upstream) end to the 3' (downstream) end. As used herein, the term "DNA sequence", "nucleotide sequence" or "polynucleotide sequence" refers to the nucleotide sequence of a DNA molecule. The nomenclature used herein is that required by Title 37 of the United States Code of Federal Regulations §1.822 and set forth in the tables in WIPO Standard ST.25 (1998), Appendix 2, Tables 1 and 3. By convention, the nucleotide sequences of the invention provided as SEQ ID NO: 1-8 and fragments thereof are disclosed with reference to only one strand of the two complementary nucleotide sequence strands. By implication, the complementary sequences (i.e. the sequences of the complementary strand), also referred to in the art as the reverse complementary sequences, are within the scope of the invention and are expressly intended to be within the scope of the subject matter claimed.

The nucleotide sequence corresponding to the complete nucleotide sequence of the inserted transgenic DNA and substantial segments of the soybean genome DNA flanking either end of the inserted transgenic DNA is provided herein as SEQ ID NO: 6. A subsection of this is the inserted transgenic DNA provided as SEQ ID NO: 5. The nucleotide sequence of the soybean genome DNA physically linked by phosphodiester bond linkage to and therefore flanking the 5' end of the inserted transgenic DNA is set forth as shown in

SEQ ID NO: 3. The nucleotide sequence of the soybean genome DNA physically linked by phosphodiester bond linkage to and therefore flanking the 3' end of the inserted transgenic DNA is set forth as shown in SEQ ID NO: 4.

The soybean event MON 87708 further comprises two regions, one spanning the 5' location and one spanning the 3' location where the transgenic DNA is inserted into the genomic DNA, referred to herein as the 5' and 3' junction, respectively. A "junction sequence" or "junction region" refers to the DNA sequence and/or corresponding DNA molecule that spans the inserted transgenic DNA and the adjacent flanking genomic DNA. The junction sequences may be arbitrarily represented by the two 60 nucleotide sequences provided as SEQ ID NO: 1 and SEQ ID NO: 2, each representing 30 nucleotides of the flanking genomic DNA adjacent to and contiguous with 30 nucleotides of insert DNA. Alternatively, the junction sequences may be arbitrarily represented by the two 100 nucleotide sequences provided as SEQ ID NO: 7 and SEQ ID NO: 8, each representing 50 nucleotides of the flanking genomic DNA adjacent to and contiguous with 50 nucleotides of insert DNA. These nucleotides are connected by phosphodiester linkage and in soybean event MON 87708 are present as part of the genome. In soybean the identification of one or more of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and SEQ ID NO: 8 in a sample derived from a soybean plant, seed, or plant part is determinative that the DNA was obtained from soybean event MON 87708 and is diagnostic for the presence in a sample of DNA from soybean event MON 87708. The invention thus provides a DNA molecule that contains at least the nucleotide sequence as set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and/or SEQ ID NO: 8. Any segment of DNA derived from transgenic soybean event MON 87708 that is sufficient to include SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and/or SEQ ID NO: 8 is within the scope of The invention. In addition, any polynucleotide comprising a sequence complementary to any of the sequences described within this paragraph is within the scope of the invention. FIG. 1 illustrates the physical arrangement of SEQ ID NO: 1-5 and 7-8 relative to SEQ ID NO: 6 arranged from 5' to 3'.

The invention provides exemplary DNA molecules that can be used either as primers or probes for diagnosing the presence of DNA derived from soybean plant event MON 87708 in a sample. Such primers or probes are specific for a target nucleic acid sequence and as such are useful for the identification of soybean event MON 87708 nucleic acid sequence by the methods of the invention described herein.

A "primer" is typically a highly purified, isolated polynucleotide that is designed for use in specific annealing or hybridization methods that involve thermal amplification. A pair of primers may be used with template DNA, such as a sample of soybean genomic DNA, in a thermal amplification, such as polymerase chain reaction (PCR), to produce an amplicon, where the amplicon produced from such reaction would have a DNA sequence corresponding to sequence of the template DNA located between the two sites where the primers hybridized to the template. As used herein, an "amplicon" is a piece or fragment of DNA that has been synthesized using amplification techniques. An amplicon of the invention comprises at least SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and/or SEQ ID NO: 8. A primer is typically designed to hybridize to a complementary target DNA strand to form a hybrid between the primer and the target DNA strand, and the presence of the primer is a point of recognition by a polymerase to begin extension of the primer (i.e., polymerization of additional nucleotides into a

lengthening nucleotide molecule) using as a template the target DNA strand. Primer pairs, as used in the invention, are intended to refer to use of two primers binding opposite strands of a double stranded nucleotide segment for the purpose of amplifying linearly the polynucleotide segment between the positions targeted for binding by the individual members of the primer pair, typically in a thermal amplification reaction or other conventional nucleic-acid amplification methods. Exemplary DNA molecules useful as primers are provided as SEQ ID NO: 9-10. The primer pair provided as SEQ ID NO: 9 and SEQ ID NO: 10 are useful as a first DNA molecule and a second DNA molecule that is different from the first DNA molecule, and both are each of sufficient length of contiguous nucleotides of either SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6 to function as DNA primers that, when used together in a thermal amplification reaction with template DNA derived from soybean event MON 87708, produce an amplicon comprising SEQ ID NO: 2.

A "probe" is an isolated nucleic acid that is complementary to a strand of a target nucleic acid. Probes according to the invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and the detection of such binding can be useful in diagnosing, discriminating, determining, or confirming the presence of that target DNA sequence in a particular sample. A probe may be attached to a conventional detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. An exemplary DNA molecule useful as a probe is provided as SEQ ID NO: 11.

Probes and primers according to the invention may have complete sequence identity with the target sequence, although primers and probes differing from the target sequence that retain the ability to hybridize preferentially to target sequences may be designed by conventional methods. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of transgenic DNA from soybean event MON 87708 in a sample. Probes and primers are generally at least about 11 nucleotides, at least about 18 nucleotides, at least about 24 nucleotides, or at least about 30 nucleotides or more in length. Such probes and primers hybridize specifically to a target DNA sequence under stringent hybridization conditions. Conventional stringency conditions are described by Sambrook et al., 1989, and by Haymes et al., In: *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, D.C. (1985). As used herein, two nucleic acid molecules are capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure. A nucleic acid molecule is the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are "minimally complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional "low-stringency" conditions. Similarly, the molecules are "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional "high-stringency" conditions.

Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure.

As used herein, the term "isolated" refers to at least partially separating a molecule from other molecules normally associated with it in its native or natural state. In one embodiment, the term "isolated" refers to a DNA molecule that is at least partially separated from the nucleic acids which normally flank the DNA molecule in its native or natural state. Thus, DNA molecules fused to regulatory or coding sequences with which they are not normally associated, for example as the result of recombinant techniques, are considered isolated herein. Such molecules are considered isolated even when integrated into the chromosome of a host cell or present in a nucleic acid solution with other DNA molecules.

Any number of methods well known to those skilled in the art can be used to isolate and manipulate a DNA molecule, or fragment thereof, disclosed in the invention. For example, PCR (polymerase chain reaction) technology can be used to amplify a particular starting DNA molecule and/or to produce variants of the original molecule. DNA molecules, or fragment thereof, can also be obtained by other techniques such as by directly synthesizing the fragment by chemical means, as is commonly practiced by using an automated oligonucleotide synthesizer.

The DNA molecules and corresponding nucleotide sequences provided herein are therefore useful for, among other things, identifying soybean event MON 87708, selecting plant varieties or hybrids comprising soybean event MON 87708, detecting the presence of DNA derived from the transgenic soybean event MON 87708 in a sample, and monitoring samples for the presence and/or absence of soybean event MON 87708 or plant parts derived from soybean event MON 87708.

The invention provides soybean plants, progeny, seeds, plant cells, plant parts (such as pollen, ovule, pod, flower tissue, root tissue, stem tissue, and leaf tissue), and commodity products. These plants, progeny, seeds, plant cells, plant parts, and commodity products contain a detectable amount of a polynucleotide of the invention, i.e., such as a polynucleotide having at least one of the sequences provided as SEQ ID NO: 1-8. Plants, progeny, seeds, plant cells, and plant parts of the invention may also contain one or more additional transgenes. Such transgene may be any nucleotide sequence encoding a protein or RNA molecule conferring a desirable trait including but not limited to increased insect resistance, increased water use efficiency, increased yield performance, increased drought resistance, increased seed quality, improved nutritional quality, and/or increased herbicide tolerance, in which the desirable trait is measured with respect to a soybean plant lacking such additional transgene.

The invention provides soybean plants, progeny, seeds, plant cells, and plant part such as pollen, ovule, pod, flower, root or stem tissue, and leaves derived from a transgenic soybean plant event MON 87708. A representative sample of soybean event MON 87708 seed has been deposited according to the Budapest Treaty for the purpose of enabling the invention. The repository selected for receiving the deposit is the American Type Culture Collection (ATCC) having an address at 10801 University Boulevard, Manassas, Va. USA, Zip Code 20110. The ATCC repository has assigned the accession No. PTA-9670 to the event MON 87708 seed.

The invention provides a microorganism comprising a DNA molecule having SEQ ID NO: 1 and SEQ ID NO: 2 present in its genome. An example of such a microorganism is a transgenic plant cell. Microorganisms, such as a plant cell of The invention, are useful in many industrial applications, including but not limited to: (i) use as research tool for scientific inquiry or industrial research; (ii) use in culture for producing endogenous or recombinant carbohydrate, lipid, nucleic acid, or protein products or small molecules that may be used for subsequent scientific research or as industrial products; and (iii) use with modern plant tissue culture techniques to produce transgenic plants or plant tissue cultures that may then be used for agricultural research or production. The production and use of microorganisms such as transgenic plant cells utilizes modern microbiological techniques and human intervention to produce a man-made, unique microorganism. In this process, recombinant DNA is inserted into a plant cell's genome to create a transgenic plant cell that is separate and unique from naturally occurring plant cells. This transgenic plant cell can then be cultured much like bacteria and yeast cells using modern microbiology techniques and may exist in an undifferentiated, unicellular state. The new plant cell's genetic composition and phenotype is a technical effect created by the integration of the heterologous DNA into the genome of the cell. Another aspect of the invention is a method of using a microorganism of the invention. Methods of using microorganisms of the invention, such as transgenic plant cells, include (i) methods of producing transgenic cells by integrating recombinant DNA into the genome of the cell and then using this cell to derive additional cells possessing the same heterologous DNA; (ii) methods of culturing cells that contain recombinant DNA using modern microbiology techniques; (iii) methods of producing and purifying endogenous or recombinant carbohydrate, lipid, nucleic acid, or protein products from cultured cells; and (iv) methods of using modern plant tissue culture techniques with transgenic plant cells to produce transgenic plants or transgenic plant tissue cultures.

Plants of the invention may pass along the event DNA, including the transgene, to progeny. As used herein, "progeny" includes any plant, seed, plant cell, and/or regenerable plant part comprising the event DNA derived from an ancestor plant and/or a polynucleotide having at least one of the sequences provided as SEQ ID NO: 1 and SEQ ID NO: 2. Plants, progeny, and seeds may be homozygous or heterozygous for the transgene. Progeny may be grown from seeds produced by a soybean event MON 87708 plant and/or from seeds produced by a plant fertilized with pollen from a soybean event MON 87708 plant.

Progeny plants may be self-pollinated (also known as "selfing") to generate a true breeding line of plants, i.e., plants homozygous for the transgene. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes.

Alternatively, progeny plants may be outcrossed, e.g., bred with another unrelated plant, to produce a varietal or a hybrid seed or plant. The other unrelated plant may be transgenic or nontransgenic. A varietal or hybrid seed or plant of the invention may thus be derived by crossing a first parent that lacks the specific and unique DNA of the soybean event MON 87708 with a second parent comprising soybean event MON 87708, resulting in a hybrid comprising the specific and unique DNA of the soybean event MON 87708. Each parent can be a hybrid or an inbred/variatal, so long as the cross or breeding results in a plant or seed of the invention, i.e., a seed having at least one allele containing

the specific and unique DNA of soybean event MON 87708 and/or SEQ ID NO: 1 and SEQ ID NO: 2. Two different transgenic plants may thus be mated to produce hybrid offspring that contain two independently segregating, added, exogenous genes. For example, the MON 87708 dicamba tolerant soybean can be crossed with other transgenic soybean plant to produce a plant having the characteristics of both transgenic parents. One example of this would be a cross of MON 87708 dicamba tolerant soybean with a plant having one or more additional traits such as herbicide tolerance (e.g., soybean event 40-3-2 or soybean event MON89788 (U.S. Patent Application Publication No. 20060282915)), insect control (e.g. soybean event MON87701 (U.S. Patent Application Publication No. 20090130071)), and/or other desirable traits (e.g. enhanced oil composition such as soybean event MON87769 (PCT Patent Publication WO2009102873)), resulting in a progeny plant or seed that is tolerant to dicamba and has one or more additional traits. Herbicides for which transgenic plant tolerance has been demonstrated and the method of the invention can be applied, include but are not limited to: glyphosate, glufosinate, sulfonylureas, imidazolinones, bromoxynil, delapon, cyclohexanedione, protoporphyrionogen oxidase inhibitors, and isoxasflutole herbicides. Nucleotide molecules encoding proteins involved in herbicide tolerance are known in the art and include, but are not limited to, a nucleotide molecule encoding: glyphosate-tolerant 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) (see, for example, U.S. Pat. Nos. 5,627,061; 5,633,435; 6,040,497; 5,094,945; 5,804,425; 6,248,876; 7,183,110; RE39,247); glyphosate oxidoreductase (GOX) (see, for example, U.S. Pat. No. 5,776,760); glyphosate-n-acetyltransferase (GAT); an herbicide-tolerant acetolactate synthase (ALS, also known as acetohydroxyacid synthase (AHAS)) for tolerance to sulfonylureas, imidazolinones, triazolopyrimidines, pyrimidinyl oxybenzoates, sulfonylamino carbonyl triazolinones, and/or heteroaryl ethers; an herbicide-tolerant acetyl coenzyme A carboxylase (AC-Case) or R-2,4-dichlorophenoxypropionate dioxygenase (rdpA) for tolerance to an aryloxyphenoxypropionate (AOPP) (such as haloxyfop, quizalofop, dichlorofop, and diclofop); a detoxification protein such as a 2,4-D dioxygenase (tfdA), R-2,4-dichlorophenoxypropionate dioxygenase (rdpA), Aryloxy Alkanoate Dioxygenase (AAD), and/or S-2,4-dichlorprop dioxygenase (sdpA) for tolerance to synthetic auxin herbicides; a bromoxynil nitrilase (Bxn) for Bromoxynil tolerance (see, for example, U.S. Pat. No. 4,810,648); a phytoene desaturase (crtI) for tolerance to norflurazon; the bialaphos resistance (bar) or phosphinothricin acetyltransferase (PAT) protein (see, for example, U.S. Pat. Nos. 5,646,024 and 5,276,268) for tolerance to glufosinate and bialaphos; and a protein for triketone (mezotrione, tembotrione, topromezone, isoxazole) herbicide-tolerance such as tolerant 4-HydroxyPhenylPyruvate Dioxygenase (HPPD), a detoxifying cytochrome P450, or an HPPD pathway bypass such as *Artrobacter globiformis* HPP oxidase (HPP0) and *Pseudomonas acidovorans* 4-HPA 1-hydroxylase (HPAH) and NADH oxidoreductase (HPAC). Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcox J. ed., American Society of Agronomy, Madison Wis. (1987).

The invention provides a plant part that is derived from soybean event MON 87708. As used herein, a "plant part"

refers to any part of a plant which is comprised of material derived from a soybean event MON 87708 plant. Plant parts include but are not limited to pollen, ovule, pod, flower, root or stem tissue, fibers, and leaves. Plant parts may be viable, nonviable, regenerable, and/or nonregenerable.

The invention provides a commodity product that is derived from soybean event MON 87708. As used herein, a "commodity product" refers to any composition or product which is comprised of material derived from a soybean event MON 87708 plant, seed, plant cell, or plant part. Commodity products may be sold to consumers and may be viable or nonviable. Nonviable commodity products include but are not limited to nonviable seeds and grains; processed seeds, seed parts, and plant parts; dehydrated plant tissue, frozen plant tissue, and processed plant tissue; seeds and plant parts processed for animal feed for terrestrial and/or aquatic animals consumption, oil, meal, flour, flakes, bran, fiber, milk, cheese, paper, cream, wine, and any other food for human consumption; and biomasses and fuel products. Viable commodity products include but are not limited to seeds and plant cells. The soybean event MON 87708 can thus be used to manufacture any commodity product typically acquired from soybean. Any such commodity product that is derived from the soybean event MON 87708 may contain at least a detectable amount of the specific and unique DNA corresponding to soybean event MON 87708, and specifically may contain a detectable amount of a polynucleotide containing at least 15 contiguous nucleotides of SEQ ID NO: 1 or SEQ ID NO: 2. Any standard method of detection for nucleotide molecules may be used, including methods of detection disclosed herein. A commodity product is within the scope of the invention if there is any detectable amount of SEQ ID NO: 1 or SEQ ID NO: 2 in the commodity product.

The plants, progeny, seeds, plant cells, plant parts (such as pollen, ovule, pod, flower, root or stem tissue, and leaves), and commodity products of the invention are therefore useful for, among other things, growing plants for the purpose of producing seed and/or plant parts of soybean event MON 87708 for agricultural purposes, producing progeny of soybean event MON 87708 for plant breeding and research purposes, use with microbiological techniques for industrial and research applications, and sale to consumers.

The invention provides methods for controlling weeds and methods for producing plants using dicamba herbicide and soybean event MON 87708. A method for controlling weeds in a field is provided and consists of planting soybean event MON 87708 varietal or hybrid plants in a field and applying a herbicidally effective dose of dicamba to the field for the purpose of controlling weeds in the field without injuring the MON 87708 plants. Such application of dicamba herbicide may be pre-emergence, i.e., any time after MON 87708 seed is planted and before MON 87708 plants emerge, or post-emergence, i.e., any time after MON 87708 plants emerge. Another method for controlling weeds in a field is also provided and consists of applying an effective dose of dicamba herbicide to control weeds in a field and then planting soybean event MON 87708 in the field. Such application of dicamba herbicide would be pre-planting, i.e., before MON 87708 seed is planted, and could be done any time pre-planting including, but not limited to, about 14 days pre-planting to about 1 day pre-planting. The invention also provides a method for producing soybean seed essentially free of the seeds of toxic weed species by planting seeds of a dicamba tolerant soybean variety MON 87708 in a field, applying a post-

emergence effective dose of dicamba herbicide sufficient to kill the toxic weed species to the field, and harvesting seed from the field. A herbicidally effective dose of dicamba for use in the field should consist of a range from about 0.005 pounds per acre to about 8 pounds of dicamba per acre over a growing season. Multiple applications of dicamba may be used over a growing season, for example, two applications (such as a pre-planting application and a post-emergence application or a pre-emergence application and a post-emergence application) or three applications (such as a pre-planting application, a pre-emergence application, and a post-emergence application).

Methods for producing an herbicide tolerant soybean plant comprising the DNA sequences specific and unique to the transgenic event MON 87708 of the invention are provided. Transgenic plants used in these methods may be homozygous or heterozygous for the transgene. Progeny plants produced by these methods may be varietal or hybrid plants; may be grown from seeds produced by a soybean event MON 87708 plant and/or from seeds produced by a plant fertilized with pollen from a soybean event MON 87708 plant; and may be homozygous or heterozygous for the transgene. Progeny plants may be subsequently self-pollinated to generate a true breeding line of plants, i.e., plants homozygous for the transgene, or alternatively may be outcrossed, e.g., bred with another unrelated plant, to produce a varietal or a hybrid seed or plant.

A soybean plant that tolerates application of dicamba herbicide may be produced by sexually crossing an event MON 87708 plant comprising a nucleotide molecule comprising the sequence of SEQ ID NO: 1 and SEQ ID NO: 2 with another soybean plant and thereby producing seed, which is then grown into progeny plants. These progeny plants may then be treated with dicamba herbicide to select for progeny plants that are tolerant to dicamba herbicide. Alternatively, these progeny plants may be analyzed using diagnostic methods to select for progeny plants that contain the event MON 87708 DNA. The other plant used in the crossing may or may not be tolerant to dicamba herbicide and may or may not be transgenic. The progeny plant and/or seed produced may be varietal or hybrid seed. In practicing this method, the step of sexually crossing one plant with another plant, i.e., cross-pollinating, may be accomplished or facilitated by human intervention, for example: by human hands collecting the pollen of one plant and contacting this pollen with the style or stigma of a second plant; by human hands and/or actions removing, destroying, or covering the stamen or anthers of a plant (e.g., by detasseling or by application of a chemical gametocide) so that natural self-pollination is prevented and cross-pollination would have to take place in order for fertilization to occur; by human placement of pollinating insects in a position for "directed pollination" (e.g., by placing beehives in orchards or fields or by caging plants with pollinating insects); by human opening or removing of parts of the flower to allow for placement or contact of foreign pollen on the style or stigma (e.g., in soy which naturally has flowers that hinder or prevent cross-pollination, making them naturally obligate self-pollinators without human intervention); by selective placement of plants (e.g., intentionally planting plants in pollinating proximity); and/or by application of chemicals to precipitate flowering or to foster receptivity (of the stigma for pollen).

A soybean plant that tolerates application of dicamba herbicide may be produced by selfing an event MON 87708 plant comprising a nucleotide molecule comprising the sequence of SEQ ID NO: 1 and SEQ ID NO: 2 and thereby producing seed, which is then grown into progeny plants.

These progeny plants may then be treated with dicamba herbicide to select for progeny plants that are tolerant to dicamba herbicide. Alternatively, these progeny plants may be analyzed using diagnostic methods to select for progeny plants that contain the event MON 87708 DNA. In practicing this method, the step of sexually crossing one plant with itself, i.e., self-pollinating or selfing, may be accomplished or facilitated by human intervention, for example: by human hands collecting the pollen of the plant and contacting this pollen with the style or stigma of the same plant and then optionally preventing further fertilization of the plant; by human hands and/or actions removing, destroying, or covering the stamen or anthers of other nearby plants (e.g., by detasseling or by application of a chemical gametocide) so that natural cross-pollination is prevented and self-pollination would have to take place in order for fertilization to occur; by human placement of pollinating insects in a position for "directed pollination" (e.g., by caging a plant alone with pollinating insects); by human manipulation of the flower or its parts to allow for self-pollination; by selective placement of plants (e.g., intentionally planting plants beyond pollinating proximity); and/or by application of chemicals to precipitate flowering or to foster receptivity (of the stigma for pollen).

Progeny soybean plants and seeds encompassed by these methods and produced by using these methods will be distinct from other soybean plants, for example because the progeny soybean plants and seeds: are recombinant and as such created by human intervention; are dicamba herbicide tolerant; contain at least one allele that consists of the transgene DNA of the invention; and/or contain a detectable amount of a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 and SEQ ID NO: 2. A seed may be selected from an individual progeny plant, and so long as the seed comprises SEQ ID NO: 1 and SEQ ID NO: 2, it will be within the scope of the invention.

In practicing the invention, two different transgenic plants can be crossed to produce hybrid offspring that contain two independently segregating heterologous genes. Selfing of appropriate progeny can produce plants that are homozygous for both genes. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcox J. ed., American Society of Agronomy, Madison Wis. (1987).

The plants and seeds used in the methods disclosed herein may also contain one or more additional transgenes. Such transgene may be any nucleotide sequence encoding a protein or RNA molecule conferring a desirable trait including but not limited to increased insect resistance, increased water use efficiency, increased yield performance, increased drought resistance, increased seed quality, improved nutritional quality, and/or increased herbicide tolerance, in which the desirable trait is measured with respect to a soybean plant lacking such additional transgene.

The methods of the invention are therefore useful for, among other things, controlling weeds in a field while growing plants for the purpose of producing seed and/or plant parts of soybean event MON 87708 for agricultural or research purposes, selecting for progeny of soybean event MON 87708 for plant breeding or research purposes, and producing progeny plants and seeds of soybean event MON 87708.

The plants, progeny, seeds, plant cells, plant parts (such as pollen, ovule, pod, flower, root or stem tissue, and leaves), and commodity products of the invention may be evaluated for DNA composition, gene expression, and/or protein expression. Such evaluation may be done by using any standard method such as PCR, northern blotting, southern analysis, western blotting, immuno-precipitation, and ELISA or by using the methods of detection and/or the detection kits provided herein.

Methods of detecting the presence of DNA derived from a soybean cell, tissue, seed, or plant of soybean event MON 87708 in a sample are provided. One method consists of (i) extracting a DNA sample from at least one soybean cell, tissue, seed, or plant, (ii) contacting the DNA sample with a primer pair that is capable of producing an amplicon from event MON 87708 DNA under conditions appropriate for DNA amplification, (iii) performing a DNA amplification reaction, and then (iv) detecting the amplicon molecule and/or confirming that the nucleotide sequence of the amplicon comprises a nucleotide sequence specific for event MON 87708, such as one selected from the group consisting of SEQ ID NO: 1-8. The amplicon should be one that is specific for event MON 87708, such as an amplicon that comprises SEQ ID NO: 1 or SEQ ID NO: 2. The detection of a nucleotide sequence specific for event MON 87708 in the amplicon is determinative and/or diagnostic for the presence of the soybean event MON 87708 specific DNA in the sample. An example of a primer pair that is capable of producing an amplicon from event MON 87708 DNA under conditions appropriate for DNA amplification is provided as SEQ ID NO: 10-11. Other primer pairs may be readily designed by one of skill in the art and would comprise at least one fragment of SEQ ID NO: 6. Another method of detecting the presence of DNA derived from a soybean cell, tissue, seed, or plant of soybean event MON 87708 in a sample consists of (i) extracting a DNA sample from at least one soybean cell, tissue, seed, or plant, (ii) contacting the DNA sample with a DNA probe specific for event MON 87708 DNA, (iii) allowing the probe and the DNA sample to hybridize under stringent hybridization conditions, and then (iv) detecting hybridization between the probe and the target DNA sample. An example of the sequence a DNA probe that is specific for event MON 87708 DNA is provided as SEQ ID NO: 11. Other probes may be readily designed by one of skill in the art and would comprise at least one fragment of SEQ ID NO: 6. Detection of probe hybridization to the DNA sample is diagnostic for the presence of soybean event MON 87708 specific DNA in the sample. Absence of hybridization is alternatively diagnostic of the absence of soybean event MON 87708 specific DNA in the sample.

DNA detection kits are provided that are useful for the identification of soybean event MON 87708 DNA in a sample and can also be applied to methods for breeding soybean plants containing the appropriate event DNA. Such kits contain DNA primers and/or probes comprising fragments of SEQ ID NO: 1-8. One example of such a kit comprises at least one DNA molecule of sufficient length of contiguous nucleotides of SEQ ID NO: 6 to function as a DNA probe useful for detecting the presence and/or absence of DNA derived from transgenic soybean event MON 87708 in a sample. The DNA derived from transgenic soybean event MON 87708 would comprise SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and/or SEQ ID NO: 8. A DNA molecule sufficient for use as a DNA probe is provided that is useful for determining, detecting, or diagnosing the presence and/or absence of soybean event MON 87708 DNA in a sample is provided as SEQ ID NO: 11. Other probes may

be readily designed by one of skill in the art and should comprise at least 15 contiguous nucleotides of SEQ ID NO: 6 and be sufficiently unique to soybean event MON 87708 DNA in order to identify DNA derived from the event. Another type of kit comprises a primer pair useful for producing an amplicon useful for detecting the presence and/or absence of DNA derived from transgenic soybean event MON 87708 in a sample. Such a kit would employ a method comprising contacting a target DNA sample with a primer pair as described herein, then performing a nucleic acid amplification reaction sufficient to produce an amplicon comprising SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and/or SEQ ID NO: 8, and then detecting the presence and/or absence of the amplicon. Such a method may also include sequencing the amplicon or a fragment thereof, which would be determinative of, i.e. diagnostic for, the presence of the soybean event MON 87708 specific DNA in the target DNA sample. Other primer pairs may be readily designed by one of skill in the art and should comprise at least 15 contiguous nucleotides of SEQ ID NO: 6 and be sufficiently unique to soybean event MON 87708 DNA in order to identify DNA derived from the event.

Nucleic-acid amplification can be accomplished by any of the various nucleic-acid amplification methods known in the art, including thermal amplification methods. Many techniques are known in the art for detecting, quantifying, and/or sequencing the amplicon produced by these methods. One exemplary technique useful in practicing this invention is TAQMAN® (PE Applied Bio systems, Foster City, Calif.).

The kits and detection methods of the invention are useful for, among other things, identifying soybean event MON 87708, selecting plant varieties or hybrids comprising soybean event MON 87708, detecting the presence of DNA derived from the transgenic soybean event MON 87708 in a sample, and monitoring samples for the presence and/or absence of soybean event MON 87708 or plant parts derived from soybean event MON 87708.

The sequence of the heterologous DNA insert, junction sequences, or flanking sequences from soybean event MON 87708 (with representative seed samples deposited as ATCC PTA-9670) can be verified (and corrected if necessary) by amplifying such sequences from the event using primers derived from the sequences provided herein followed by standard DNA sequencing of the amplicon or of the cloned DNA.

As used herein, the term “comprising” means “including but not limited to”.

The following examples are included to demonstrate examples of certain preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent approaches the inventors have found function well in the practice of the invention, and thus can be considered to constitute examples of preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLES

Example 1

Transformation of Soybean A3525 and MON 87708 Event Selection

The soybean plant MON 87708 was produced by Agrobacterium-mediated transformation of soybean. Soybean

cells were transformed and regenerated into intact soybean plants and individual plants were selected from the population of plants that showed integrity of the plant expression cassette and resistance to dicamba. From this population, soybean plant event MON 87708 was selected and characterized.

The transgenic dicamba tolerant soybean plant MON 87708 was developed through Agrobacterium-mediated transformation of soybean meristem tissue utilizing transformation vector PV-GMHT4355. The method was described in U.S. Pat. No. 6,384,301 (herein incorporated by reference), which allows for the generation of transformed plants without utilization of callus. Briefly, meristem tissues were excised from the embryos of germinated A3525 soybean seed (Asgrow, St Louis, Mo.). After co-culturing with Agrobacterium carrying the vector, the meristems were placed on selection medium containing glyphosate (Monsanto, St Louis, Mo.), carbenicillin disodium salt, cefotaxime sodium salt, and ticarcillin disodium salt/potassium clavulanate mixture to inhibit the growth of untransformed plant cells and excess Agrobacterium. The meristems were then placed in media conducive to shoot and root development. Rooted plants with normal phenotypic characteristics were selected and transferred to soil for growth and further assessment.

The R0 plants generated through the above transformation were transferred to soil for growth and then selfed to produce R1 seed. During subsequent selfing of the R0 plants to produce the R1 generation, the unlinked insertions of T-DNA I (dmo expression cassette) and T-DNA II (cp4 epsps expression cassette) were segregated. A non-lethal dose of glyphosate was applied to R1 plants. The plants with minor injuries were selected for further analyses, whereas plants showing no injury, i.e., containing T-DNA II (cp4 epsps expression cassette) were eliminated from subsequent development. Subsequently, R0 plants containing only a single T-DNA I insert (i.e., dmo gene cassette) were identified. The T-DNA I expression cassette comprised the Peanut Chlorotic Streak Virus (PC1SV) promoter with a duplicated enhancer region (P-PC1SV.FLT-enh); operably linked to a DNA leader derived from RNA transcript of Tobacco Etch Virus (L-TEV); operably linked to a DNA molecule encoding an N-terminal chloroplast transit peptide from ribulose 1,5-bisphosphate carboxylase small subunit (SSU) from *Pisum sativum* (TS-RbcS-3C); operably linked to part of the mature protein from ribulose 1,5-bisphosphate carboxylase small subunit (SSU) from *Pisum sativum* (CR-RbcS-3C); operably linked to a DNA molecule encoding a dicamba mono-oxygenase (DMO) from *Stenotrophomonas maltophilia* (*Pseudomonas maltophilia* was the original name of the source of the DMO gene. This source organism was subsequently reclassified first as *Xanthomonas maltophilia* and then as *Stenotrophomonas maltophilia*); operably linked to a 3' UTR DNA molecule derived from the ribulose 1,5-bisphosphate carboxylase small subunit gene of *Pisum sativum* (T-Ps.RbcS2-E9). Plants were selected by a combination of analytical techniques, including TaqMan, PCR analysis, and herbicide spray. The MON 87708 event was selected from among approximately 2,400 individual transgenic events based on its superior phenotypic characteristics, a comprehensive molecular profile analysis, and its desirable haplotype association. Event MON 87708 was then crossed with event MON 89788 (glyphosate tolerant). The progeny of this cross were treated with dicamba (Clarity®, BASF, Research Triangle Park, N.C.), glyphosate (Roundup WeatherMAX®, Monsanto Co., St Louis, Mo.), or a combination of dicamba and glyphosate. The treatments

were done at pre-plant, post-plant at the vegetative 3 growth stage (V3), and post-plant at the reproduction 1 stage (R1). Treated plants were scored for percent growth inhibition at 14 days after treatment (DAT) for the pre-plant herbicide treatment, 3 DAT for post-emergence treatment at the VE stage, and 3 DAT post-emergence treatment at the R1 stage. The herbicide(s) were applied at various rates per acre as shown in Table 1. Percent inhibition measurements represent an average of the repetitions.

TABLE 1

Dicamba and/or Roundup WeatherMAX® tolerance testing with MON89788 × MON 87708			
Herbicide (a.e. Rate gm/ha (lb/a))	% inhibition at 14 DAT PRE	% inhibition at 3 DAT POST (V3)	% inhibition at 3 DAT POST (R1)
Untreated/No herbicide	0.0	0.0	0.0
Roundup WeatherMAX® (3364 (3.0))	0.0	0.0	0.0
Clarity® (2244 (2.0))	0.0	10.0	20.0
Clarity® 561 (0.5) and Roundup WeatherMAX® (841 (0.75))	0.0	5.0	10.0
Clarity® (1120 (1.0)) and Roundup WeatherMAX® (1682 (1.5))	0.0	7.5	12.5
Clarity® (2244 (2.0)) and Roundup WeatherMAX® (3364 (3.0))	0.0	22.5	25.0

The dicamba tolerance transgene was mapped in soybean event MON 87708 to linkage group 9 at approximately map position 143.5. The associated haplotype window 19743 and 19767 has no effect on yield, maturity, height or lodging. Haplotype association information is provided in Table 2 where GM_A92205 indicates event MON 87708.

TABLE 2

Haplotype association LG9, Pos 143.5								
Event	Haplotype Window	Haplotype ID	Yield	Maturity	Height	Lodging	Haplotype sequence	Linkage group
GM_A92205	19743	1573355	0.00	-0.03	0.06	0.04	CGCTG	9
GM_A92205	19743	1573357	0.00	0.07	-0.03	-0.04	CGCTA	9
GM_A92205	19743	1573371	0.00	-0.09	-0.41	-0.09	CCCTG	9
GM_A92205	19743	1573373	0.00	-0.20	-0.01	-0.03	TG*GG	9
GM_A92205	19743	1573374	0.00	-0.08	-0.07	0.05	TG*GA	9
GM_A92205	19743	1573375	0.00	-0.15	0.05	0.04	CCCTA	9
GM_A92205	19743	1573376	0.00	-0.45	-0.14	0.00	TC*GG	9
GM_A92205	19767	1573486	0.00	0.00	-0.03	0.00	TACGGTC	9
GM_A92205	19767	1573493	0.00	0.00	0.22	0.00	AACAATT	9
GM_A92205	19767	1573494	0.00	0.00	0.03	0.00	TACAATC	9
GM_A92205	19767	1573495	0.00	0.00	0.07	0.00	TGAAACC	9
GM_A92205	19767	1573497	0.00	0.00	0.41	0.00	TACGGTT	9
GM_A92205	19767	1573499	0.00	0.00	-0.01	0.00	TGAAACT	9
GM_A92205	19767	1573500	0.00	0.00	0.06	0.00	TGAGACC	9
GM_A92205	19767	1573502	0.00	0.00	-0.07	0.00	AACAATC	9
GM_A92205	19767	1573503	0.00	0.00	0.08	0.00	AACGATC	9
GM_A92205	19767	1573504	0.00	0.00	0.07	0.00	TACAGTC	9
GM_A92205	19767	1573506	0.00	0.00	-0.03	0.00	AACGATT	9
GM_A92205	19767	1573507	0.00	0.00	0.20	0.00	TGAAATT	9

Example 2

Characterization of MON 87708 DNA Sequences

The DNA inserted into the genome of soybean plant MON 87708 and the flanking sequence was characterized by detailed molecular analyses. These analyses included: the insert sequence, the insert number (number of integration sites within the soybean genome), the copy number (number

of copies of transgene DNA within one locus), the integrity of the inserted gene cassette, the flanking sequences, and the association of the insertion with haplotype regions of the soybean genome.

Molecular DNA probes were used that included the intact coding region and its respective regulatory elements, the promoters, introns, and polyadenylation sequences of the plant expression cassettes. The analysis showed that MON 87708 contains a single transgene DNA insertion with one copy of the expression cassette. Inverse PCR and DNA sequence analyses were performed to determine the 5' and 3' insert-to-plant genome junctions, confirm the organization of the elements within the insert (FIG. 1), and determine the complete DNA sequence of the insert in soybean plant MON 87708 (provided herein as SEQ ID NO: 5). A soybean plant that comprises in its genome the linked transgene genetic elements shown in FIG. 1 and is resistant to dicamba is an aspect of the invention.

Sequences flanking the transgene DNA insertion in MON 87708 were determined using inverse PCR as described in Ochman et al., 1990 (PCR Protocols: A guide to Methods and Applications, Academic Press, Inc.) and/or genome walker techniques. Plant genomic DNA was isolated from both A3525 and the transgenic soybean lines from tissue grown under standard greenhouse conditions. Approximately 1 gram of young leaf tissue was combined with liquid nitrogen and ground to a fine powder using a mortar and pestle. DNA was extracted using a Nucleon™ PhytoPure™ Genomic DNA extraction kit (RPN8511, Amersham, Piscataway, N.J.) according to the manufacturer's protocol. After the final precipitation step, DNA was resuspended in 0.5 ml of TE (10 mM Tris-HCl pH 8.0, 1 mM EDTA). This method can be modified by one skilled in the art to extract DNA from any tissue of soybean, including, but not limited

to seed. An aliquot of DNA was digested with restriction endonucleases selected based upon restriction analysis of the transgene DNA. After self-ligation of restriction fragments, PCR was performed using primers designed from the transgene DNA sequence that would amplify sequences extending away from the 5' and 3' ends of the transgene DNA. PCR products were separated by agarose gel electrophoresis and purified using a QIAGEN gel purification kit (Qiagen, Valencia, Calif.). The subsequent DNA products were

sequenced directly using standard DNA sequencing protocols. The 5' flanking sequence which extends into the right border sequence of the expression cassette transgene DNA is presented as SEQ ID NO: 3 ([C], see FIG. 1). The 3' flanking sequence which extends into the left border sequence of the expression cassette transgene DNA is presented as SEQ ID NO: 4 ([D], see FIG. 1). The portion of the expression cassette DNA that was fully integrated into the A3525 genomic DNA is presented as SEQ ID NO: 5 ([E], see FIG. 1).

Isolated DNA molecule sequences were compared to the transgene DNA sequence to identify the flanking sequence and the co-isolated transgene DNA fragment. Confirmation of the presence of the expression cassette was achieved by PCR with primers designed based upon the deduced flanking sequence data and the known transgene DNA sequence. The wild type sequence corresponding to the same region in which the transgene DNA was integrated in the transformed line was isolated using primers designed from the flanking sequences in MON 87708. The PCR reactions were performed using the Elongase® amplification system (Invitrogen, Carlsbad, Calif.). The flanking DNA sequences in MON 87708 and the A3525 wild type sequence were analyzed against multiple nucleotide and protein databases. This information was used to examine the relationship of the transgene to the plant genome and to look for the insertion site integrity. The flanking sequence and wild type sequences were used to design primers for TAQMAN® endpoint assays used to identify the events. Zygosity assays were developed using this information.

Example 3

Event Specific Endpoint TAQMAN® Assays

This example describes an event specific endpoint TAQMAN® thermal amplification method developed to identify event MON 87708 in a sample. Examples of conditions useful with the event MON 87708 Specific Endpoint TAQMAN® method are as follows: Step 1: 18 megohm water adjusted for final volume of 10 µl. Step 2: 5.0 µl of 2× Universal Master Mix (dNTPs, enzyme, buffer) to a 1× final concentration. Step 3: 0.5 µl Event Primer-1 (SQ13570) and Event Primer-2 (SQ13571) Mix (resuspended in 18 megohm water to a concentration of 20 µM for each primer) to 1.0 µM final concentration (for example in a microcentrifuge tube, the following should be added to achieve 500 µl at a final concentration of 20 µM: 100 µl of Primer SQ13570 (SEQ ID NO: 9) at a concentration of 100 µM; 100 µl of Primer SQ13571 (SEQ ID NO: 10) at a concentration of 100 µM; 300 µl of 18 megohm water). Step 4: 0.2 µl Event 6-FAM™ MGB Probe PB4655 (resuspended in 18 megohm water to a concentration of 10 µM (SEQ ID NO: 11) to 0.2 µM final concentration. Step 5: 0.5 µl Internal Control Primer-1 and Internal Control Primer-2 Mix (resuspended in 18 megohm water to a concentration of 20 µM for each primer) to 1.0 µM final concentration. Step 6: 0.2 µl Internal Control VIC™ Probe to 0.2 µM final concentration (resuspended in 18 megohm water to a concentration of 10 µM) Step 7: 3.0 µl Extracted DNA (template) for each sample with one each of the following comprising 1. Leaf Samples to be analyzed; 2. Negative control (non-transgenic DNA); 3. Negative water control (no template); 4. Positive control MON 87708 DNA. Step 8: Thermocycler Conditions as follows: One Cycle at 50° C. for 2 minutes; One Cycle at 95° C. for 10 minutes; Ten Cycles of 95° C. for 15 seconds then 64° C. for 1 minute

with -1° C./cycle; Thirty Cycles of 95° C. for 15 seconds then 54° C. 1 minute; final cycle of 10° C.

The DNA primers used in the endpoint assay are primers SQ13570 (SEQ ID NO: 9), SQ13571 (SEQ ID NO: 10), and 6-FAM™ labeled probe PB4655 (SEQ ID NO: 11). 6-FAM™ is a fluorescent dye product of Applied Biosystems (Foster City, Calif.) attached to the DNA probe. For TAQMAN® MGB™ probes, the 5' exonuclease activity of Taq DNA polymerase cleaves the probe from the 5'-end, between the fluorophore and quencher. When hybridized to the target DNA strand, quencher and fluorophore are separated enough to produce a fluorescent signal, thus releasing fluorescence. SQ13570 (SEQ ID NO: 9) and SQ13571 (SEQ ID NO: 10) when used with these reaction methods with PB4655 (SEQ ID NO: 11) produce a DNA amplicon that is diagnostic for event MON 87708 DNA. The controls for this analysis should include a positive control from soybean containing event MON 87708 DNA, a negative control from non-transgenic soybean, and a negative control that contains no template DNA. Additionally, a control for the PCR reaction includes Internal Control Primers and an Internal Control Probe, specific to a single copy gene in the Glycine genome. One of skill in the art will know how to design primers specific to a single copy gene in the Glycine genome. These assays are optimized for use with either an Applied Biosystems GeneAmp® PCR System 9700 (run at maximum speed) or MJ Research DNA Engine PTC-225 thermal cycler. Other methods and apparatus known to those skilled in the art that produce amplicons that identify the event MON 87708 DNA is within the skill of the art.

R0 plants demonstrating the presence of the expression cassette were allowed to develop into fully mature plants. Probes designed based on the sequences of the dicamba tolerance transgene cassette were used to probe Southern blots to determine linkage. The R0 plants were also evaluated for copy number of the expression cassette using a combination of Southern analysis and endpoint TAQMAN®.

A zygosity assay is useful for determining if a plant comprising an event is homozygous for the event DNA; that is comprising the exogenous DNA in the same location on each chromosome of a chromosomal pair; or heterozygous for an event DNA, that is comprising the exogenous DNA on only one chromosome of a chromosomal pair; or is null for the event DNA, that is wildtype. The endpoint TAQMAN® thermal amplification method was also used to develop zygosity assays for event MON 87708. This example describes an event specific endpoint TAQMAN® thermal amplification method developed to determine the zygosity of event MON 87708 in a sample. For this assay, a three primer assay was employed wherein primer SQ20632 (SEQ ID NO: 12) hybridizes and extends specifically from the 3' junction of the inserted exogenous DNA and genomic DNA, primer SQ20636 (SEQ ID NO: 13) hybridizes and extends specifically from the DNA flanking the 3' side of the inserted exogenous DNA, and primer SQ20637 (SEQ ID NO: 14) hybridizes and extends specifically from genomic DNA into which was integrated the inserted exogenous DNA. The three primers are diagnostic for the event. In this example, primer SQ20636 (SEQ ID NO: 13) and primer SQ20632 (SEQ ID NO: 12) and the 6-FAM™-labeled oligonucleotide probe PB10130 (SEQ ID NO: 15) are diagnostic when there is a copy of the inserted exogenous DNA. In this example, SQ20636 (SEQ ID NO: 13) and primer SQ20637 (SEQ ID NO: 14) and the VIC™-labeled oligonucleotide probe PB10131 (SEQ ID NO: 16) are diagnostic when there is no copy of the inserted exogenous DNA present in the genomic DNA, i.e. wildtype.

When the three primers and two probes are mixed together in a PCR reaction with DNA extracted from a plant homozygous for event MON 87708, there is a fluorescent signal only from the 6-FAMTM-labeled oligonucleotide probe PB10130 (SEQ ID NO: 15) which is indicative of and diagnostic a plant homozygous for event MON 87708. When the three primers and two probes are mixed together in a PCR reaction with DNA extracted from a plant heterozygous for event MON 87708, there is a fluorescent signal from both the 6-FAMTM-labeled oligonucleotide probe PB10130 (SEQ ID NO: 15) and the VICTM-labeled oligonucleotide probe PB10131 (SEQ ID NO: 16) which is indicative of and diagnostic a plant heterozygous for event MON 87708. When the three primers and two probes are mixed together in a PCR reaction with DNA extracted from a plant which is null for event MON 87708 (i.e. wildtype), there is a fluorescent signal from only the VICTM-labeled oligonucleotide probe PB10131 (SEQ ID NO: 16) which is indicative of and diagnostic a plant null for event MON 87708, i.e. wildtype. Examples of conditions useful with this method are as follows. Step 1: 18 megohm water adjusted for final volume of 10 μ l. Step 2: 5.0 μ l of 2 \times Universal Master Mix (Applied Biosystems cat #4304437; dNTPs, enzyme, buffer) to a 1 \times final concentration. Step 3: 0.5 μ l of Zygoty Primers SQ20632, SQ20636, SQ20637 (resuspended in 18 megohm water to a concentration of 20 μ M for each primer) to a final concentration of 1.0 μ M. Step 4: 0.2 μ l 6-FAMTM Probe PB10130 (SEQ ID NO: 15) (resuspended in 18 megohm water to a concentration of 10 μ M) to 0.2 μ M final concentration. Step 5: 0.2 μ l VICTM Probe PB10131 (SEQ ID NO: 16) (resuspended in 18 megohm water to a concentration of 10 μ M) to 0.2 μ M final concentration. Step 6: 3.0 μ l Extracted DNA (template) for each sample with one each of the following comprising 1. Leaf Samples to be analyzed (4-80 ng of genomic DNA diluted in water); 2. Negative control (nontransgenic soybean DNA; 4 ng diluted in water); 3. Negative water control (no template; solution in which DNA was resuspended); 4. Positive control MON 87708 genomic DNA from known heterozygous event (4 ng diluted in water); 5. 4. Positive control MON 87708 genomic DNA from known homozygous event (4 ng diluted in water). Step 7: Gently mix. Step 8: Thermocycler Conditions when using Applied Biosystems GeneAmp[®] PCR System 9700 (run at maximum speed) or MJ Research DNA Engine PTC-225 thermal cycler are as follows: One Cycle at 50 $^{\circ}$ C. for 2 minutes; one cycle at 95 $^{\circ}$ C. for 10 minutes; Ten Cycles of (95 $^{\circ}$ C. for 15 seconds then 64 $^{\circ}$ C. for 1 minute (-1 $^{\circ}$ C./cycle); Thirty Cycles of (95 $^{\circ}$ C. for 15 seconds then 54 $^{\circ}$ C. for 1 minute); Optional additional 10 to 20 cycles (95 $^{\circ}$ C. for 15 seconds then 64 $^{\circ}$ C. for 1 minute (-1 $^{\circ}$ C./cycle) may provide more distinct population separation during EndPoint TaqMan[®] analysis; One cycle at 10 $^{\circ}$ C. hold.

Example 4

Identification of Event MON 87708 in any MON 87708 Breeding Activity

The following example describes how one may identify the MON 87708 event within progeny of any breeding activity using soybean event MON 87708.

DNA event primer pairs are used to produce an amplicon diagnostic for soybean event MON 87708. An amplicon diagnostic for MON 87708 comprises at least one junction sequence, provided as SEQ ID NO: 1 or SEQ ID NO: 2 or SEQ ID NO: 7 or SEQ ID NO: 8. Event primer pairs that

will produce a diagnostic amplicon for MON 87708 include primer pairs based upon the flanking sequences and the inserted expression cassette. To acquire a diagnostic amplicon in which SEQ ID NO: 1 is found, one would design a forward primer molecule based upon SEQ ID NO: 3 from bases 1 through 1126 and a reverse primer molecule based upon the inserted expression cassette DNA sequence (SEQ ID NO: 5 from positions 1 through 3003) in which the primer molecules are of sufficient length of contiguous nucleotides to specifically hybridize to SEQ ID NO: 3 and SEQ ID NO: 5. To acquire a diagnostic amplicon in which SEQ ID NO: 2 is found, one would design a forward primer molecule based upon the inserted expression cassette DNA sequence (SEQ ID NO: 5 from positions 1 through 3003) and a reverse primer molecule based upon the 3' flanking sequence (SEQ ID NO: 4 from bases 131 through 1947), in which the primer molecules are of sufficient length of contiguous nucleotides to specifically hybridize to SEQ ID NO: 4 and SEQ ID NO: 5. For practical purposes, one should design primers which produce amplicons of a limited size range, for example, between 100 to 1000 bases. Smaller (shorter polynucleotide length) sized amplicons in general are more reliably produced in PCR reactions, allow for shorter cycle times, and can be easily separated and visualized on agarose gels or adapted for use in endpoint TAQ-MAN[®]-like assays. Smaller amplicons can be produced and detected by methods known in the art of DNA amplicon detection. In addition, amplicons produced using the primer pairs can be cloned into vectors, propagated, isolated, and sequenced or can be sequenced directly with methods well established in the art. Any primer pair derived from the combination of SEQ ID NO: 3 and SEQ ID NO: 5 or the combination of SEQ ID NO: 4 and SEQ ID NO: 5 that are useful in a DNA amplification method to produce an amplicon diagnostic for MON 87708 or progeny thereof is an aspect of the invention. Any single isolated DNA polynucleotide primer molecule comprising at least 11 contiguous nucleotides of SEQ ID NO: 3, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for MON 87708 or progeny thereof is an aspect of the invention. Any single isolated DNA polynucleotide primer molecule comprising at least 11 contiguous nucleotides of SEQ ID NO: 4, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for MON 87708 or progeny thereof is an aspect of the invention. Any single isolated DNA polynucleotide primer molecule comprising at least 11 contiguous nucleotides of SEQ ID NO: 5, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for MON 87708 or progeny thereof is an aspect of the invention.

An example of the amplification conditions for this analysis is illustrated in Example 3. However, any modification of these methods or the use of DNA primers homologous or complementary to SEQ ID NO: 3 or SEQ ID NO: 4 or DNA sequences of the genetic elements contained in the transgene insert (SEQ ID NO: 5) of MON 87708 that produce an amplicon diagnostic for MON 87708 is within the art. A diagnostic amplicon comprises a DNA molecule homologous or complementary to at least one transgene/genomic junction DNA (SEQ ID NO: 1 or SEQ ID NO: 2 or SEQ ID NO: 7 or SEQ ID NO: 8), or a substantial portion thereof.

An analysis for event MON 87708 plant tissue sample should include a positive tissue control from event MON 87708, a negative control from a soybean plant that is not event MON 87708 (for example, but not limited to A3525), and a negative control that contains no soybean genomic

DNA. A primer pair that will amplify an endogenous soybean DNA molecule will serve as an internal control for the DNA amplification conditions. Additional primer sequences can be selected from SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5 by those skilled in the art of DNA amplification methods, and conditions selected for the production of an amplicon by the methods shown in Example 3 may differ, but result in an amplicon diagnostic for event MON 87708 DNA. The use of these DNA primer sequences with modifications to the methods of Example 3 are within the scope of the invention. The amplicon produced by at least one DNA primer sequence derived from SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5 that is diagnostic for MON 87708 is an aspect of the invention.

DNA detection kits contain at least one DNA primer of sufficient length of contiguous nucleotides derived from SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5, that when used in a DNA amplification method produces a diagnostic amplicon for MON 87708 or its progeny is an aspect of the invention. A MON 87708 soybean plant, plant part, plant cell, seed, or commodity product that will produce an amplicon diagnostic for MON 87708 when tested in a DNA

amplification method is an aspect of the invention. The assay for the MON 87708 amplicon can be performed by using an Applied Biosystems GeneAmp® PCR System 9700 (run at maximum speed) or MJ Research DNA Engine PTC-225 thermal cycler or any other amplification system that can be used to produce an amplicon diagnostic of MON 87708 as shown in Example 3.

A deposit of a representative sample of soybean event MON 87708 seed disclosed above and recited in the claims has been made under the Budapest Treaty with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110. The ATCC accession number for this deposit is PTA-9670. The deposit will be maintained in the depository for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer, and will be replaced as necessary during that period.

Having illustrated and described the principles of the invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 16

<210> SEQ ID NO 1

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric DNA molecule of soybean genomic DNA and transgene DNA

<400> SEQUENCE: 1

ttgatctcca tgagccattt agtctcacct tcaaactctg atagtttaaa ctgaaggcgg 60

<210> SEQ ID NO 2

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric DNA molecule of soybean genomic DNA and transgene DNA

<400> SEQUENCE: 2

tcattgctga tccatgtaga tttcccggac tttagctcaa aatgcatgta tttattagcg 60

<210> SEQ ID NO 3

<211> LENGTH: 1300

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric DNA molecule of soybean genomic DNA and transgene DNA

<400> SEQUENCE: 3

gtacaaataa aactttatct gtacatttcg ttagttaaata tatattttgt ccatcaaatt 60

gttccttttaa attaaaatct aaaaactaac tttaccgtaa agaaatatgc attcatgtat 120

accaaataaa attttgacaa gagtttaagt tttatattat gattcttaag gaaatcatta 180

tcattataat ttataaaaat aaataaattg ataattatat gctatatcaa tttatgattt 240

gatgagtatg tatgttttaa atgcgagatt ctgccgccgt tcgatatagt tagcagtaga 300

gcctgttct caccctcaca cctgctcagt gtgaacttta aaagggactt tgttgacaaa 360

-continued

```

tgtaggac gtcgtcttct ttgcaataa aaaatthtca tctgtthaaa acgtthttat 420
agtaaaatta taaatagaaa atthagttgt aaaatthgaa atataaattt aattagaatg 480
cattcacatt gtaattcttt tacattatta thttattacct aattataaat tatcaacaat 540
aaaatctgac atagtatatg thtagattaa aatthgtaa tgtaagtttg aattaaaaaa 600
gtatattata aacttgagtt tggataata thttthtca tcagactttg gtataatag 660
agttgatcta aaagtaagtt gaaggatacc aaagggtaat ctaaacaatgc atgagaaatg 720
ttggggaata tctthtagtg aaacaaaaag cthtataaat tatcatgtca tactatata 780
gacaaatggt ggtthttgta ataacttht taaatatgaa caacaggatt thtthtttac 840
cgtcaaaata tgaataaaaag thaaactctt aaattattga tcataggttt gcaatthttt 900
thtatagaga ggtthgcaat thctgagttc tcaataactg atgattaaat gcgcatcggt 960
tgcacatgcg aaaaataaatt taagaggtaa cthttgaagt ctacacttca tgcctggggc 1020
tgccaggaaa gcttagatct ccatgagcat ccacgagctt atccacgagc atccacgagc 1080
ttatccgatt tgagcattga thctcatgag cthtttagtc thcacttcaa aactgatag 1140
thtaactga aggcgggaaa cgacaactct atccccatca agctagcttc tgcaggtcct 1200
gctcgagcgg ccgcatctct tgagccaatc aaagaggagt gatgtagacc taaagcaata 1260
atggagccat gacgtaaggg cthacgcca tacgaaataa 1300

```

<210> SEQ ID NO 4

<211> LENGTH: 1947

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric DNA molecule of soybean genomic DNA and transgene DNA

<400> SEQUENCE: 4

```

atgtactttc atthtataat aacgctgcgg acatctacat thttgaattg aaaaaaatt 60
ggtaattact cthtctthtt thccatattg accatcatac thattgctga thcatgtaga 120
thtcccgac thtagctcaa aatgcatgta thttattagc thctgtcttt thgttaattt 180
gttctcatca taatatttg acaaaaatat agctaggaaa gctthccatg catatthttg 240
aagcaatgaa gtatatagtg gatgcaatgt thctatatac thcactagtc agaaaattgc 300
ggacagttct gagattgatt ggcttcatgg cctacggtg catctattac cattgtcttt 360
gcatthgtca taaaaaagg tggaccatat thcatgttat taaaaacaa caaatcacg 420
cagtgcacat cthctgcaga atgtgtaggt taaccttatt acacttgatt aagthaagt 480
thcatgccatt agthtgagat tgaactthaa thctthaatc aagatcttag atatggaaaa 540
aattgtaatt cthtaaaga taataagatt thtgataga aattaattat caatthtaca 600
thtaatacat aataatthga agaaaaaag taagggtcat aatcacta accagagtaa 660
thtgacacgt gaaggggaca ctatgaaagc aaactthtt thgttctta aggttaggca 720
aggaaagaa agaatthgca cthtaattag actatthtca aaattattat gthtcttht 780
cttatcttgc thaaatthg cthattgtgt thttattatt attattgta thcatgatca 840
attatctatc aaagatcgat thccaacctg ccaggaaatc cgctgattg thtgcttcca 900
atgtgagaga thcaagatca gaatthctga aggtagtgct gactaccaag gtagcaaaat 960
aatgatattg ggaaggtga aaaaatgta gtactagtag thctactaca aaatthcaa 1020
aagggtthtg tgattgtgc ataagaatct thttgcatth gtctgtaagc thgaaatta 1080

```

-continued

```

cacgtggcac aagtcacttg cagccaaaga acctttctgt gaccaattat gttccctgag 1140
ctgaatagtg gttcttattc taatctcatc aatatctaat tacctagtga atatactact 1200
agactattgc agtggttatta atatcttaat gatagactat tgcagcagac agaaattaca 1260
ggtattatta tatactaata tacaattctg cattttccac acttttcccc tgcccatgct 1320
tccatgccac tgaagtctga aaccacattg gcagatcttg ctatctagaa attaaataac 1380
aatataagtt tgtatattta tatttcatat ttttttagta catttttatt ttgcacactc 1440
tataattcca tgattccttg attatcggag aatgatgtga tatgcaaacc acgagttaga 1500
accatcaaat caagcaaaga tatggatgga atgcctttaa tggaaagatt aattcaaagg 1560
ggcagaaaact ggtaattttt tcttcaactg aatgctatgc agtatgcagc agatctttca 1620
ttacagaat atctgcaaaa ccttgtgttg gagatcttac ctattgaata atgatatagg 1680
taaaataaag tatttaattt caccataact tttaagatga tgttaaaatg atctatgcaa 1740
ttcatgttgg atcgaatatt aaagatgtca catctaataa tactatgata aaaataaagt 1800
ataatttctg atcttataag tcaaaataaa tcatgtaaat ataaattaat tctcttctta 1860
taaattaatt ttatataatt aagatagatc caatgtgaac tctaagacca tgcatatata 1920
aaaatcatta tcaagtgaat atgcaac 1947

```

<210> SEQ ID NO 5

<211> LENGTH: 3003

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Transgene insert

<400> SEQUENCE: 5

```

tcaaactg atagtttaa ctgaaggcgg gaaacgaca tctgatcccc atcaagctag 60
cttctgcagg tctgctcga gggccgcag atcttgagcc aatcaaagag gagtgatgta 120
gacctaaagc aataatggag ccatgacgta agggcttacg cccatacga ataattaaag 180
gctgatgtga cctgtcggtc tctcagaacc tttacttttt atgtttggcg tgtattttta 240
aatttccacg gcaatgacga tgtgacccaa cgagatcttg agccaatcaa agaggagtga 300
tgtagaccta aagcaataat ggagccatga cgtaagggct tacgccata cgaaataatt 360
aaaggctgat gtgacctgct ggtctctcag aacctttact ttttatattt ggctgtatt 420
tttaaatttc cacggcaatg acgatgtgac ctgtgcatcc gctttgccta taaataagtt 480
ttagtttgta ttgatcgaca cggctgagaa gacacggcca taagcttgga tctcagagaa 540
ttctcaacac aacatataca aaacaaacga atctcaagca atcaagcatt ctacttctat 600
tgcagcaatt taaatcattt cttttaaagc aaaagcaatt ttctgaaaat tttcaccatt 660
tacgaacgat agccatggct tctatgatat cctcttcgcg tgtgacaaca gtcagccgtg 720
cctctagggg gcaatccgcc gcaatggctc cattcggcgg cctcaaatcc atgactggat 780
tcccagtgag gaaggtcaac actgacatta cttccattac aagcaatggt ggaagagtaa 840
agtgcattgca ggtgtggcct ccaattggaa agaagaagtt tgagactctt tctatttgc 900
caccattgac gagagattcc cgggccatgg ccaccttctg ccgcaatgcc tggatgtgg 960
cggcgtgccc cgaggaactg tccgaaaagc cgctcggccg gacgattctc gacacaccgc 1020
tcgctcteta ccgccagccc gacgggtgtg tcgctggcgt gctcagatc tgtccgcacc 1080
gcttcggccc gctgagcgac ggcacctctg tcaacggcca tctccaatgc ccctatcacg 1140

```

-continued

```

ggctggaatt cgatggcggc gggcagtgcg tccataaacc gcacggcaat ggcgcccgcc 1200
cggttccgct caacgtccgc tccttcccgg tggtagagcg cgacgcgctg atctggatct 1260
gtcccggcga tccggcgcgtg gccgatcctg gggcgatccc cgacttcggc tgccgcgctc 1320
atcccgccta tcggaccgtc ggcggctatg ggcatgtcga ctgcaactac aagctgctgg 1380
tcgacaacct gatggacctc ggccacgccc aatatgtcca tcgcgccaac gccagaccg 1440
acgccttcga ccggctggag cgcgaggtga tcgtcggcga cggtagata caggcgcgta 1500
tgaagattcc cggcggcacg ccgagcgtgc tgatggccaa gttcctgcgc ggcgcccaata 1560
cccccgtcga cgcttggaac gacatccgct ggaacaagg ggcgcgatg ctcaacttca 1620
tcgcggtggc gccggaagg accccgaagg agcagagcat cactcgcgc ggtaccata 1680
tcctgacccc cgagacggag gcgagctgcc attatttctt cggctcctcg cgcaatttcg 1740
gcatcgacga tccggagatg gacggcgtgc tcgcagctg gcaggctcag gcgctggta 1800
aggaggacaa ggtcgtcgtc gaggcgatcg agcgcgcgc cgcctatgtc gaggcgaatg 1860
gcatccgccc ggcgatgctg tcgtgcgacg aagccgcagt ccgtgtcagc cgcgagatcg 1920
agaagcttga gcagctcgaa gccgcctgaa ccggcttatg ctgcacgggc ggggcggggc 1980
ggtttcgatc ggctcgctg tcccggcgat attctagagc tttcgttctg atcatcggtt 2040
tcgacaacgt tcgtcaagt caatgcatca gtttcattgc gcacacacca gaatcctact 2100
gagtttgagt attatggcat tgggaaaact gttttctctg taccatttgt tgtgcttgta 2160
atctactgtg tttttatc ggttttcgt atcgaactgt gaaatggaaa tggatggaga 2220
agagttaatg aatgatatgg tccttttggt cattctcaa ttaatattat ttgtttttc 2280
tcttatttgt tgtgtgtga atttgaaatt ataagagata tgcaaacatt ttgttttgag 2340
taaaaatgtg tcaaatcgtg gcctctaatt accgaagtta atatgaggag taaaacactt 2400
gtagttgtac cattatgctt attcactagg caacaaatat attttcagac ctgaaaaagc 2460
tgcaaatggt actgaataca agtatgtcct cttgtgtttt agacatttat gaactttcct 2520
ttatgtaatt ttccagaatc cttgtcagat tctaactatt gctttataat tatagttata 2580
ctcatggatt tgtagttgag tatgaaaata ttttttaag cattttatga cttgccaaatt 2640
gattgacaac atgcatcaat cgcggccgct ctagaactag tggatcccc cctttaaggg 2700
ggctgcagga attcgatatc aagctttggc gcgccaaatc gtgaagtttc tcatctaagc 2760
ccccatttgg acgtgaatgt agacacgtcg aaataaagat ttccgaatta gaataatttg 2820
tttattgctt tcgcctataa atacgacgga tcgtaatttg tcgttttatc aaaatgtact 2880
ttcattttat aataacgtg cggacatcta catttttgaa ttgaaaaaaa attggtaatt 2940
actctttctt tttctcata ttgaccatca tactcattgc tgatccatgt agatttcccg 3000
gac 3003

```

<210> SEQ ID NO 6

<211> LENGTH: 5946

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric DNA molecule of soybean genomic DNA
and transgene DNA

<400> SEQUENCE: 6

```

gtacaaataa aactttatct gtacatttcg ttagttaaat tatattttgt ccatcaaatt 60
gttcctttaa attaaaatct aaaaactaac tttaccgtaa agaaatatgc attcatgtat 120

```

-continued

accaaataaa	atdddgacaa	gagtttaagt	tttatattat	gattcttaag	gaaatcatta	180
tcattataat	ttataaaaat	aaataaattg	ataattatat	gctatatcaa	tttatgattt	240
gatgagtatg	tatgttttaa	atgcgagatt	ctgccgccgt	tcgatatagt	tagcagtaga	300
gcctgttct	caccctcaca	cctgctcagt	gtgaacttta	aaagggactt	tgttgacaaa	360
tgttaggatc	gtcgtcttct	tttgcaataa	aaaatdddca	tctgtttaaa	acgtdddttat	420
agtaaaatta	taaatagaaa	athtagttgt	aaaatdddga	atataaattt	aattagaatg	480
cattcacatt	gtaattcttt	tacattatta	tttattacct	aattataaat	tatcaacaat	540
aaaatctgac	atagtatatg	tttagattaa	aatttgtaaa	tgtaagtttg	aattaaaaaa	600
gtatattata	aacttgagtt	tggtataata	ttttttatca	tcagactttg	gtataaatatg	660
agttgatcta	aaagtaagtt	gaaggatacc	aaagggtaat	ctaaacatgc	atgagaaatg	720
ttggggaata	tcttttagtgt	aaacaaaaag	ccttataaat	tatcatgtca	tactatatat	780
gacaaatggt	ggtttttgta	ataatcattt	taaatatgaa	caacaggatt	ttctdddttac	840
cgtaaaaata	tgaataaaaag	ttaaactctt	aaattattga	tcataggttt	gcaatdddttt	900
tttatagaga	ggtttgcaat	ttctgagttc	tcaataactg	atgattaaat	gcgcatcgtt	960
tgcatgcatg	aaaataaatt	taagaggtaa	catttgaagt	ctcaccttca	tgtccggggc	1020
tgccaggaaa	gcttagatct	ccatgagcat	ccacgagctt	atccacgagc	atccacgagc	1080
ttatccgatt	tgagcattga	tctccatgag	ccatttagtc	tcaccttcaa	acactgatag	1140
tttaaaactga	aggcgggaaa	cgacaatctg	atccccatca	agctagcttc	tgcaggctct	1200
gctcgagcgg	ccgcagatct	tgagccaatc	aaagaggagt	gatgtagacc	taaagcaata	1260
atggagccat	gacgtaaggg	cttacgcccc	tacgaaataa	ttaaaggctg	atgtgacctg	1320
tcggtctctc	agaaccttta	ctdddttatgt	ttggcgtgta	ttdddtaaatt	tccacggcaa	1380
tgacgatgtg	acccaacgag	atcttgagcc	aatcaaagag	gagtgatgta	gacctaaagc	1440
aataatggag	ccatgacgta	agggcttacg	cccatacgaa	ataattaaag	gctgatgtga	1500
cctgtcggtc	tctcagaacc	tttactdddtt	atatttgccg	tgtatdddttt	aatdddccacg	1560
gcaatgacga	tgtgacctgt	gcatccgctt	tgctataaaa	taagtttttag	tttgtattga	1620
tcgacacggg	cgagaagaca	cggccataag	cttggatcct	cgagaattct	caacacaaca	1680
tatacaaaac	aaacgaatct	caagcaatca	agcattctac	ttctattgca	gcaatdddtaa	1740
tcatttcttt	taaagcaaaa	gcaatdddct	gaaaatdddca	accatdddacg	aacgatagcc	1800
atggcttctc	tgatatcctc	ttccgctgtg	acaacagtea	gccgtgctc	tagggggcaa	1860
tccgccgcaa	tggtccatt	cggcggcctc	aatccatga	ctggattccc	agtgaggaag	1920
gtcaaacactg	acattacttc	cattacaagc	aatggtggaa	gagtaaagtg	catgcaggtg	1980
tggcctccaa	ttggaagaa	gaagtttgag	actcttctct	attdgccacc	attgacgaga	2040
gattcccggg	ccatggccac	cttcgctccg	aatgcctggg	atgtggcggc	gctgcccag	2100
gaactgtccg	aaaagccgct	cggccggacg	attctcgaca	caccgctccg	gctctaccgc	2160
cagcccagcg	gtgtggtcgc	ggcgctgctc	gacatctgtc	cgcaccgctt	cgcgccgctg	2220
agcgacggca	tcctcgtcaa	cggccatctc	caatgcccct	atcacgggct	ggaattcgat	2280
ggcggcgggc	agtgcgtcca	taaccgcac	ggcaatggcg	cccgcccggc	ttcgctcaac	2340
gtccgctcct	tcccgggtgt	ggagcgcgac	gcgctgatct	ggatctgtcc	cggcgatccg	2400
gcgctggccg	atcctggggc	gatccccgac	ttcggtgcc	gcgtcgatcc	cgcctatcgg	2460
accgtcggcg	gctatgggca	tgtcgactgc	aactacaagc	tgctggtcga	caacctgatg	2520

-continued

gacctcgcc	acgccaata	tgtccatcgc	gccaacgccc	agaccgacgc	cttcgaccgg	2580
ctggagcgcg	aggtgatcgt	cggcgacggt	gagatacagg	cgctgatgaa	gattcccggc	2640
ggcacgccga	gcgtgctgat	ggccaagttc	ctgcgcggcg	ccaatacccc	cgtcgacgct	2700
tggaacgaca	tccgctgga	caaggtgagc	gcgatgctca	acttcacgc	ggtggcgccg	2760
gaaggcacc	cgaaggagca	gagcatccac	tcgcgcggta	cccatatcct	gacccccgag	2820
acggaggcga	gctgccatta	tttcttcggc	tcctcgcgca	atttcggcat	cgacgatccg	2880
gagatggacg	gcgtgctgcg	cagctggcag	gctcagggcg	tggtaagga	ggacaaggtc	2940
gtcgtcgagg	cgatcgagcg	cgcgcgcgcc	tatgtcgagg	cgaatggcat	ccgcccggcg	3000
atgctgtcgt	gcgacgaagc	cgcagtcctg	gtcagccgcg	agatcgagaa	gcttgagcag	3060
ctcgaagccg	cctgaaccgg	cttatgctgc	acgggcgggg	cggggcggtt	tcgatcggct	3120
cgctgtccc	ggcgatattc	tagagctttc	gttcgatca	tcggtttcga	caacgttcgt	3180
caagttcaat	gcatcagttt	cattgcgcac	acaccagaat	cctactgagt	ttgagtatta	3240
tggcattggg	aaaactgttt	ttcttgacc	attgtttg	cttgaattt	actgtgtttt	3300
ttattcgggt	ttcgctatcg	aactgtgaaa	tggaaatgga	tggagaagag	ttaatgaatg	3360
atatggctct	ttgttcatt	ctcaaattaa	tattatttgt	ttttctctt	attgtttgtg	3420
tgttgaattt	gaaattataa	gagatatgca	aacattttgt	tttgagtaa	aatgtgtcaa	3480
atcgtggcct	ctaatgaccg	aagttaatat	gaggagtaa	acactttag	ttgtaccatt	3540
atgcttattc	actaggcaac	aatatattt	tcagacctag	aaaagctgca	aatgttactg	3600
aatacaagta	tgtcctcttg	tgttttagac	atttatgaac	tttcctttat	gtaattttcc	3660
agaatccttg	tcagattcta	atcattgctt	tataattata	gttatactca	tggatttgta	3720
gttgagtatg	aaaatattt	ttaatgcatt	ttatgacttg	ccaattgatt	gacaacatgc	3780
atcaatcgcg	gccgctctag	aactagtgga	ttccccctt	taagggggt	gcaggaattc	3840
gatatcaagc	tttggcgcg	caaatcgtga	agtttctcat	ctaagcccc	atttgacgct	3900
gaatgtagac	acgtcgaaat	aaagatttcc	gaattagaat	aatttgttta	ttgctttcgc	3960
ctataaatac	gacggatcgt	aatttgcgt	tttatcaaaa	tgtactttca	ttttataata	4020
acgctgcgga	catctacatt	tttgaattga	aaaaaaattg	gtaattactc	tttcttttcc	4080
tccatattga	ccatcact	cattgctgat	ccatgtagat	ttcccggact	ttagctcaaa	4140
atgcatgtat	ttattagcgt	tctgtctttt	cgtaatttg	ttctcatcat	aatattgtga	4200
caaaaatata	gctaggaaag	ctttccatgc	atattttgta	agcaatgaag	tatatagtgg	4260
atgcaatgtc	tctatatatt	cactagtcga	gaaaattgcg	gacagttctg	agattgattg	4320
gcttcatggc	cctacgggtc	atctattacc	attgtctttg	catttgtcat	acaaaaaggt	4380
ggaccatatt	catgttatgt	aaaaacaaac	aaaatcacgc	agtgacatc	ttctgcagaa	4440
tgtgtaggtt	aaccttatta	cacttgatta	agttaagtgt	catgccatta	gtttgagatt	4500
gaacttaaaa	tctttaatca	agatcttaga	tatggaaaa	attgtaattc	cattaaagat	4560
aataagattt	ttgatagaa	attaattatc	aattttacat	taataacata	ataatttgaa	4620
gaaaaaagt	aagggtcata	atcactacta	ccagagtaat	ttgacacgtg	aaggggacac	4680
tatgaaagca	aattactttt	ggttcctaaa	ggttaggcaa	gggaaagaaa	gaatttgac	4740
ttaattagca	ctattttcaa	aattattatg	tttcttttcc	ttatcttget	taaaatttgc	4800
ttattgtgtt	attattatta	ttattgttat	gcatgatcaa	ttattcatca	aagatcgatc	4860

-continued

```

tccaacctgc caggaaatcc gctgatttgt ttgcttccaa tgtgagagat ccaagatcag 4920
aattctggaa ggtagtgctg actaccaagg tagcaaaata atgatattgg ggaagggtgaa 4980
aaatatgtag tactagtact tctactacaa aatttcaaaa agggttttgt gatttgtgca 5040
taagaatctt tttgcatttg tctgtaagct tgaaaattac acgtggcaca agtcacttgc 5100
agccaaagaa cctttctgtg accaattatg ttccctgagc tgaatagtgg ttcttattct 5160
aatctcatca atatctaatt acctagttaa tatactacta gactattgca gtgttattaa 5220
tatcttaatg atagactatt gcagcagaca gaaattacag gtattattat atactaatat 5280
acaattctgc atttccaca cttttcccct gccatgctt ccatgccact gaagtctgaa 5340
accacattgg cagattttgc tatctagaaa ttaaataaca atataagttt gtatatttat 5400
atctcatatt ttttagtac atttttattt tgcacactct ataattccat gattccttga 5460
ttatcggaga atgatgtgat atgcaaacca cgagttagaa ccatcaaatc aagcaaagat 5520
atggatggaa tgcctttaat ggaaagatta attcaaaggg gcagaaactg gtaatttttt 5580
cttcaactga atgctatgca gtatgcagca gatctttcat ttacagaata tctgcaaac 5640
cttgtgttgg agatcttacc tattgaataa tgatataggt aaaataaagt atttaatttc 5700
accataactt ttaagatgat gttaaaatga tctatgcaat tcatggttga tcgaatatta 5760
aagatgtcac atctaataat actatgataa aaataaagta taatttctga tcttataagt 5820
caaaataaat catgtaaata taaattaatt ctcttcttat aaattaattt tatataatta 5880
agatagatcc aatgtgaact ctaagacat gcatatataa aatcattat caagtgaata 5940
tgcaac 5946

```

```

<210> SEQ ID NO 7
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric DNA molecule of soybean genomic DNA
and transgene DNA

```

```

<400> SEQUENCE: 7

```

```

gagcttatcc gatttgagca ttgatctcca tgagccattt agtctcacct tcaaactctg 60
atagtttaaa ctgaaggcgg gaaacgacaa tctgatcccc 100

```

```

<210> SEQ ID NO 8
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric DNA molecule of soybean genomic DNA
and transgene DNA

```

```

<400> SEQUENCE: 8

```

```

ctccatattg accatcatac tcattgctga tccatgtaga tttcccggac tttagctcaa 60
aatgcatgta tttattagcg ttctgtcttt tcgttaattt 100

```

```

<210> SEQ ID NO 9
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
Primer

```

```

<400> SEQUENCE: 9

```

-continued

ttttctccat attgaccatc atactca 27

<210> SEQ ID NO 10
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
 Primer

<400> SEQUENCE: 10

agacagaacg ctaataaata catgcatt 28

<210> SEQ ID NO 11
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
 Primer

<400> SEQUENCE: 11

ttcccggact ttagctca 18

<210> SEQ ID NO 12
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
 Primer

<400> SEQUENCE: 12

gatttccgg actttagctc aa 22

<210> SEQ ID NO 13
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
 Primer

<400> SEQUENCE: 13

tcacaatatt atgatgagaa caaattaacg 30

<210> SEQ ID NO 14
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
 Primer

<400> SEQUENCE: 14

gccctaatag gattttaac ttttc 25

<210> SEQ ID NO 15
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
 Primer

<400> SEQUENCE: 15

atgcatgtat ttattagcgt tc 22

-continued

<210> SEQ ID NO 16
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide PCR
 Primer

<400> SEQUENCE: 16

acatgtgtga ctacttct

18

We claim:

1. A recombinant DNA molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: [1-8] 1-4, 6-8, and full complements thereof.

2. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:1.

3. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:2.

4. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:3.

5. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:4.

6. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:5.]

7. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:6.

8. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:7.

9. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:8.

10. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:1 and SEQ ID NO:2.

11. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:3 and SEQ ID NO:4.

12. The recombinant DNA molecule of claim 1, comprising the nucleotide sequence of SEQ ID NO:7 and SEQ ID NO:8.

13. The recombinant DNA molecule of claim 1, wherein said recombinant DNA molecule is from transgenic soybean event MON 87708, a representative sample of seed comprising said event having been deposited as ATCC PTA-9670.

14. The recombinant DNA molecule of claim 1, wherein said recombinant DNA molecule is in a soybean plant, soybean plant cell, soybean seed, soybean plant part, or commodity product.

15. A DNA primer or probe comprising a nucleotide sequence of sufficient length of contiguous nucleotides of SEQ ID NO:6, or full complements thereof, wherein said DNA primer or probe is diagnostic for event MON 87708 and wherein said DNA primer or probe hybridizes under stringent hybridization conditions with a DNA molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: [1-8] 1-4 and 6-8 and does not hybridize under the stringent hybridization conditions with a DNA molecule not comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: [1-8] 1-4 and 6-8.

16. A pair of DNA molecules comprising of a first DNA molecule and a second DNA molecule different from the first

15 DNA molecule, wherein said first and second DNA molecules each comprise a nucleotide sequence of sufficient length of contiguous nucleotides of SEQ ID NO:6, or a full complement thereof, to function as DNA primers that when used together in an amplification reaction with DNA from event MON 87708 produce an amplicon diagnostic for soybean event MON 87708 DNA in a sample.

17. A method of detecting the presence of a DNA molecule from soybean event MON 87708 in a sample, said method comprising:

a) contacting said sample with the DNA probe of claim 15;

b) subjecting said sample and said DNA probe to stringent hybridization conditions;

20 and c) detecting hybridization of said DNA probe to a DNA molecule in said sample, wherein the hybridization of said DNA probe to said DNA molecule indicates the presence of a DNA molecule from soybean event MON 87708 in said sample.

25 18. A method of detecting the presence of a DNA molecule from soybean event MON 87708 in a sample, said method comprising: a) contacting said sample with the pair of DNA molecules of claim 16;

b) performing an amplification reaction sufficient to produce a DNA amplicon comprising a sequence selected from the group consisting of SEQ ID NO: [1-8] 1-4, 6-8, and full complements thereof; and

c) detecting the presence of said DNA amplicon in said reaction, wherein the presence of said DNA amplicon in said reaction indicates the presence of a DNA molecule from soybean event MON 87708 in said sample.

35 19. A method of determining the zygosity of a soybean plant or seed comprising event MON 87708 comprising:

a) contacting a sample comprising soybean DNA with a primer set and a probe set capable of producing a first amplicon and a second amplicon;

b) performing a nucleic acid amplification reaction with said sample, primer set, and a probe set;

c) detecting in said nucleic acid amplification reaction said first amplicon that is diagnostic for event MON 87708 and said second amplicon that diagnostic for native soybean genomic DNA not comprising event MON 87708; and

d) analyzing for the presence and/or absence of said first amplicon and said second amplicon in said nucleic acid amplification reaction, wherein the presence of both amplicons indicates said sample is heterozygous for event MON 87708 and the presence of only said first amplicon indicates said sample is homozygous for event MON 87708.

20. The method of claim 19, wherein the primer set comprises SEQ ID NO:12, SEQ ID NO:13 and SEQ ID NO:14.

21. The method of claim 19, wherein the probe set comprises SEQ ID NO:15 and SEQ ID NO:16. 5

22. A DNA detection kit comprising a pair of DNA primers or at least one DNA probe, wherein a) the pair of DNA primers comprises at least one DNA molecule comprising a nucleotide sequence of sufficient length of contiguous nucleotides of SEQ ID NO:6, or full complements 10 thereof, wherein the pair of DNA primers is capable of producing an amplicon that is diagnostic for event MON 87708; and

b) the at least one DNA probe is diagnostic for event MON 87708. 15

* * * * *