

US00RE48097E

(19) **United States**
 (12) **Reissued Patent**
Maiti et al.

(10) **Patent Number: US RE48,097 E**
 (45) **Date of Reissued Patent: Jul. 14, 2020**

(54) **CEPHEM COMPOUNDS, THEIR PRODUCTION AND USE**

(71) Applicant: **NAEJA-RGM PHARMACEUTICALS ULC**, Alberta (CA)

(72) Inventors: **Samarendra Nath Maiti**, Alberta (CA); **Dai Quoc Nguyen**, Alberta (CA); **Andhe V. N. Reddy**, Alberta (CA); **Judy Yip**, Alberta (CA); **Chan Minh Ha**, Alberta (CA); **Rong Ling**, Alberta (CA); **Rudong Shan**, Alberta (CA); **Madhava Reddy Madala**, Alberta (CA)

(73) Assignee: **NAEJA-RGM PHARMACEUTICALS ULC**, Alberta (CA)

(21) Appl. No.: **16/225,892**

(22) Filed: **Dec. 19, 2018**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **9,751,894**
 Issued: **Sep. 5, 2017**
 Appl. No.: **15/369,181**
 Filed: **Dec. 5, 2016**

U.S. Applications:

(60) Provisional application No. 62/265,625, filed on Dec. 10, 2015.

(51) **Int. Cl.**

C07D 501/56 (2006.01)
A61K 31/352 (2006.01)
A61K 31/423 (2006.01)
A61K 31/428 (2006.01)
A61K 31/546 (2006.01)
A61K 31/69 (2006.01)
C07D 501/04 (2006.01)

(52) **U.S. Cl.**

CPC **C07D 501/56** (2013.01); **A61K 31/352** (2013.01); **A61K 31/423** (2013.01); **A61K 31/428** (2013.01); **A61K 31/546** (2013.01); **A61K 31/69** (2013.01); **C07D 501/04** (2013.01)

(58) **Field of Classification Search**

CPC **C07D 501/56**; **C07D 501/04**; **A61K 31/69**; **A61K 31/423**; **A61K 31/428**; **A61K 31/352**; **A61K 31/546**; **A61K 2300/00**; **A61P 31/04**

See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

4,406,899 A * 9/1983 Aburaki C07D 501/46
 514/202
 4,748,171 A * 5/1988 Yamauchi C07D 471/08
 514/202
 4,864,022 A * 9/1989 Miyake A07D 277/587
 540/222

4,910,301 A * 3/1990 Kaplan C07D 501/46
 540/222
 5,010,188 A * 4/1991 Yamauchi C07D 471/08
 540/221
 5,071,979 A * 12/1991 Lattrell C07D 501/46
 540/225
 5,173,485 A * 12/1992 Sakane C07D 519/00
 514/202
 5,215,982 A * 6/1993 Sakane C07D 501/00
 514/202
 5,998,611 A 12/1999 Shinohara et al.
 7,384,928 B2 * 6/2008 Nishitani A61K 31/546
 514/202
 8,809,314 B1 8/2014 He et al.
 9,751,894 B2 9/2017 Maiti et al.
 2012/0264727 A1 10/2012 Cho et al.
 2013/0079319 A1 3/2013 Yamawaki et al.
 2014/0256697 A1 9/2014 Yamawaki et al.

FOREIGN PATENT DOCUMENTS

EP 332156 9/1989
 EP 0832893 4/1998
 EP 1489084 12/2004
 EP 2341053 7/2011
 EP 2557082 2/2013

(Continued)

OTHER PUBLICATIONS

Extended Search Report for European Patent Application No. 16871861.7, dated Apr. 23, 2019, 7 pages.

Garigipati "An Efficient Conversion of Nitriles to Amidines," Tetrahedron Letters, 1990, vol. 31, No. 14, pp. 1969-1972.

Official Action for Canada Patent Application No. 3,005,779, dated Jun. 6, 2018, 5 pages.

Official Action for Canada Patent Application No. 3,005,779, dated Sep. 7, 2018, 4 pages.

Official Action with English Translation for Korea Patent Application No. 10-2018-7015784, dated Oct. 25, 2018, 6 pages.

Judkins et al. "A Versatile Synthesis of Amidines From Nitriles Via Amidoximes," Synthetic Communications, 1996, vol. 26, No. 23, pp. 4351-4367.

(Continued)

Primary Examiner — Alan D Diamond

(74) *Attorney, Agent, or Firm* — Sheridan Ross P.C.

(57)

ABSTRACT

Cephem compounds, pharmaceutically acceptable salts thereof, and methods of using same, wherein the compound has a bicyclic nitrogen-containing aromatic heterocyclic ring as the quaternary ammoniomethyl group at the 3-position and one or both of a terminal amidine residue (substituted or unsubstituted) attached to an aryl or a 5- or 6-membered heteroaryl group (substituted or unsubstituted) which is further attached through a spacer to the free N-atom of the quaternary nitrogen-containing bicyclic ring at the 3-side chain, or a terminal guanidine residue attached to an aryl or a 5- or 6-membered heteroaryl group (substituted or unsubstituted) which is further attached through a spacer to the free N-atom of the quaternary nitrogen-containing bicyclic ring at the 3-side chain.

14 Claims, No Drawings

(56)

References Cited

FOREIGN PATENT DOCUMENTS

EP	2703406	3/2014
EP	2706062	3/2014
WO	WO 2004/039814	5/2004
WO	WO 2007/119511	10/2007
WO	WO 2011/136268	11/2011
WO	WO 2013/052568	4/2013
WO	WO 2014/068388	5/2014
WO	WO 2014/087165	6/2014

OTHER PUBLICATIONS

Nadrah et al. "Preparation of Amidines by Amidoxime Reduction with Potassium Formate," *Synlett*, May 2007, Iss. 8, pp. 1257-1258.

Ohki et al. "Studies on 3'-Quaternary Ammonium Cephalosporins II)," *The Journal of Antibiotics*, Sep. 1995, vol. 48, No. 9, pp. 1049-1051.

Pfaff et al. "A Lewis acid-promoted Pinner reaction," *Beilstein Journal of Organic Chemistry*, 2013, vol. 9, pp. 1572-1577.

Pinner et al. "Unwandlung der Nitrile in Imide." *Berichte der deutschen chemischen Gesellschaft*, Jul.-Dec. 1877, vol. 10, No. 2, pp. 1889-1897.

Weintraub et al. "A Convenient General Synthesis of Amidines," *The Journal of Organic Chemistry*, Apr. 1968, vol. 33, No. 4, pp. 1679-1681.

Yamawaki et al. "A novel series of parenteral cephalosporins exhibiting potent activities against *Pseudomonas aeruginosa* and other Gram-negative pathogens: Synthesis and structure-activity relationships," *Bioorganic & Medicinal Chemistry*, Nov. 2007, vol. 15, No. 21, pp. 6716-6732.

* cited by examiner

**CEPHEM COMPOUNDS, THEIR
PRODUCTION AND USE**

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue; a claim printed with strikethrough indicates that the claim was canceled, disclaimed, or held invalid by a prior post-patent action or proceeding.

CROSS-REFERENCE TO RELATED
APPLICATIONS

This Application is a Reissue of U.S. application Ser. No. 15/369,181, filed Dec. 5, 2016, now U.S. Pat. No. 9,751,894, which claims priority to U.S. Provisional Patent Application No. 62/265,625, filed Dec. 10, 2015, entitled "Cephem Derivatives As Broad-Spectrum Antibacterial Agents", both of which [is] are hereby incorporated by reference in [its] their entirety.

TECHNICAL FIELD

New cephem compounds, their pharmaceutically acceptable salts, their use, and the methods for preparation of these new compounds are provided. More particularly, cephem compounds having improved antibacterial activity, and combinations of the present cephem antibiotics with β -lactamase inhibitors that are active against a number of resistant pathogenic microorganisms are provided.

BACKGROUND

Cephem antibiotics have been widely used for the treatment of bacterial infections both in hospitals and in the general public. As such, it is highly desirable to use cephem antibiotics showing activity against both gram-positive and gram-negative bacteria. Unfortunately, due to the existence of multiple-drug resistant gram-negative and gram-positive organisms, many bacteria have become highly resistant to a number of β -lactam drugs, including the bacteria that constitute the ESKAPE organisms, generally encompassed by the following six pathogens: *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacter* species. This includes a resistance to known cepheims whereby the bacteria are producing β -lactamases, including Class A (ESBL) and Class D (serine β -lactamases) types, which have an extended substrate spectrum. As a result, the morbidity and mortality caused by bacterial infections in both hospital and community settings continues to rise, and has become a significant public health concern.

There is a demand for development of new cephem compounds which exhibit more potent antimicrobial activity in particular more effective against a variety of β -lactamase producing gram-positive and gram-negative bacteria.

A variety of cephem compounds having a quaternary ammonium group at the 3-side chain and 2-(2-aminothiazol-4-yl)-2-hydroxy-(or substituted hydroxyl)aminoacetamido group at 7-position have been published such as, for example, U.S. Pat. Nos. 4,864,022, 5,173,485, 5,215,982, 5,071,979, 4,406,899, 4,910,301, 4,748,171, 5,010,188, International Patent Application Nos. PCT/JP2003/013684,

PCT/JP2007/056136, PCT/IB2013/002423, European Patent Application No. 2 703 406, European Patent Application No. 2 557 082; European Patent No. 2 341 053; European Patent Application No. 2 70606, United States Patent Application No. 2013/0079319 (Shionogi), United States Patent Application No. 2012/0264727, and European Patent Application No. 2706062.

The above mentioned patents and patent applications disclose cephem compounds whose quaternary ammonium-ethyl group at the 3-position has a monocyclic or bicyclic nitrogen-containing heterocyclic or aromatic heterocyclic ring which is optionally substituted with an organic group other than benzamidine, heteroaryl amidine, arylguanidine and heteroaryl guanidine as the terminal residues.

There remains a need for new cephem compounds having increased antibiotic efficacy, particularly in highly resistant gram-positive and gram-negative bacteria, the compounds having structural features are significantly different from the compounds described in the patent references cited above

SUMMARY

According to embodiments herein, cephem compounds and pharmaceutically acceptable salts thereof are provided having a bicyclic nitrogen containing aromatic heterocyclic ring as the quaternary ammoniomethyl group at the 3-position and the free N-atom of the said bicyclic aromatic heterocyclic ring further attached through one or two carbon spacers to a residue like benzamidine, heteroarylamidine, arylguanidine and heteroaryl guanidine or similar basic functionality bearing terminal residues. The present compounds, and pharmaceutically acceptable salts thereof, comprise different structural characteristics to known cephem compounds, providing improved antibacterial activity, particularly when used in combination with one or more β -lactamase inhibitors. It may be an advantage of the present compounds, and pharmaceutically acceptable salts thereof, to provide antibacterial activity against "third-generation" and "fourth-generation" cephalosporin-resistant bacteria including gram-negative and gram-positive strains.

In some embodiments, the present cephem compounds, and pharmaceutically acceptable salts thereof, may comprise the following structural features:


(1) A bicyclic nitrogen containing aromatic heterocyclic ring as the quaternary ammoniomethyl group at the 3-position, and

(2) A terminal amidine residue (substituted or unsubstituted) attached to an aryl or a 5- or 6-membered heteroaryl group (substituted or unsubstituted) which is further attached through a spacer to the free N-atom of the quaternary nitrogen containing bicyclic aromatic heterocyclic ring at the 3-side chain, or

(3) A terminal guanidine residue attached to an aryl or a 5- or 6-membered heteroaryl group (substituted or unsubstituted) which is further attached through a spacer to the free N-atom of the quaternary nitrogen containing bicyclic aromatic heterocyclic ring at the 3-side chain.

In some embodiments, the present cephem compound may be represented by the general formula (I) or a pharmaceutically acceptable salt thereof:

3



Accordingly, in some embodiments, cephem compounds and pharmaceutically acceptable salts thereof, i.e., compounds of formula (I) and pharmaceutically acceptable salts thereof are provided. In some aspects, these compounds may exhibit activity against pathogenic microorganisms, therefore useful in the treatment of bacterial infections in humans or animals either alone or in combination with other β -lactam and/or non β -lactam β -lactamase inhibitors.

In other embodiments, pharmaceutical compositions comprising one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, and a pharmaceutically acceptable carrier or diluent are provided. In some aspects, these compositions may exhibit activity against pathogenic organisms.

In other embodiments, processes for the preparation of the new cephem compounds and salts thereof, i.e., the compounds of formula (I) and pharmaceutically acceptable salts thereof are provided.

In other embodiments, pharmaceutical compositions comprising (i) one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, (ii) one or more β -lactamase inhibitors, and (iii) a pharmaceutically acceptable carrier or diluent are provided. In some aspects, these compounds may exhibit activity against pathogenic microorganisms.

In other embodiments, methods for treating bacterial infections in a subject, comprising providing or administering to a subject in need thereof:

- (i) a therapeutically effective amount of one or more compounds of formula (I), or pharmaceutically acceptable salts thereof;
- (ii) a therapeutically effective amount of a pharmaceutical composition comprising one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, and a pharmaceutically acceptable carrier or diluent;
- (iii) a therapeutically effective amount of a combination comprising (a) one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, and (b) one or more β -lactamase inhibitors; or
- (iv) a therapeutically effective amount of a pharmaceutical composition comprising (a) one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, (b) one or more β -lactamase inhibitors, and (c) a pharmaceutically acceptable carrier or diluent.

In other embodiments, the use of a compound comprising formula (I) for bacterial infections in a subject is provided. In some aspects, the compound may further comprise a β -lactamase inhibitor, and may involve the preparation of a therapeutically effective medicament.

According to embodiments herein, the present subject may be a human being or an animal, or any other organism


4

in which the present compounds and compositions may provide a beneficial antibacterial effect.


DETAILED DESCRIPTION OF THE EMBODIMENTS

According to embodiments herein, cephem compounds of general formula (I) and pharmaceutically acceptable salts of the compounds of formula (I) are provided, wherein the compounds comprise antibiotics suitable for use either alone or in combination with β -lactamase inhibitors for the treatment of bacterial infections. The term "antibiotic" refers to a compound or composition which decreases the viability of a microorganism, or which inhibits the growth or proliferation of a microorganism and is further intended to include an antimicrobial, bacteriostatic or bactericidal agent.

More specifically, the present cephem compounds, and pharmaceutically acceptable salts thereof, may be represented by the following general formula (I):




In the formula (I), A is defined by the formula (Ia):



20

where X is N, C(H), C(F) or C(Cl);

B is defined as hydrogen, methyl, ethyl or represented by the formula (Ib)




wherein, R¹ and R² is independently hydrogen or lower alkyl, or R¹ and R² together may form a 3 to 6-membered spiro ring system; and m is 0 or 1.

C is defined as a quaternized bicyclic nitrogen containing aromatic heterocyclic ring.

Further, regarding object (1a), it is understood that said objects include syn isomer (Z form), anti isomer (E form) and a mixture thereof.


In some preferred embodiments, the quaternized bicyclic nitrogen containing aromatic heterocyclic rings representing C, may comprise (Ic-Iz):

5










6

-continued



7

-continued




where D is represented by CH_2 , CH_2CH_2 or CH_2CO ; and

E signifies an optionally substituted benzene ring or an optionally substituted 5- or 6-membered aromatic heterocyclic ring having at least one heteroatom selected from O, S and N. Such heteroaromatic rings include pyrrolyl, imidazolyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazolyl, isoxazolyl, oxazolyl, oxadiazolyl, isothiazolyl, thiazolyl, thiadiazolyl, furyl, thienyl and the like.

In the above formula (I), E may even more preferably selected from aryl or 5- and 6-membered aromatic heterocyclic rings as indicated below:

8



In the above definition, optional substituents include chloro, fluoro, cyano, hydroxy, carboxy, acetyl, methoxy, ethoxy, trifluoromethyl, pyrrolidinyloxy, piperidinyloxy and the like.


Preferably, the substituent is selected from chloro, fluoro, hydroxy, methoxy, trifluoromethyl, pyrrolidinyloxy.

Having further regard to the formula (I), in the definition of E the aryl or aromatic heterocyclic ring can accommodate up to 3 substituents selected from: F, Cl, CH_3 , C_2H_5 , CF_3 , NH_2 , NHCH_3 , CONH_2 , CONHCH_3 , OCH_3 .

In the formula (I), F is optionally substituted amidine or optionally substituted guanidine, and









G is hydrogen, methyl, ethyl, C_{3-6} alkyl, C_{3-6} cycloalkyl or an optionally substituted 5- or 6-membered aliphatic or an optionally substituted 5- or 6-membered aromatic heterocyclic ring, in which the heterocyclic ring is substituted with at least 1-2 hetero atoms selected from N, O, and S (α or β).

In the formula (I), some preferred examples of “-C-D-E-F” include the fragments (1 to 277) as shown below:










9

-continued



10

-continued



13

14

15

16









17

18

19








11

-continued




12

-continued




13

-continued










14

-continued










15

-continued









16

-continued









17

-continued









18

-continued








19

-continued




20

-continued




21

-continued











22

-continued



23

-continued



101

5

102

10

15

103

20

104

25

30

105

35

40

106

45

50

107

55








108

60

65

24

-continued



109

5

110

10

15

111

20

112

113

35

40

114

45

50

115

55


108

60

65


25

-continued









26

-continued










27

-continued



28








-continued



65









29

-continued



30

-continued



145

5

10

146

15

147

20

25

148

30

149

40

150

50

151

55


151

60

65


31

-continued










32

-continued












33

-continued









34

-continued









35

-continued









36

-continued



37

-continued



38

-continued

199

5

10

200

15

20

201

25

30

202

35

40

203

45

50

55

204

60

65

205

206








207

208

209








210

211










39

-continued



40








-continued



65








41

-continued



42







-continued



65








43

-continued




44

-continued




45

-continued









46

-continued









47

-continued



48

-continued



60

Regarding the compounds of general formula (I), it is to be understood that said compounds include syn-isomer, anti-isomer and a mixture thereof All tautomeric isomers are contemplated, however, for the purposes of example, the


49

object compound (I). As used herein the term 'isomers' refers to different compounds that have the same molecular formula but differ in arrangement and configuration of the atoms, such as geometrical isomers and optical isomers. For a given compound herein, it is to be understood that a substituent may be attached at a chiral center of a carbon atom. Therefore, enantiomers, diastereoisomers or racemates of the compound are contemplated.

It is further understood that isotopically labeled compounds, which are identical to those recited in Formula (I), so but for the fact one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number predominantly found in nature are also contemplated. Examples of isotopes that can be incorporated into the present compounds may include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as, but not limited to, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S and ^{18}F respectively. The present compounds, prodrugs thereof, and pharmaceutically acceptable salts of said compounds which contain the above mentioned isotopes and and/or other isotopes of other atoms are contemplated. Isotopically labeled compounds and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the examples below, by substituting a readily available isotopically labeled agent.


The term " β -lactamase inhibitor" refers to a compound which is capable of inhibiting β -lactamase activity, where inhibiting β -lactamase activity means inhibiting the activity of a class A, C or D β -lactamase. The term " β -lactamase" denotes an enzyme capable of inactivating a β -lactam antibiotic like a cephem antibiotic. The β -lactamase inhibitor may be, but is not limited to, the following groups:

Group 1: An oxapenam derivative of the general formula (II):




in which R^3 signifies OR^4 , $\text{S}(\text{O})_n\text{R}^4$ or a 5-6 membered heteroaromatic ring which may be substituted with 1 to 5 substituents selected from alkyl, hydroxyl, alkoxy, amino, alkylamino, dialkylamino and halogen; whereby $n=0, 1,$ or 2 and R^4 is hydrogen, alkyl, $(\text{C}_2\text{-C}_7)$ alkene, $(\text{C}_2\text{-C}_7)$ alkyne or a 5-6 membered heteroaromatic ring which may be substituted with 1 to 5 substituents selected from alkyl, hydroxyl, alkoxy, amino, alkylamino, dialkylamino and halogen, or a pharmaceutically acceptable salt thereof.

Example of a specific compound from Group 1 is clavulanic acid (IIa):





50


Group 2: A penam sulfone derivative of the general formula (III):



In which R^5 signifies hydrogen or halogen; R^6 signifies CH_2R^7 ; $\text{CH}=\text{CHR}^7$ wherein R^7 is hydrogen, halogen, cyano, carboxylic acid, acyl such as acetyl, carboxamide which may be substituted, alkoxy, carbonyl or a 5-6 membered heteroaromatic ring which is optionally substituted with 1 to 5 substituents selected from alkyl, hydroxyl, alkoxy, amino, alkylamino, dialkylamino and halogen; or which is optionally fused with a 5-6 membered heteroaromatic ring; $\text{CH}=\text{NR}^8$ where R^8 is amino, alkylamino, dialkylamino, aminocarbonyl, acylamino such as acetyl, hydroxyl, alkoxy, or a pharmaceutical acceptable salt thereof. Examples of two specific compounds from Group 2 are sulbactam (IIIa) and tazobactam (IIIb):




Group 3: A penem derivative of the general formula (IV):




In which R^9 signifies a 5-6 membered heteroaromatic ring which may be substituted with 1 to 5 substituents selected from alkyl, hydroxyl, alkoxy, amino, alkylamino, dialkylamino and halogen; or which is optionally fused with a 5-6 membered heteroaromatic ring; or a pharmaceutical acceptable salt thereof. Example of a specific compound from Group 3 is BRL-42,715 (IVa):

51




(IVa)

Group 4: A cyclic boronic acid derivative of the general formula (V):




(V)

Wherein, R^{10} signifies a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted carbocyclyl and substituted or unsubstituted heterocyclyl. The substituent is selected from alkyl, hydroxyl, alkoxy, amino, alkylamino, dialkylamino and halogen; or a pharmaceutical acceptable salt thereof. Example of a specific compound from Group 4 is RPX-7009 (Va):



(Va)


Group 5: A cyclic boronic acid of the general formula (VI):



(VI)

Wherein R^{11} signifies alkoxy, substituted or unsubstituted thioheteroaryl or substituted carboxamide; or a pharmaceutically acceptable salt thereof. Examples of specific compounds from Group 5 are RPX 7262 (VIa), RPX 7282 (VIb), RPX 7381 (VIc), and RPX 7400 (VIId):

52




5

RPX 7262 (VIa)

10


15



RPX 7282 (VIb)

20


25



RPX 7381 (VIc)

30

35




RPX 7400 (VIId)

Group 6: A diazabicyclooctane derivative of the general formula (VII):

(Va)

45



(VII)


Wherein, R^{12} signifies carboxylic acid, alkoxy, heteroaryl, nitrile, or carboxamide which may be substituted; or a pharmaceutical acceptable salt thereof. Examples of several specific compounds from Group 6 are (VIIa-VIIg):

(VI)

55

60

65




(VIIa, NXL-104)


53

-continued


(VIIb, MK-7655)




(VIIc)




(VIIId)




(VIIe)



(VIIf)



(VIIg)



It has been found that the efficacy of cephem compounds of the formula (I) herein against gram-negative bacteria can be potentiated by co-using a β -lactamase inhibitor selected from any one of the formula (II) to (VII).

A "pharmaceutically acceptable salt" of formula (I) refers to a salt which possesses the desired pharmacological activity of the parent compound. Examples of the groups for forming a pharmaceutically acceptable salt include: inorganic base salts, ammonium salts, organic base salts, basic amino acid salts, inorganic acid addition salts, and organic acid addition salts. Inorganic bases that can form the inorganic base salts include alkali metals (e.g., sodium, potassium, and lithium) and alkaline earth metals (e.g., calcium and magnesium). Organic bases that can form the organic

54

base salts include n-propylamine, n-butylamine, cyclohexylamine, benzylamine, octylamine, ethanolamine, diethanolamine, diethylamine, triethylamine, dicyclohexylamine, procaine, choline, picoline, N,N-dibenzylethylenediamine, N-methylglucamine, morpholine, pyrrolidine, pyridine, piperidine, N-ethylpiperidine and N-methylmorpholine. Basic amino acids that can form the basic amino acid salts include lysine, arginine, ornithine and histidine. As will be appreciated by one skilled in the art, the compounds of formula (I) containing a basic nitrogen atom are capable of forming acid addition salts. Such salts with pharmaceutically acceptable acids are included herein. Examples of such acids are hydrochloric, hydrobromic, phosphoric, sulfuric, citric, oxalic, maleic, fumaric, glycolic, mandelic, tartaric, aspartic, succinic, malic, formic, acetic, trifluoroacetic, methanesulfonic, ethanesulfonic, trifluoromethanesulfonic, benzenesulfonic, p-toluenesulfonic and the like.

Moreover, some compounds of formula (I) when they contain a basic group such as NH, NH_2 or pyridine and the like may form an inner zwitterionic salt with COOH group. Such inner salts are also contemplated herein.

Pharmaceutically acceptable solvates of the compounds of formula (I) are contemplated herein. 'Pharmaceutically acceptable solvate' refers to a molecular complex of a compound with one or more solvent molecules in a stoichiometric or non-stoichiometric amount. Such solvent molecules are those commonly used in the pharmaceutical art, which are known to be innocuous to recipient, e.g., water, ethanol, and the like. A molecular complex of a compound or moiety of a compound and a solvent can be stabilized by non-covalent intra-molecular forces such as, for example, electrostatic forces, Van der Waals forces or hydrogen bonds. The term hydrate refers to a complex where the one or more solvent molecules are water.

A combination of one or more of the compounds of formula (I) and one or more β -lactamase inhibitors are provided, where the β -lactamase inhibitors may be selected from formula (II) to (VII). Such combinations may exhibit a synergistic effect when used in the treatment of bacterial infections.

A combination of pharmaceutical compositions comprising one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, and one or more pharmaceutically acceptable carriers or diluents are provided. Further, pharmaceutical compositions comprising (i) one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, (ii) one or more β -lactamase inhibitors, and (iii) one or more pharmaceutically acceptable carriers or diluents are also provided. Preferably, the β -lactamase inhibitors may be selected from formula (II) to (VII). It is understood that any of the compositions and combinations herein may be administered to a subject such as, for example, by parenteral, in particular intramuscular route, oral, sublingual, rectal, aerosol or by local route in a topical application on the skin and the mucous membranes. Suitable pharmaceutically acceptable carriers and diluents include excipients such as starch, glucose, lactose, sucrose, gelatin, gum Arabic, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene glycol, water, ethanol, and the like. Compositions of the present disclosure, if desired, can also contain minor amounts of wetting, dispersing or emulsifying agents, or pH buffering agents, and preservatives. In addition, auxiliary, stabilizing, thickening, lubricating, and coloring agents can be included. Pharmaceutical compositions can be formulated in a conventional manner. Proper formulation is dependent upon the route of admin-

istration chosen. The present pharmaceutical compositions can take the form of injectable preparations, suspensions, emulsions, sugar-coated tablets, pellets, gelatin-capsules, capsules containing liquids, powders, granules, sustained-release formulations, suppositories, aerosols, sprays, ointments, creams or any other form suitable for use.

In a pharmaceutical composition containing the present compounds, the weight ratio of active ingredient to carrier will normally be in the range of 1:20 to 20:1.

As above, methods for treating bacterial infections in a subject are also provided, said methods comprising administering to a subject in need thereof:

- (i) a therapeutically effective amount of one or more compounds formula (I), or pharmaceutically acceptable salts thereof;
- (ii) a therapeutically effective amount of a pharmaceutical composition comprising one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, and a pharmaceutically acceptable carrier or diluent;
- (iii) a therapeutically effective amount of a combination comprising (a) one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, and (b) one or more β -lactamase inhibitors; or
- (iv) a therapeutically effective amount of a pharmaceutical composition comprising (a) one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, (b) one or more β -lactamase inhibitors, and (c) a pharmaceutically acceptable carrier or diluent.

In some embodiments, the present methods for preventing bacterial infections in a subject comprise providing a subject in need thereof:

- (i) a therapeutically effective amount of one or more compounds formula (I), or pharmaceutically acceptable salts thereof;
- (ii) a therapeutically effective amount of a pharmaceutical composition comprising one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, and a pharmaceutically acceptable carrier or diluent;
- (iii) a therapeutically effective amount of a combination comprising (a) one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, and (b) one or more β -lactamase inhibitors; or
- (iv) a therapeutically effective amount of a pharmaceutical composition comprising (a) one or more compounds of formula (I), or pharmaceutically acceptable salts thereof, (b) one or more β -lactamase inhibitors, and (c) a pharmaceutically acceptable carrier or diluent.

The present methods for treating or preventing a bacterial infection in a subject may comprise administering to a subject in need thereof (i) a therapeutically effective amount of one or more compounds formula (I), or pharmaceutically acceptable salts thereof, and (ii) a therapeutically effective amount of one or more β -lactamase inhibitors. It will thus be apparent that in the treating or preventing of bacterial infections, compounds of formula (I) and β -lactamase inhibitors may be administered to the subject in the same pharmaceutical formulation (e.g., a pharmaceutical composition comprising compounds of formula (I), β -lactamase inhibitors, and a carrier or diluent) or different pharmaceutical formulations (e.g., a first pharmaceutical composition—comprising compounds of formula (I) and a carrier or diluent; and a second pharmaceutical composition—comprising β -lactamase inhibitors and a carrier or diluent). When administered in different formulations, the first and

second pharmaceutical compositions may be administered simultaneously, sequentially, or separated in time.

In other embodiments, the use, in the manufacture of a medicament, of a compound of formula (I) as an active ingredient is provided, wherein the active ingredient may be provided in an antibacterial composition in admixture with a carrier. In some embodiments, the use, in the manufacture of a medicament, may further comprise the compound of formula (I) in combination with one or more β -lactamase inhibitors as active ingredients, in an antibacterial composition in admixture with a carrier. In other embodiments, the use, in the manufacture of a medicament, may further comprise the compound of formula (I) in combination with one or more antibiotics β -lactamase inhibitors as active ingredients, or in combination with one or more antibiotics (e.g., a β -lactam antibiotic or some other antibiotic) as active ingredients, in an antibacterial composition in admixture with a carrier.

The parenteral administration which includes intramuscular, intraperitoneal, subcutaneous and intravenous use, sterile solutions of the active ingredients are usually prepared and the pH of the solutions are suitably adjusted and buffered. For intravenous use, the total concentration of solutes should be controlled to render the preparation isotonic. Suitable solvents include saline solution (e.g., 0.9% NaCl solution) and a pyrogenic sterile water. Pharmaceutical compositions for oral delivery can be, for example, in the form of tablets, lozenges, aqueous or oily suspensions, granules, powders, emulsions, capsules, syrups, or elixirs. Orally administered compositions can contain one or more optional agents, for example, sweetening agents such as fructose, aspartame, or saccharin, flavoring agents such as peppermint, oil of wintergreen, cherry, coloring agents, and preserving agents to provide a pharmaceutically palatable preparation. Moreover, when in tablet form, the compositions can be coated to delay disintegration and absorption in the gastrointestinal tract, thereby providing a sustained action over an extended period of time. Oral compositions can include standard vehicles such as mannitol, lactose, starch, magnesium stearate, sodium saccharin, cellulose, magnesium carbonate, and the like. For oral liquid preparations, for example, suspensions, elixirs, and solutions, suitable carriers, excipients, or diluents include water, saline, alkylene glycols (e.g. propylene glycol), polyalkylene glycols (e.g., polyethylene glycol), oils, alcohols, slightly acidic buffers ranging from about pH 4 to about pH 6 (e.g., acetate, citrate, ascorbate ranging from about 5 mM to about 50 mM), and the like. Additionally, flavoring agents, preservatives, coloring agents, bile salts, acylcarnitines, and the like can be added.

For topical formulations of the present compounds, creams, gels, ointments or viscous lotions can be used as appropriate delivery forms. Topical delivery systems also include transdermal patches containing at least one compound of formula (I) to be administered. Delivery through the skin can be achieved by diffusion or by more active energy sources such as iontophoresis or electrotransport. Formulations of a compound herein, for topical use, such as in creams, ointments, and gels, can include an oleaginous or water soluble ointment base, for example, topical compositions can include vegetable oils, animal fats, and in certain embodiments, semisolid hydrocarbons obtained from petroleum. Topical compositions can further include white ointment, yellow ointment, cetyl esters wax, oleic acid, olive oil, paraffin, petrolatum, white petrolatum, spermaceti, starch glycerite, white wax, yellow wax, lanolin, and glyceryl monostearate. Various water-soluble ointment bases can also

be used, including glycol ethers and derivatives, polyethylene glycols, polyoxyl 40 stearate, and polysorbates.

In a pharmaceutical composition containing the present cephem compounds, the weight ratio of active ingredient to carrier will normally be in the range of 1:20 to 20:1.

The therapeutically effective amount of the compounds of formula (I) and pharmaceutically acceptable salts thereof and the amounts sufficient to achieve the stated goals of the methods disclosed herein vary depending upon the physical characteristics of the subject, the severity of the subject's symptoms, the formulation and the means used to administer the drug, and the method being practiced. The specific dose for a given subject is usually set by the judgment of the attending physician. However, a therapeutically effective and/or sufficient amount of the compounds and salts of the present compositions is typically between about 1 mg/kg body weight to 500 mg/kg body weight, including from 1 to 100 mg/kg, from 1 to 75 mg/kg, from 1 to 50 mg/kg, from 1 to 25 mg/kg, from 25 to 150 mg/kg, from 25 to 125 mg/kg, from 25 to 100 mg/kg, from 25 to 75 mg/kg, from 25 to 50 mg/kg, from 50 to 150 mg/kg, from 50 to 125 mg/kg, and from 50 to 100 mg/kg, regardless of the formulation. In equally preferred aspects, a therapeutically effective amount is about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 or 35 mg/kg body weight, regardless of the formulation. In some situations, a dose less than 1 mg/kg body weight or greater than 500 mg/kg body weight may be effective.

In a particular oral formulation for use in the present methods, the compound of formula (I) or a pharmaceutically acceptable salt thereof may be in the form of a capsule containing the compound or salt. Suitable amounts of the compound or salt may range from about 10 to about 3000 mg, with preferred amounts including about 100, 125, 150, 175, 200, 225, 250, 275, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450 and 1500 mg.

In a particular intravenous (IV) formulation for use in the present methods, the compound of formula (I) or a pharmaceutically acceptable salt thereof is administered in a dosage of between about 100 mg and 2000 mg, preferably about 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 450, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1500 or more mg, by IV infusion over approximately 60, 90, 120 or more minutes, every 6, 12, 18 or 24 hours for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more days. The compound of formula (I) or a pharmaceutically acceptable salt thereof may be reconstituted in sterile water for injection (WFI) or be diluted in 5% dextrose in water, for example.

The terms "dose", "unit dose", "unit dosage", or "effective dose" refer to physically discrete units that contain a predetermined quantity of active ingredient calculated to produce a desired therapeutic effect. These terms are synonymous with the therapeutically effective amounts and amounts sufficient to achieve the stated goals of the methods disclosed herein.

'Therapeutically effective amount' refers to the amount of a compound that, when administered to a subject for treating a disease, or at least one of the clinical symptoms of a disease, is sufficient to affect such treatment of the disease, disorder, or symptom. The therapeutically effective amount can vary depending, for example, on the compound, the disease, disorder, and/or symptoms of the disease, severity of the disease, disorder, and/or symptoms of the disease, the age, weight, and/or health of the patient to be treated, and the judgement of the prescribing physician.

Administration frequencies of doses for the treatment of a bacterial infection include 4, 3, 2 or once daily, every other day, every third day, every fourth day, every fifth day, every sixth day, once weekly, every eight days, every nine days, every ten days, bi-weekly, monthly and bi-monthly. Depending on the means of administration, the dosage may be administered all at once, such as with an oral formulation in a capsule, or slowly over a period of time, such as with an intravenous administration. For slower means of administration, the administering period can be a matter of minutes, such as about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120 or more minutes, or a period of hours, such as about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5 or more hours.

The weight ratio of (i) the present compounds and (ii) a β -lactamase inhibitor or an antibiotic (if it is being administered with a β -lactamase inhibitor or an antibiotic, e.g., a β -lactam antibiotic or some other antibiotic) will normally be in the range from about 1:20 to about 20:1.

In some embodiments, it is an aim to provide an improved method for the treatment of bacterial infections caused by β -lactamase producing bacteria in a patient in need of such treatment comprising administering to the patient a therapeutically effective amount of at least one compound chosen from formula (I) or a pharmaceutically acceptable salt thereof, either alone or in combination with one or more β -lactamase inhibitors, including the β -lactamase inhibitor of formula (II) to (VII) provided above. In such an embodiment, the compounds may increase the antibacterial effectiveness of β -lactamase susceptible β -lactam antibiotics, that is, they may increase the effectiveness of the antibiotic against infections caused by β -lactamase producing microorganisms in mammalian subjects, particularly in humans. As such, the present compounds of formula (I) and pharmaceutically acceptable salts thereof may be valuable for co-administration with β -lactamase inhibitors. In this regard, the present compounds of formula (I) or a pharmaceutically acceptable salt thereof can be mixed with the β -lactamase inhibitors, and the two agents thereby administered simultaneously. Alternatively, the two agents can be administered sequentially, either one immediately after the other or separated in time by 1, 5, 10, 15, 30, 45 or 60 minutes, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18 or more hours, or 1, 2, 3, 4, 5 or more days. When co-administered with a β -lactamase inhibitor, the present compounds and the β -lactamase inhibitor may, in combination, provide a synergistic effect. The term 'synergistic effect' refers to the effect produced when two or more agents are co-administered is greater than the effect produced when the agents are administered individually. Alternatively, the compound of formula (I) or a salt thereof can be administered as a separate agent during a course of treatment with the β -lactamase inhibitor.

In other embodiments, it is an aim to provide an improved method for the treatment of bacterial infections caused by β -lactamase producing bacteria in a patient in need of such treatment comprising administering to the patient a therapeutically effective amount of at least one compound chosen from formula (I) or a pharmaceutically acceptable salt thereof, either alone or in combination with one or more antibiotics, including β -lactam antibiotics and non β -lactam antibiotics. In such an embodiment, the compounds may increase the antibacterial effectiveness of β -lactamase susceptible β -lactam antibiotics, that is, they may increase the effectiveness of the antibiotic against infections caused by β -lactamase producing microorganisms in mammalian subjects, particularly in human. In that regard, the compounds

59

of formula (I) and pharmaceutically acceptable salts thereof may be valuable for co-administration with β -lactam antibiotics. In the treatment of a bacterial infection, the present so compounds of formula (I) or a pharmaceutically salt thereof can be mixed with the β -lactam antibiotic, and the two agents thereby administered simultaneously. Alternatively, the two agents can be administered sequentially, either one immediately after the other or separated in time by 1, 5, 10, 15, 30, 45 or 60 minutes, or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18 or more hours, or 1, 2, 3, 4, 5 or more days. When co-administered with a β -lactam

60

antibiotic, the combination of the present compound and the antibiotic can provide a synergistic effect. The term 'synergistic effect' refers to the effect produced when two or more agents are co-administered is greater than the effect produced when the agents are administered individually. Alternatively, the compound of formula (I) or a salt thereof can be administered as a separate agent during a course of treatment with the antibiotic. Examples of compounds of the formula (I), without limiting to the specified compounds, are provided in the following Table 1.

TABLE 1

Compound No.	Structure	Chemical Name
1		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-3-[[1-(4-carbamimidoyl)benzyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
2		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino}acetyl]-amino]-8-oxo-3-[[1-(4-[N-(piperidin-3-yl)carbamimidoyl]benzyl)-1H-pyrazolo[4,3-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
3		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino}-acetyl]amino]-8-oxo-3-[[1-(4-[N-(piperidin-3-yl)carbamimidoyl]benzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
4		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]-imino}acetyl]amino]-3-[[1-(4-carbamimidamidobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
5		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
6		(6R,7S)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-(2-carbamimidamidoethoxy)benzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
7		(6R,7R)-3-[[1-(4-[N-(2-aminoethyl)carbamimidoyl]benzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
8		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-(4,5-dihydro-1H-imidazol-2-yl)benzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
9		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-2,3-dihydro-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
10		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-(2-carbamimidamidoethoxy)benzyl)-2,3-dihydro-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
11		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
12		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-[2-(4-carbamimidamidophenyl)ethyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
13		(6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(4-carbamimidoylfuran-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl}methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
14		(6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
15		(6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
16		(6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(4-carbamimidoyl-1,3-thiazol-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl}methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
17		(6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(3-[(diaminomethylidene)amino]-1,2-oxazol-5-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
18		(6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(4-carbamimidoylthiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
19		(6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(4-(N-methoxycarbamimidoyl)benzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
20		(6R,7R)-7-{[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
21		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
22		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[(carboxymethoxy)imino]acetyl]amino)-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
23		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino)-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
24		(6R,7R)-7-[[[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino)-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
25		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoylfuran-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
26		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoyl-1,3-thiazol-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
27		(6R,7R)-7-[[[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
28		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
34		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[(4-carbamimidamidothiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
35		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(hydroxyimino)acetyl]amino]-3-({1-[(4-carbamimidamido-1,3-thiazol-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
36		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[(4-carbamimidamido-1,3-thiazol-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
37		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(hydroxyimino)acetyl]amino]-3-({1-[(4-carbamimidamido-benzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
38		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[(4-carbamimidamido-benzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
39		(6R,7R)-1-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(hydroxyimino)acetyl]amino]-3-({1-[(4-carbamimidamidofuran-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
40		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[(4-carbamimidamidofuran-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
41		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-{{(2-carboxypropan-2-yl)oxy}imino}acetyl]amino]-3-({1-[(4-carbamimidamidothiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
42		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-{{(2-carboxypropan-2-yl)oxy}imino}acetyl]amino]-3-({1-[(4-carbamimidamidofuran-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
43		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[2-(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidamido-benzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
44		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[2-(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidamidofuran-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
45		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[2-(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[4-(hydrazinylcarboximidoyl)-benzyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl)-8-oxo-5-thia-1-azabicyclo-[4.2.0]oct-2-ene-2-carboxylate
46		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[2-(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(3-carbamimidamidopropyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo-[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
47		(6R,7R)-7-{[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-{[1-(3-carbamimidamido-propyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
48		(6R,7R)-7-{[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(hydroxyimino)acetyl]amino}-3-{[1-(3-carbamimidamido-propyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
49		(6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-{(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-{[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
50		(6R,7R)-7-{[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-{[1-(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
51		(6R,7R)-7-{[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-{[1-(4-carbamimidoylthiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
52		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-{[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl}]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
53		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-{[1-(4-carbamimidoyl-1,3-thiazol-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl}]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
54		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-{[1-(4-{N-[(3R)-piperidin-3-yl]carbamidoyl}benzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl}]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
55		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-{[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl}]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
56		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-{[1-(4-carbamimidoyl-furan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl}]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
57		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[2-(2-carboxypropan-2-yl)oxy]imino)acetyl]amino]-3-[[1-({3-[(diaminomethylidene)amino]-1,2-oxazol-5-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
58		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
59		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-[[4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
60		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-[[4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
61		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-[[4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
62		Preparation of (6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[[1-[[4-(N-phenylcarbamimidoyl)thiophen-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
63		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[[1-[[4-(pyrrolidin-1-ylcarbamimidoyl)thiophen-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
64		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[[1-[[4-[N-(1,3-triazol-2-yl)carbamimidoyl]thiophen-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
65		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-[[2-(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-8-oxo-3-[[1-[[4-(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
66		(6R,7R)-7-[[[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
67		(6R,7R)-7-[[[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
68		(6R,7R)-7-[[[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
69		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued




Compound No.	Structure	Chemical Name
70		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
71		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
72		(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxyimino]acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate
73		(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

TABLE 1-continued

Compound No.	Structure	Chemical Name
74		(6R,7R)-7-[[[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[[[(2-carboxypropan-2-yl)oxy]-imino]acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

Methods of Preparation:

The compounds of general formula (I) can be prepared as described in the following schemes (Scheme 1 and Scheme 2) which illustrate the general method of preparation only and are not intended to be limiting to any specific compound described herein.



Process I (Scheme 1):

(a) Coupling Step:

Reaction of (VIII, $q=0$, $Y=\text{chloride}$) with the intermediate (IX) is carried out in a conventional solvent such as water, alcohol (e.g., methanol, ethanol etc.), acetone, dioxane, acetonitrile, chloroform, methylene chloride, ethylene chloride, tetrahydrofuran, ethylacetate, N,N-dimethylformamide, pyridine or any other organic solvent which does not adversely influence the reaction. These conventional solvents may also be used as a co-solvent in a mixture with water. In this reaction, when $Y=\text{Cl}$, the reaction is preferably carried out in the presence of a conventional alkali metal halide such as potassium iodide and in the presence of N,N-dimethylformamide or dimethylacetamide. The reaction is usually carried out ranging from -20°C . to 40°C .


(b) Hydrolysis Step:

The hydrolysis is preferably carried out in the presence of a base or an acid including Lewis acid. Suitable base may include an inorganic base such as an alkali metal (e.g., sodium, potassium etc.); an alkali earth metal (e.g., magnesium, calcium etc.), the hydroxide or carbonate or bicarbonate thereof, and an organic base trialkylamine (e.g., trimethylamine, triethylamine, etc.), picoline or the like. Suitable acid may include an organic acid (e.g., formic acid, acetic acid, propionic acid, trichloroacetic acid, trifluoroacetic acid) and an inorganic acid (e.g., hydrochloric acid, hydrobromic acid, sulfuric acid). The hydrolysis using Lewis acid such as trihaloacetic acid (e.g., trichloroacetic acid, trifluoroacetic acid) or the like and the reaction is preferably carried out in the presence of cation trapping agents (e.g., anisole, phenol etc.). The hydrolysis using Lewis acid such

95

as aluminum trichloride is carried out in a solvent like nitromethane. The hydrolysis reaction is usually carried out in a non-aqueous solvent such as methylene chloride, tetrahydrofuran, a mixture of solvents thereof, or any other solvent that does not adversely influence the reaction. The reaction temperature may be room temperature, or any other temperature as may be appropriate.

Scheme 2



Process II (Scheme 2):

(a) Coupling Step:

Reaction of (VIII, $q=1$, $Y=\text{iodide}$) with the intermediate (IX) is carried out in a conventional solvent such as water, alcohol (e.g., methanol, ethanol etc.), acetone, dioxane, acetonitrile, chloroform, methylene chloride, ethylene chloride, tetrahydrofuran, ethyl acetate, N,N-dimethylformamide, pyridine or any other organic solvent which does not adversely influence the reaction. These conventional solvents may also be used as a co-solvent in a mixture with water. In this reaction, when $Y=\text{Cl}$, the reaction is preferably carried out in the presence of a conventional alkali metal halide such as potassium iodide and in the presence of N,N-dimethylformamide or dimethylacetamide. The reaction temperature is not critical, and usually carried out under cooling to room temperature, even more preferably ranging from -20°C . to 40°C .

(b) Reduction Step:

Reduction is carried out in a conventional manner. Suitable reducing agents to be used in chemical reduction are a

96


combination of a metal (e.g., tin, zinc, iron etc.) and an organic or inorganic acid (e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.). Reduction can also be carried out using a combination of alkali metal halide (e.g., potassium iodide) and acetyl chloride. The reduction is usually carried out in a conventional solvent which does not adversely influence the reaction such as N,N-dimethylformamide, diethyl ether, dioxane, tetrahydrofuran, etc. or a mixture of solvents thereof. The reduction is carried out under cooling to warming, more specifically in the range from -40°C . to 0°C .

(c) Hydrolysis Step:

The hydrolysis is preferably carried out in the presence of a base or an acid including Lewis acid. Suitable base may include an inorganic base such as an alkali metal (e.g., sodium, potassium etc.); an alkali earth metal (e.g., magnesium, calcium etc.), the hydroxide or carbonate or bicarbonate thereof, and an organic base, trialkylamine (e.g., trimethylamine, triethylamine, etc.), picoline or the like. Suitable acid may include an organic acid (e.g., formic acid, acetic acid, propionic acid, trichloroacetic acid, trifluoroacetic acid) and an inorganic acid (e.g., hydrochloric acid, hydrobromic acid, sulfuric acid). The hydrolysis using Lewis acid such as trihaloacetic acid (e.g., trichloroacetic acid, trifluoroacetic acid) or the like and the reaction is preferably carried out in the presence of cation trapping agents (e.g., anisole, phenol etc.). The hydrolysis using Lewis acid such as aluminum trichloride is carried out in a solvent like nitromethane. The hydrolysis reaction is usually carried out in a non-aqueous solvent such as methylene chloride, tetrahydrofuran, a mixture of solvents thereof or any other solvent that does not adversely influence the reaction. The reaction temperature is not critical, but is usually carried out at room temperature.

In the formula (VIII), A and B have the same definitions as described before.

Thus, in formula (VIII), A is defined by the formula (Ia);




(Ia)

Object (Ia) includes syn isomer (Z form) anti isomer (E form) and a mixture thereof.

Wherein X is N, C(H), C(F) or C(Cl);

B is defined as hydrogen, methyl, ethyl or represented by the formula (Ib)



(Ib)

Wherein, R^1 and R^2 is independently hydrogen or lower alkyl, or R^1 and R^2 together may form a 3 to 6-membered spiro ring system; and

m is 0 or 1.


Furthermore, in the formula (Ib), R^1 and R^2 together may form a 3- to 6-membered spiro ring system;

Y is halogen, even more preferably chloro or iodo;

q is 0 or 1;


97

In the organic residue as designated by (IX), objects C, D, E and F have the same definitions as described before, thus C is selected from the following quaternized bicyclic aromatic heterocyclic rings represented by the formulae (Ic) to (Iz);











98

-continued




99

-continued




D represents CH_2 , CH_2CH_2 or CH_2CO ;

E is selected from aryl or a 5- to 6-membered aromatic heterocyclic ring as shown below;



100

-continued



F is optionally substituted amidine or an optionally substituted guanidine; and

(Iu) G is hydrogen, methyl, ethyl, C_{3-6} alkyl, C_{3-6} cycloalkyl or an optionally substituted 5- or 6-membered aliphatic or an optionally substituted 5- or 6-membered aromatic heterocyclic ring, in which the heterocyclic ring is substituted with at least 1-2 hetero atoms selected from N, O, and S (α or β).

(Iv) In the formula (VIII), P^1 is a suitable carboxy protecting group frequently used in (3-lactam chemistry. Suitable examples may be the ones such as lower alkyl ester (e.g., methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, t-pentyl, hexyl, 1-cyclopropylethyl etc.); lower alkenyl ester (e.g., vinyl, allyl); lower alkynyl ester (e.g., ethynyl, propynyl); lower alkoxyalkyl ester (e.g., methoxymethyl, ethoxymethyl, isopropoxymethyl, 1-methoxyethyl, 1-ethoxyethyl etc.); lower alkylthioalkyl ester (e.g., methylthiomethyl, ethylthiomethyl, ethylthioethyl, isopropylthiomethyl); mono (or di or tri-) halo(lower) alkyl ester (e.g., 2-iodoethyl, 2,2,2-trichloroethyl etc.); lower alkanoyloxy(lower)alkyl ester (e.g., acetoxymethyl, propionyloxymethyl, butyryloxymethyl, valeryloxymethyl, pivaloyloxymethyl, hexanoyloxymethyl, 2-acetoxyethyl, 2-propionyloxyethyl etc.); lower alkanesulfonyl(lower)alkyl ester (e.g., mesylmethyl ester, 2-mesyloxyethyl ester etc.); ar(lower)alkyl ester, for example, phenyl(lower)alkyl ester which may have one or more suitable substituent(s) (e.g., benzyl, 4-methoxybenzyl, 4-nitrobenzyl, phenethyl, trityl, benzhydryl, bis(methoxyphenyl)methyl, 3,4-dimethoxybenzyl, 4-hydroxy-3,5-di-t-butylbenzyl, etc.); aryl ester which may have one or more suitable substituent(s) such as substituted or unsubstituted phenyl ester (e.g., phenyl, tolyl, t-butyl phenyl, xylyl, mesityl, 4-chlorophenyl, 4-methoxyphenyl, etc.); tri(lower)alkyl silyl ester; lower alkylthioester (e.g., methylthioester, ethylthioester, etc.) and the like.

(Ix) It is understood that in the formula (VIII), when an amino group is present in the molecule, it is to be protected with a suitable protecting group commonly used in the β -lactam chemistry such as benzyl, trityl, t-butoxycarbonyl or the like. Similarly, it is to be understood that in the formula (VIII), when a carboxyl group is present in the molecule, it is to be protected with a suitable protecting group selected from the group as described for P^2 above.

(Iz) The following are provided for illustrative purposes only, and are not intended to be limiting of the present compositions and methods in any way.

EXAMPLES

60 In the examples the following abbreviations have been used.

Boc: N-tert-butoxycarbonyl

Br s: broad singlet

CDCl_3 : deuterated chloroform

65 CD_3OD : deuterated methanol

CH_3NO_2 : nitromethane


D_2O : deuterium oxide

101


d: doublet
 DCM: dichloromethane
 DMAP: 4-dimethylaminopyridine
 DMF: N, N'-dimethylformamide
 DMSO-d₆: deuterated dimethylsulfoxide
 ES: electron spray
 g: gram(s)
 h: hour(s)
 HPLC: high-performance liquid chromatography
 Hz: Hertz
 J: coupling constant
 m: multiplet
 mmol: millimole(s)
 MHz: megahertz
 MS: mass spectrometry

Example 1 (Table 1, Compound 1)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]-amino]-3-[[1-(4-carbamimidoylbenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



Step 1: 4-(1H-Pyrrolo[3,2-c]pyridin-1-ylmethyl) benzonitrile



To a solution of 1H-pyrrolo[3,2-c]pyridine (1.0 g, 8.5 mmol) in DMF (40 mL) under nitrogen at 0° C. was added NaH (60% in mineral oil, 0.51 g, 12.8 mmol) in portions and after stirring for 15 min was added 4-(bromomethyl)benzonitrile (2.5 g, 12.8 mmol) in portions. The reaction mixture was stirred at 0° C. for 2 h then at room temperature for 1 h and quenched with saturated ammonium chloride solution then diluted with ethyl acetate (100 mL) and water (100 mL). The organic layer was separated, washed with brine (3x100 mL), dried (Na₂SO₄) and concentrated.


The crude product was purified by silica gel column chromatography using methylene chloride: ethyl acetate:

102

methanol (5:3:2) to afford 4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzonitrile (1.2 g, 60%) as a solid.

¹H NMR (400 MHz, CDCl₃): δ 5.38 (s, 2H), 6.68 (d, 1H, J=3.2 Hz), 7.11-7.17 (m, 4H), 7.54 (d, 2H, J=8.0 Hz), 8.27 (d, 1H, J=5.6 Hz), 8.94 (s, 1H).

Step 2: Methyl 4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride




Through a solution of 4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzonitrile (from step 1, 0.7 g, 3.0 mmol) in anhydrous methanol (25 mL) in a pressure vessel at 0° C. was bubbled a stream of anhydrous hydrogen chloride gas

for 10 min. The reaction vessel was stoppered and stirred at room temperature for 18 h and the contents were transferred into a flask and evaporated under reduced pressure. The crude product was triturated with ether, then dried under vacuum to provide methyl 4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (0.85 g, 94%) as a solid.

¹H NMR (400 MHz, CD₃OD): δ 4.34 (s, 3H), 5.50 (s, 2H), 7.20 (s, 1H), 7.50 (d, 2H, J=8.0 Hz), 7.96-8.11 (m, 4H), 8.39 (d, 1H, J=6.8 Hz), 9.22 (s, 1H). One proton was not observed in CD₃OD.

Step 3: 4-(1H-Pyrrolo[3,2-c]pyridin-1-ylmethyl) benzenecarboximidamide





103

A pressure vessel containing a suspension of methyl 4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (from step 2, 0.9 g, 2.85 mol) in methanol (25 mL) was saturated with ammonia gas and stoppered. The reaction mixture was stirred at room temperature for 24 h, then excess ammonia was vented out and the contents were concentrated under reduced pressure. The residue was triturated with hexanes and dried under vacuum to give 4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidamide (1.0 g) as a solid.

¹H NMR (400 MHz, CD₃OD): δ 5.76 (s, 2H), 7.05 (d, 1H, J=3.2 Hz), 7.45 (d, 2H, J=8.0 Hz), 7.79-7.89 (m, 4H), 8.30 (s, 1H), 9.08 (s, 1H). Three protons were not observed in CD₃OD.

Step 4: tert-Butyl {[4-(1H-pyrrolo[3,2-c]pyridinylmethyl)phenyl]carboximidoyl}carbamate



104

A solution of 4-(1H-Pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidamide (from step 3, 0.25 g, 1.0 mmol) in 1,4-dioxane (30 mL) was treated with a saturated sodium carbonate solution (20 mL) followed by di-tert-butyl dicarbonate (1.1 g, 5.0 mmol) and stirred at room temperature for 40 h. The reaction mixture was concentrated under reduced pressure to remove the volatiles and the remaining solution was dissolved in ethyl acetate (60 mL), then washed with water (60 mL), brine solution (60 mL), dried and concentrated. The crude product was purified by silica gel column chromatography using ethyl acetate:methanol:ammonium hydroxide (60:39:1) as eluent to afford tert-butyl {[4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)phenyl]carboximidoyl}carbamate (0.12 g, 34%).


¹H NMR (400 MHz, CDCl₃): δ 1.54 (s, 9H), 5.36 (s, 2H), 6.60 (br s, 1H), 6.66 (d, 1H, J=3.6 Hz), 7.11-7.14 (m, 5H), 7.79 (d, 2H, J=8.4 Hz), 8.23 (d, 1H, J=6.0 Hz), 8.93 (s, 1H).

Step 5: 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-Butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[1-tert-butoxy-2-methyl-1-oxopropan-2-yl]oxy]imino}acetyl]amino]-2-[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl methyl]-1-4-[N-(tert-butoxycarbonyl)carbamimidoyl]benzyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide

To a solution of tert-butyl {[4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)phenyl]carboximidoyl}carbamate (from step 4, 0.07 g, 0.20 mmol) in dimethylacetamide (1.3 mL) was added 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[1-tert-butoxy-2-methyl-1-oxopropan-2-yl]oxy]imino}acetyl]amino]-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.156 g, 0.20 mmol) at 0° C. The reaction mixture was then degassed under reduced pressure for 0.5 h and was added sodium iodide (0.06 g, 0.40 mmol). After stirring at 15° C. for 16 h, the reaction mixture was slowly added to 5% aqueous sodium chloride and sodium thiosulfate solution cooled under ice. The suspension was then filtered, washed with water and the solid was vacuum dried to get a yellow solid (0.22 g) which was used in the next step without further purification.

105

Step 6: (6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




To a solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-{5-[(tert-butoxy-carbonyl)amino]-1,2,4-thiadiazol-3-yl}-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino]acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabi-
cyclo[4.2.0]oct-2-en-3-yl]methyl]-1-4-[N-(tert-butoxycar-
bonyl)carbamimidoyl]benzyl]-1H-pyrrolo[3,2-c]pyridin-5-
ium iodide (from step 5, 0.22 g) in dry dichloromethane (4.5
mL) at -40° C. was added anisole (0.36 mL, 3.34 mmol)
followed by 2M AlCl_3 in CH_3NO_2 (1.7 mL, 3.52 mmol). The

106

$J=14.4$ Hz), 5.53 (s, 2H), 5.63 (d, 1H, $J=4.8$ Hz), 6.76 (s, 1H), 6.94 (s, 1H), 7.22 (d, 2H, $J=7.2$ Hz), 7.55 (d, 2H, $J=7.6$ Hz), 7.68 (m, 2H), 8.23 (d, 1H, $J=7.2$ Hz), 9.06 (s, 1H). Seven protons were not observed in D_2O .

Example 2 (Table 1, Compound 9)


(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-2,3-dihydro-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



liquid was stirred at 0° C. for 30 min. To the reaction mixture were added di-isopropyl ether (5 mL) and water (0.5 mL), and the resultant mixture was stirred to generate a precipitate. The supernatant was removed by decantation. To the insoluble matter adhering to the vessel were added dilute aqueous hydrochloric acid solution (2 mL) and acetonitrile (5 mL) and was stirred to dissolve the matter completely. Thereto was added HP20 resin (0.5 g), stirred for 30 min. and then filtered. The filtrate was concentrated and freeze-dried to give a crude product, which was purified by HPLC to obtain (6R,7R)-7-[[[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (0.02 g, 15.5%).

^1H NMR (D_2O): δ 1.25 (s, 6H), 2.94 (d, 1H, $J=17.6$ Hz), 3.38 (d, 1H, $J=18.0$ Hz), 5.08-5.13 (m, 2H), 5.33 (d, 1H,

Step 1: 4-((2,3-Dihydro-1H-imidazo[1,2-b]pyrazol-1-yl)methyl)benzonitrile




To a solution of 2,3-dihydro-1H-imidazo[1,2-b]pyrazole (1.399 g, 12.84) in dimethyl formamide (10 mL) was added 4-(bromomethyl)benzonitrile (2.86 g, 12.84 mmol) followed by potassium carbonate (2.66 g, 19.26 mmol) and the reaction mixture was stirred under nitrogen 20 h, then

107

diluted with ethyl acetate (100 mL) and water (100 mL). The organic layer was separated, washed with water (3×100 mL) followed by brine solution (200 mL), dried over Na₂SO₄ and concentrated under reduced pressure. Purification of the crude product by silica gel column chromatography using ethyl acetate as eluent gave 4-((2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-yl)methyl)benzimidamide (2.1 g, 72.9%) as a solid.

¹H NMR (400 MHz, DMSO-d₆): δ 3.66 (t, 2H), 4.07 (t, 2H), 4.31 (s, 2H), 5.22 (d, 2H, J=2.0 Hz), 7.19 (d, 1H, J=1.5 Hz), 7.58 (d, 2H, J=8.2 Hz), 7.85 (d, 2H, J=8.2 Hz)


Step 2: Ethyl 4-(2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidate hydrochloride



Through a solution of 4-((2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-yl)methyl)benzimidamide (from step 1, 1.5 g, 6.69 mmol) in anhydrous methanol in a pressure vessel at 0° C. was bubbled a stream of anhydrous hydrogen chloride gas for 15 min. The reaction vessel was stoppered and stirred at room temperature for 20 h, the contents were transferred to a flask and evaporated in vacuo. The crude product was triturated with ether and dried under vacuum to get ethyl 4-(2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidate hydrochloride (1.8 g, 87.8%) as a solid.

¹H NMR (400 MHz, DMSO-d₆): δ 1.49 (t, 3H), 3.77 (dd, 2H, J=4.2 Hz), 4.17 (t, 2H), 4.43 (dd, 2H, 10.0 Hz), 4.63 (q, 2H, J=7.1 Hz), 5.51 (d, 1H, J=2.3 Hz), 6.00 (br s, 2H), 7.63 (d, 2H, J=8.0 Hz), 8.14 (d, 2H, J=8.0 Hz), 12.20 (br s, 2H).

Step 3: 4-(2,3-Dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidamide




108

A pressure vessel containing a suspension of ethyl 4-(2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidate hydrochloride (from step 2, 1.8 g, 5.87 mol) in methanol (25 mL) was saturated with ammonia gas and stoppered. The reaction mixture was stirred at room temperature for 24 h, then the excess ammonia was vented out and the contents were concentrated under reduced pressure. The residue was triturated with hexanes and dried under vacuum to give 4-(2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidamide (0.95 g, 66.9%) as a solid.

¹H NMR (400 MHz, DMSO-d₆): δ 3.66 (t, 2H), 4.06 (t, 2H), 4.31 (t, 2H), 5.22 (d, 1H, J=2.0 Hz), 7.17 (d, 1H, J=1.6 Hz), 7.71 (d, 2H, J=8.2 Hz), 7.83 (d, 2H, J=8.2 Hz)

Mass: ES⁺ 242.10

Step 4: tert-Butyl {[4-(2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)phenyl]carboximidoyl} carbamate




To a suspension of 4-(2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidamide (from step 3, 0.95 g, 3.94 mmol) in a mixture of sodium bicarbonate (1.60 g, 19.7 mmol), 1,4-dioxane (25 mL) and water (10 mL) was added di-tert-butyl dicarbonate (2.74 g, 12.54 mmol) and stirred at room temperature for 24 h. The reaction mixture was concentrated under reduced pressure to remove the volatiles and the remaining solution was dissolved in ethyl acetate (80 mL), washed with water (80 mL), brine solution (80 mL), dried and concentrated. The crude product was purified by silica gel column chromatography using ethyl acetate as eluent to afford tert-butyl {[4-(2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)phenyl]carboximidoyl} carbamate (0.45 g, 33.5%).

¹H NMR (400 MHz, DMSO-d₆): δ 1.42 (s, 9H), 3.63 (t, 2H), 4.24 (t, 2H), 5.18 (d, 1H, J=2.0 Hz), 7.16 (d, 1H, J=2.0 Hz), 7.45 (d, 2H, J=8.0 Hz), 7.92 (d, 2H, J=8.0 Hz).

109


110

Step 5: 5-[[[(6R,7R)-7-[[[(2Z)-2-[[[(tert-Butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[[4-[N-(tert-butoxycarbonyl)carbamimidoyl]benzyl]-2,3-dihydro-1H-imidazo[1,2-b]pyrazol-5-ium iodide



To a solution of 4-methoxybenzyl (6R,7R)-7-((Z)-2-(((1-(tert-butoxy)-2-methyl-1-oxopropan-2-yl)oxy)imino)-2-(((tert-butoxycarbonyl)amino)thiazol-4-yl)acetamido)-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.35 g, 0.449 mmol) in dimethylformamide (8 mL) at room temperature was added potassium iodide (0.149 g, 0.898 mmol) and after stirring for 10 min was added tert-butyl [[4-(2,3-dihydro-1H-imidazo[1,2-b]pyrazol-1-ylmethyl)phenyl]carbamimidoyl]carbamate (from step 4, 153 mg, 0.449 mmol). The reaction mixture was then stirred under nitrogen at room temperature for 16 h and diluted with a 1:1 mixture of sodium thio sulfate and brine (15 mL) solution. The suspension was then filtered, washed with water and the solid was vacuum dried to get a yellow solid (0.448 g), which was used in the next step without further purification.

Step 6: (6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-2,3-dihydro-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



5

30

35

40

45

To a solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-[[[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[[4-[N-(tert-butoxycarbonyl)carbamimidoyl]benzyl]-2,3-dihydro-1H-imidazo[1,2-b]pyrazol-5-ium iodide (from step 5, 0.45 g) in dry dichloromethane (15 mL) under nitrogen at room temperature was added anisole (0.8 mL) followed by trifluoroacetic acid (3 mL) in one portion and stirred for 3 h. The reaction mixture was concentrated under reduced pressure and the residue was suspended in distilled water (20 mL), stirred for 20 min and filtered. The filtrate was lyophilized and the product was purified by HPLC to obtain (6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-2,3-dihydro-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as a formic acid salt (0.022 g, 8.4%).


111

¹H NMR (400 MHz, D₂O): δ 1.33 (s, 6H), 3.08 (d, 1H, J=18.0 Hz), 3.31 (d, 1H, J=18.0 Hz), 3.86 (t, 2H), 4.12 (q, 1H, J=8.6 Hz), 4.22 (q, 1H, J=8.2 Hz), 4.44 (q, 2H, J=3.5 Hz), 4.85 (dd, 2H, J=15.6 and 19.7 Hz), 5.07 (d, 1H), 5.64 (d, 1H), 5.66 (d, 1H), 6.87 (s, 1H), 7.42 (d, 2H), 7.63 (d, 2H), 7.81 (1H, d), 8.17 (1H, s).


Mass: ES⁺ 709.19

Example 3 (Table 1, Compound 2)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]-amino]-8-oxo-3-[[1-(4-{N-[(3R)-piperidin-3-yl]carbamimidoyl}benzyl)-1H-pyrazolo[4,3-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



Step 1: 4-(1H-Pyrazolo[4,3-c]pyridin-1-ylmethyl) benzonitrile





A mixture of 1H-pyrazolo[4,3-c]pyridine (0.72 g, 6.04 mmol), 4-(bromomethyl)benzonitrile (1.57 g, 8.01 mmol) and potassium carbonate (1.67 g, 12.08 mmol) in DMF (75 mL) was stirred at 60° C. for 24 h and concentrated. The residue was partitioned between ethyl acetate (100 mL) and water (100 mL). The organic layer was separated, washed with brine (3×100 mL), dried (Na₂SO₄) and concentrated. The crude product was purified by silica gel column chromatography using ethyl acetate: methanol (9:1) as eluent to afford 4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)benzonitrile (0.42 g, 30.5%) as a solid.

¹H NMR (400 MHz, CDCl₃): δ 5.66 (s, 2H), 7.24-7.28 (m, 3H), 7.61 (d, 2H, J=8.4 Hz), 8.23 (s, 1H), 8.43 (d, 1H, J=6.4 Hz), 9.15 (s, 1H).

112


Step 2: Ethyl 4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride



Through a solution of 4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)benzonitrile (from step 1, 0.7 g, 3.0 mmol) in anhydrous ethanol (30 mL) in a pressure vessel at 0° C. was bubbled a stream of anhydrous hydrogen chloride gas for 10 min. The reaction vessel was stoppered, then stirred at room temperature for 18 h and the contents were transferred into a flask and evaporated under reduced pressure. The crude product was triturated with ether and dried under vacuum to provide ethyl 4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (0.8 g, 84%) as a solid.

¹H NMR (400 MHz, CD₃OD): δ 1.59 (t, 3H, J=6.6 Hz), 4.63 (q, 2H, J=6.2 Hz), 6.03 (s, 2H), 7.60 (d, 2H, J=6.8 Hz), 8.04 (d, 2H, J=7.2 Hz), 8.37 (s, 1H), 8.58 (s, 1H), 8.84 (s, 1H), 9.59 (s, 1H). One proton was not observed in CD₃OD.

Step 3: tert-Butyl (3R)-3-([4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)phenyl]carbamimidoyl)amino piperidine-1-carboxylate





113

To a solution of ethyl 4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (from step 2, 0.57 g, 1.8 mmol) and triethylamine (0.56 mL, 4.0 mmol) in methanol (5 mL) was added tert-butyl (3R)-3-aminopiperidine-1-carboxylate (0.36 g, 1.8 mmol) in methanol (1 mL) at 0° C. The reaction mixture was stirred at room temperature for 2 days and concentrated. The residue was purified by column chromatography using methanol: ethyl acetate (1:4) as eluent to give tert-butyl (3R)-3-([4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)phenyl]carboximidoyl)amino]piperidine-1-carboxylate (0.5 g, 64%) as a solid.

¹H NMR (400 MHz, CDCl₃): δ 1.28 (s, 9H), 1.57-1.80 (m, 3H), 2.12 (s, 1H), 3.08-3.30 (m, 2H), 3.36-4.10 (m, 3H), 5.83 (s, 2H), 7.45 (d, 2H, J=8.8 Hz), 7.67-7.69 (m, 3H), 8.33 (d, 1H, J=6.0 Hz), 8.37 (s, 1H), 9.10 (s, 1H). Two protons were not observed in CD₃OD.

Step 4: tert-Butyl (3R)-3-[(tert-butoxycarbonyl){4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)phenyl}carboximidoyl]amino]piperidine-1-carboxylate



114

A solution of tert-butyl (3R)-3-([4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)phenyl]carboximidoyl)amino]piperidine-1-carboxylate (from step 3, 0.5 g, 1.15 mmol) in 1,4-dioxane (20 mL) was treated with a saturated sodium carbonate solution (10 mL) followed by di-tert-butyl dicarbonate (1.26 g, 5.78 mmol), then stirred at room temperature for 40 h. The reaction mixture was concentrated under reduced pressure to remove the volatiles and the remaining solution was dissolved in ethyl acetate (100 mL) and washed with water (100 mL), brine solution (100 mL), dried, and concentrated. The crude product was purified by silica gel column chromatography using ethyl acetate: methanol: ammonium hydroxide (60:39:1) as eluent to afford tert-butyl (3R)-3-[(tert-butoxycarbonyl){4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)phenyl]carboximidoyl]amino]piperidine-1-carboxylate (0.48 g, 78%).

¹H NMR (400 MHz, CDCl₃): δ 1.24-2.05 (m, 22H), 3.05-3.71 (m, 4H), 4.09-4.10 (m, 1H), 5.08 (br s, 1H), 5.62 (s, 2H), 7.22-7.28 (m, 3H), 7.39 (d, 2H, J=8.4 Hz), 8.19 (s, 1H), 8.38 (d, 1H, J=6.0 Hz), 9.12 (s, 1H).


Step 5: 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-2-[[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-(4-{N-(tert-butoxycarbonyl)-N-[(3R)-1-(tert-butoxycarbonyl)piperidin-3-yl]carbamimidoyl}benzyl)-1H-pyrazolo[4,3-c]pyridin-5-ium iodide

50

To a solution of tert-butyl (3R)-3-[(tert-butoxycarbonyl){4-(1H-pyrazolo[4,3-c]pyridin-1-ylmethyl)phenyl}carboximidoyl]amino]piperidine-1-carboxylate (from step 4, 0.16 g, 0.30 mmol) in dimethylacetamide (2 mL) was added 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{5-[(tert-butoxycarbonyl)amino]-1,2,4-thiadiazol-3-yl]-2-[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.23 g, 0.30 mmol) at 0° C. The reaction mixture was then degassed under reduced pressure for 0.5 h, and thereto was added sodium iodide (0.09 g, 0.60 mmol). After stirring at 15° C. for 16 h, the reaction mixture was slowly added to 5% aqueous sodium chloride and sodium thiosulfate solution cooled under ice. The suspension was then filtered, washed with water and the solid was vacuum dried to get a yellow solid (0.33 g) which was used in the next step without further purification.

115

Step 6: (6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-8-oxo-3-[[1-(4-{N-[(3R)-piperidin-3-yl]carbamimidoyl}benzyl)-1H-pyrazolo[4,3-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.1]oct-2-ene-2-carboxylate



To a solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-[[[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino]acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-(4-{N-(tert-butoxycarbonyl)-N-[(3R)-1-(tert-butoxycarbonyl)piperidin-3-yl]carbamimidoyl}benzyl)-1H-pyrazolo[4,3-c]pyridin-5-ium iodide (from step 5, 0.33 g) in dry dichloromethane (6 mL) at -40°C . was added anisole (0.50 mL, 4.64 mmol) and then 2M aluminium chloride in nitromethane (2.33 mL, 4.66 mmol). The liquid was stirred at 0°C . for 30 min. To the reaction mixture were added di-isopropyl ether (10 mL) and water (0.8 mL), and the resultant was stirred to generate a precipitate. The supernatant was removed by decantation. To the insoluble matter adhering to the vessel were added dilute aqueous hydrochloric acid solution (4 mL) and acetonitrile (15 mL). The resultant was stirred to dissolve the matter completely. Thereto was added HP20 resin (0.8 g), and stirred for 30 min, and then filtered. The filtrate was concentrated and freeze-dried to give a crude product which was purified by HPLC to obtain (6R,7R)-7-[[[(2Z)-2-(2-amino-


1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-8-oxo-3-[[1-(4-{N-[(3R)-piperidin-3-yl]carbamimidoyl}benzyl)-1H-pyrazolo[4,3-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as a formic acid salt (0.025 g, 12.5%).

$^1\text{H NMR}$ (400 MHz, D_2O): δ 1.17 (s, 6H), 1.58-1.70 (m, 2H), 1.88-1.92 (m, 1H), 2.08-2.09 (m, 1H), 2.85-2.98 (m, 3H), 3.20 (d, 1H, $J=12.0$ Hz), 3.45-3.49 (m, 2H), 3.96 (m, 1H), 5.08-5.11 (m, 2H), 5.43 (d, 1H, $J=14.8$ Hz), 5.58 (d, 1H, $J=4.4$ Hz), 5.72 (s, 2H), 6.65 (s, 1H), 7.26 (d, 2H, $J=8.0$ Hz), 7.50 (d, 2H, $J=7.2$ Hz), 7.90 (d, 1H, $J=6.8$ Hz), 8.50 (d, 1H, $J=7.2$ Hz), 8.58 (s, 1H), 9.49 (s, 1H). Seven protons were not observed in D_2O .

Mass: ES^+ 802.20


Example 4 (Table 1, Compound 5)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



117


Step 1: 4-(1H-Imidazo[1,2-b]pyrazol-1-ylmethyl)benzonitrile



To a solution of 1H-imidazo[1,2-b]pyrazole (2.158 g, 20.146 mmol) in DMF (30 mL) under nitrogen at 0° C. was added sodium hydride (60.9% in mineral oil, 1.19 g, 30.22 mmol) in portions and after stirring for 15 min was added 4-(bromomethyl)benzonitrile (3.95 g, 20.146 mmol) in portions. The reaction mixture was stirred at 0° C. for 2 h then at room temperature for 1 h and quenched with saturated ammonium chloride solution, then diluted with ethyl acetate (200 mL) and water (200 mL). The organic layer was separated, washed with water (3×200 mL), brine solution, dried (Na₂SO₄) and concentrated. The crude product was purified by silica gel column chromatography using ethyl acetate to afford 4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzonitrile (2.2 g, 49%) as a solid.

¹H NMR (400 MHz, DMSO-d₆): δ 5.25 (s, 2H), 5.62 (d, 1H, J=2.7 Hz), 7.34-7.35 (m, 1H), 7.44-7.47 (m, 3H), 7.59-7.60 (m, 1H), 7.84 (d, 2H, J=8.2 Hz).

Step 2: Ethyl 4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidate hydrochloride




Through a solution of 4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzonitrile (from step 1, 1.2 g, 5.4 mmol) in anhydrous ethanol (25 mL) in a pressure vessel at 0° C. was bubbled a stream of anhydrous hydrogen chloride gas for 15 min. The reaction vessel was stoppered and stirred at room temperature for 18 h, then the contents were transferred into a flask and evaporated under reduced pressure. The crude product was triturated with ether and dried under vacuum to provide ethyl 4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidate hydrochloride (1.6 g, 97%) as a solid.

¹H NMR (400 MHz, DMSO-d₆): δ 1.46 (t, 3H, J=7.1 Hz), 4.73 (q, 2H, J=6.9 Hz), 5.41 (s, 2H), 5.98 (d, 1H, J=2.8 Hz), 7.56-7.61 (m, 4H), 7.80 (d, 2H, J=8.6 Hz), 8.14 (d, 2H, J=8.6 Hz), 12.20 (br s, 2H).

118


Step 3: 4-(1H-Imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidamide



A pressure vessel containing a suspension of ethyl 4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidate hydrochloride (from step 2, 1.1 g, 3.61 mol) in methanol (25 mL) was saturated with ammonia gas and stoppered. The reaction mixture was stirred at room temperature for 24 h, then excess ammonia was vented out and the contents were concentrated under reduced pressure. The residue was triturated with hexanes and dried under vacuum to give 4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidamide (1.5 g, 141.5%) as a solid.

¹H NMR (400 MHz, DMSO-d₆): δ 5.27 (s, 2H), 5.64 (d, 1H, J=2.4 Hz), 7.35-7.60 (m, 5H), 7.80-7.86 (m, 2H)

Step 4: tert-Butyl {[4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)phenyl]carbonoimidoyl}carbamate




To a suspension of 4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)benzenecarboximidamide (from step 3, 1.50 g, 6.27 mmol) in a mixture of sodium bicarbonate (1.524 g, 18.81 mmol), 1,4-dioxane (30 mL) and water (10 mL) was added di-tert-butyl dicarbonate (2.74 g, 12.54 mmol) and stirred at room temperature for 20 h. The reaction mixture was concentrated under reduced pressure to remove the volatiles and the remaining solution was dissolved in ethyl acetate (80 mL), washed with water (80 mL), brine solution (80 mL), dried and concentrated. The crude product was purified by silica gel column chromatography using ethyl acetate as eluent to afford tert-butyl {[4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)phenyl]carbonoimidoyl}carbamate (0.60 g, 28.17%).

¹H NMR (400 MHz, DMSO-d₆): δ 1.41 (s, 9H), 5.18 (s, 2H), 5.58 (d, 1H, J=2.3 Hz), 7.32-7.37 (m, 3H), 7.41-7.42 (m, 1H), 7.56 (d, 1H, J=2.8 Hz), 7.88 (d, 2H, J=8.5 Hz).

Mass: ES⁻ 338.11-


119

Step 5: 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-Butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-{4-[N-(tert-butoxycarbonyl)carbamimidoyl]benzyl}]-1H-imidazo[1,2-b]pyrazol-5-ium iodide



To a solution of 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl] amino}-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0] oct-2-ene-2-carboxylate (0.35 g, 0.449 mmol) in dimethylformamide (8 mL) at room temperature was added potassium iodide (0.149 g, 0.898 mmol) and after stirring for 10 min was added tert-butyl {4-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)phenyl}carbamimidoyl} carbamate (from step 4, 152 mg, 0.449 mmol). The reaction mixture was then stirred under nitrogen at room temperature for 16 h and diluted with a 1:1 mixture of sodium thio sulfate and brine solution (15 mL). The suspension was then filtered, washed with water and the solid was vacuum dried to get a yellow solid (0.650 g), which was used in the next step without further purification.

Step 6: (6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino}acetyl] amino]-3-[[1-(4-carbamimidoylbenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



120

To a solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo [4.2.0]oct-2-en-3-yl]methyl]-1-{4-[N-(tert-butoxycarbonyl)carbamimidoyl]benzyl}]-1H-imidazo[1,2-b]pyrazol-5-ium iodide (from step 5, 0.65 g) in dry

dichloromethane under nitrogen at room temperature was added trifluoroacetic acid (5 mL) in one portion and stirred for 3 h. The reaction mixture was concentrated under reduced pressure and the residue was suspended in distilled water (20 mL), stirred for 20 min then filtered. The filtrate was lyophilized and the product was purified by HPLC to obtain (6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-3-[[1-(4-carbamimidoylbenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (0.032 g, 7.9%).


¹H NMR (400 MHz, DMSO-d₆): δ 1.29 (d, 6H, J=8.2 Hz), 2.94 (d, 1H, J=18.0 Hz), 3.25 (d, 1H, J=18 Hz), 5.05 (d, 1H, J=5.0 Hz), 5.14 (s, 2H), 5.32 (s, 2H), 5.63 (d, 1H, J=4.7 Hz), 6.13 (d, 1H, J=3.5 Hz), 6.86 (s, 1H), 7.37-7.41 (m, 3H), 7.63 (d, 2H, J=8.6 Hz), 7.87 (d, 2H, J=3.2 Hz), 8.16 (s, 1H)

Mass: ES⁺ 707.17


121

Example 5 (Table 1, Compound 11)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




Step 1: (4-Bromothiophen-2-yl)methanol



Sodium borohydride (2.08 g, 55.0 mmol) was added to a solution of 4-bromothiophene-2-carbaldehyde (10.0 g, 52.3 mmol) in anhydrous tetrahydrofuran (200 mL) at room temperature and the resulting reaction mixture was stirred at room temperature for 1.5 h. Then the reaction mixture was quenched carefully by adding saturated aq. ammonium chloride solution (50 mL). The reaction mixture was extracted with ethyl acetate (200 mL). Organic phase was separated, dried over Na₂SO₄. Solvent was evaporated, crude compound was dried under vacuum to give (4-bromothiophen-2-yl)methanol (10.56 g, crude) as white semi-solid which was used in the next step without further purification.

¹H NMR (400 MHz, CDCl₃): δ 4.79 (s, 2H), 6.92 (s, 1H), 7.17 (s, 1H).

Step 2: 5-(Hydroxymethyl)thiophene-3-carbonitrile




A mixture of (4-bromothiophen-2-yl)methanol (from step 1, 10.56 g crude, 52.3 mmol), zinc cyanide (6.14 g, 52.34 mmol) in a 500 mL round bottom flask was degassed three times with nitrogen. Then anhydrous dimethylformamide (75 mL) was added, and the mixture was degassed with nitrogen. After that was added Palladium-tetrakis(triphenylphosphine) (3.01 g, 2.61 mmol) and degassing was repeated once again. The flask was fitted with a reflux condenser, and the reaction mixture was stirred at 80° C. (bath temperature)

122

for 4 h under nitrogen. The mixture was cooled to room temperature, solvent was evaporated. The crude compound was purified by flash chromatography using RediSep silica 120 g flash column (10-30% ethyl acetate in hexanes as eluent) to give 5-(hydroxymethyl)thiophene-3-carbonitrile (5.15 g, 71% over two steps) as white solid.

¹H NMR (400 MHz, CDCl₃): δ 2.26 (t, 1H, J=5.9 Hz), 4.84 (d, 2H, J=5.9 Hz), 7.16 (s, 1H), 7.88 (d, 1H, J=1.6 Hz).


Step 3: 5-(Bromomethyl)thiophene-3-carbonitrile



A solution of 5-(hydroxymethyl)thiophene-3-carbonitrile (from step 2, 5.15 g, 37.0 mmol) in anhydrous tetrahydrofuran (150 mL) was added triphenyl phosphine (10.19 g, 38.8 mmol) followed by carbon tetrabromide (12.88 g, 38.8 mmol). The reaction mixture was stirred at room temperature for 20 h under nitrogen. Solvent was evaporated and the crude compound was purified by flash chromatography using RediSep silica 120 g flash column (0-10% ethyl acetate in hexanes as eluent) to give 5-(bromomethyl)thiophene-3-carbonitrile (6.07 g, 81%) as a white solid.

¹H NMR (400 MHz, CDCl₃): δ 4.66 (s, 2H), 7.28 (d, 1H, J=0.8 Hz), 7.92 (d, 1H, J=1.6 Hz).

Step 4: 5-(1H-Pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carbonitrile




To a solution of 1H-pyrrolo[3,2-c]pyridine (1.77 g, 15.00 mmol) in anhydrous dimethylformamide (75 mL) was added sodium hydride (60% suspension in mineral oil, 0.75 g, 18.75 mmol) at 0° C. in small portions under nitrogen. After the addition, reaction mixture was stirred for 15 minutes at 0° C. and then at room temperature for 10 minutes. Cooled to 0° C., 5-(bromomethyl)thiophene-3-carbonitrile (from step 3, 3.79 g, 18.75 mmol) was added in small portions. After the addition, reaction mixture was stirred for 10 minutes at 0° C. and then at room temperature for 1 h. Quenched with water (10 mL), solvent was evaporated under reduced pressure. Residue was partitioned between ethyl acetate (150 mL) and water (100 mL). Organic phase was separated, washed with brine and dried over Na₂SO₄. The crude compound was purified by flash chromatography using RediSep silica 80 g flash column (0-3% methanol in methylene chloride as eluent) to give 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carbonitrile (2.17 g, 60%) as a brown solid.

123

¹H NMR (400 MHz, CDCl₃): δ 5.47 (s, 2H), 6.68 (d, 1H, J=3.1 Hz), 7.10 (s, 1H), 7.15 (d, 1H, J=3.5 Hz), 7.23 (d, 1H, J=5.8 Hz), 7.83 (s, 1H), 8.35 (d, 1H, J=5.9 Hz).

Mass: ES⁺ 240.01


Step 5: Methyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride



Anhydrous hydrogen chloride gas was bubbled through a solution of 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carbonitrile (from step 4, 1.05 g, 4.38 mmol) in anhydrous methanol (30 mL) at room temperature for 45 minutes. The reaction mixture was stirred at room temperature for 20 h. Solvent was evaporated and dried under vacuum to give methyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride (1.05 g crude) as brown color gummy solid which was used in next step without further purification.

Mass: ES⁺ 272.06.

Step 6: 5-(1H-Pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidamide



Anhydrous ammonia gas was bubbled through a solution of methyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride (from step 5, 1.05 g crude, ~4.3 mmol) in anhydrous methanol (20 mL) at room


124

temperature for 45 minutes. The reaction mixture was stirred at room temperature for 16 h. Solvent was evaporated and dried under vacuum to give 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidamide (1.72 g crude) as a brown solid which was used in the next step without further purification.

¹H NMR (400 MHz, DMSO-d₆): δ 5.90 (br s, 2H), 7.00 (br s, 1H), 7.42 (br s, 2H), 7.85 (s, 1H), 8.00 (br s, 1H), 8.19 (br s, 1H), 8.43 (br s, 1H), 8.65 (s, 1H), 9.19 (br s, 1H), 9.52 (br s, 1H).

Mass: ES⁺ 257.05.

Step 7: tert-Butyl {[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophen-3-yl]carboximidoyl}carbamate




To a solution of 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidamide (from step 6, 1.72 g crude, ~4.0 mmol) in 1,4-dioxane (50 mL) was added aqueous saturated sodium bicarbonate solution followed by di-tert-butyl dicarbonate (3.50 g, 16.0 mmol). The reaction mixture was stirred at room temperature for 4 h. Solvent was evaporated, water (50 mL) was added and extracted with methylene chloride (2x50 mL). The organic phase was dried over Na₂SO₄. Solvent was evaporated and the crude compound was purified by flash chromatography using RediSep silica 40 g flash column (0-2% methanol in methylene chloride as eluent) to give tert-butyl {[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophen-3-yl]carboximidoyl}carbamate (0.17 g, 11% for three steps) as an off-white solid.

¹H NMR (400 MHz, CDCl₃): δ 1.53 (s, 9H), 5.42 (s, 2H), 6.64 (d, 1H, J=2.7 Hz), 7.15 (d, 1H, J=3.5 Hz), 7.23 (d, 1H, J=5.5 Hz), 7.41 (s, 1H), 7.86 (s, 1H), 8.28 (d, 1H, J=5.9 Hz), 8.89 (s, 1H).

Mass: ES⁺ 355.10.

125


Step 8: 5-[[[(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino]acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide



A mixture of tert-butyl [[5-(1H-pyrrolo[3,2-c]pyridin-1-yl)methyl]thiophen-3-yl]carbamimidoyl carbamate (from step 7, 107 mg, 0.30 mmol), 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-[[2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino]acetyl]amino]-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (234 mg, 0.30 mmol), and sodium iodide (90 mg, 0.60 mmol) was cooled to 0° C. in an ice-bath; dimethyl formamide (3 mL) was added, stirred for 45 minutes, then the mixture was stirred at -16° C. for 16 h under nitrogen. Sodium bisulphite (0.6 g) in sodium chloride solution (75 mL) was added at 15° C., stirred for 10 minutes. The solid separated was filtered off, washed with water and dried under vacuum to give 5-(((6R,7R)-7-((Z)-2-(2-aminothiazol-4-yl)-2-(((1-tert-butoxy)-2-methyl-1-oxopropan-2-yl)oxy)imino)acetamido)-2-(((4-methoxybenzyl)oxy)carbonyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl)methyl)-1-((4-(N-(tert-butoxycarbonyl)carbamimidoyl)thiophen-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (296 mg, crude) as yellow color solid which was used in the next step without further purification.

Mass: ES⁺ 1100.52.

Step 9: (6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[4-carbamimidoyl]thiophen-2-yl)methyl}-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



126

To a solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino]acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (from step 8, 296 mg, crude, -0.30 mmol) in


anhydrous methylene chloride (3 mL) was added anisole (0.8 mL) followed by trifluoroacetic acid (2.0 mL). The reaction mixture was stirred at room temperature for 3 h. Solvent was evaporated and diisopropyl ether (30 mL) was added. The solid separated was filtered, washed with diisopropyl ether and dried under vacuum. The light brown color solid (356 mg) was dissolved in water (20 mL) and filtered. The aqueous phase was lyophilized to give yellow solid (200 mg) which was purified by preparative HPLC (acetonitrile, water, 0.1% formic acid) to give (6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[4-carbamimidoyl]thiophen-2-yl)methyl}-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as a formic acid salt (30 mg, 14%).

¹H NMR (400 MHz, D₂O): δ 1.24 (s, 3H), 1.26 (s, 3H), 2.94 (d, 1H, J=18.0 Hz), 3.42 (d, 1H, J=18.0 Hz), 5.01-5.14 (m, 2H), 5.39 (d, 1H, J=14.5 Hz), 5.59-5.76 (m, 3H), 6.75 (s, 1H), 6.95 (d, 1H, J=3.5 Hz), 7.31 (s, 1H), 7.73 (d, 1H, J=3.5 Hz), 7.88 (d, 1H, J=7.0), 8.09 (d, 1H, J=1.6 Hz), 8.37 (d, 1H, J=7.0 Hz), 9.10 (s, 1H).

Mass: ES⁺ 724.17.


Example 6 (Table 1, Compound 13)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[4-carbamimidoyl]furan-2-yl)methyl}-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



127


Step 1: (4-Bromofuran-2-yl)methanol



Sodium borohydride (1.21 g, 31.98 mmol) was added to a solution of 4-bromofuran-2-carbaldehyde (5.33 g, 30.46 mmol) in anhydrous tetrahydrofuran (120 mL) at room temperature and the resulting reaction mixture was stirred at room temperature for 1.5 h. Then the reaction mixture was quenched carefully by adding saturated aqueous ammonium chloride solution (50 mL) and extracted with ethyl acetate (200 mL). The organic phase was separated and dried over Na_2SO_4 . Solvent was evaporated and the crude compound was dried under vacuum to give (4-bromofuran-2-yl)methanol (5.34 g) as white semi-solid, which was used in the next step without further purification.

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 2.36 (br s, 1H), 4.55 (s, 2H), 6.34 (s, 1H), 7.39 (s, 1H).


Step 2: 5-(Hydroxymethyl)furan-3-carbonitrile



A mixture of (4-bromofuran-2-yl)methanol (from step 1, 3.50 g, 19.8 mmol) and zinc cyanide (2.32 g, 19.8 mmol) in a 250 mL round bottom flask was degassed three times with nitrogen. Anhydrous dimethylformamide (20 mL) was added, and the mixture was degassed with nitrogen. After that $\text{Pd}(\text{PPh}_3)_4$ (1.38 g, 1.20 mmol) was added and degassing was repeated once again. The flask was fitted with a reflux condenser, and the reaction mixture was stirred at 80°C . for 24 h under nitrogen. The mixture was cooled to room temperature, solvent was evaporated and residue was taken in ethyl acetate (150 mL), washed with water (100 mL), brine (100 mL) and dried over Na_2SO_4 . Solvent was evaporated and the crude compound was purified by flash chromatography using RediSep silica 120 g flash column (0-30% ethyl acetate in hexanes as eluent) to give 5-(hydroxymethyl)furan-3-carbonitrile (0.58 g, 24%) as a white solid.

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 1.97 (br s, 1H), 4.64 (d, 2H, $J=6.3$ Hz), 6.53 (s, 1H), 7.92 (s, 1H).

Step 3: 5-(Bromomethyl)furan-3-carbonitrile




A solution of 5-(hydroxymethyl)furan-3-carbonitrile (from step 2, 0.92 g, 7.47 mmol) in anhydrous tetrahydrofuran (30 mL) was added triphenyl phosphine (2.15 g, 8.22

128

mmol) followed by carbon tetrabromide (2.72 g, 8.22 mmol). The reaction mixture was stirred at room temperature for 4 h under nitrogen. Solvent was evaporated and the crude compound was purified by flash chromatography using RediSep silica 80 g flash column (10% ethyl acetate in hexanes as eluent) to give 5-(bromomethyl)furan-3-carbonitrile (1.12 g, 81%) as a white solid.

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 4.43 (s, 2H), 6.60 (s, 1H), 7.94 (s, 1H).

Step 4: 5-(1H-Pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-carbonitrile




To a solution of 1H-pyrrolo[3,2-c]pyridine (0.77 g, 6.50 mmol) in anhydrous dimethylformamide (30 mL) was added sodium hydride (60% suspension in mineral oil, 0.26 g, 6.50 mmol) at 0°C . in small portions under nitrogen. After the addition, the reaction mixture was stirred for 15 minutes at 0°C . and then at room temperature for 10 minutes then cooled to 0°C . and 5-(bromomethyl)furan-3-carbonitrile (from step 3, 1.10 g, 5.91 mmol) was added in small portions. After the addition, the reaction mixture was stirred for 10 minutes at 0°C . then at room temperature for 1 h. The mixture was quenched with water (5 mL). Solvent was evaporated under reduced pressure. Residue was partitioned between ethyl acetate (100 mL) and water (50 mL). The organic phase was separated, washed with brine and dried over Na_2SO_4 . The crude compound was purified by flash chromatography using RediSep silica 40 g flash column (0-3% methanol in methylene chloride as eluent) to give 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-carbonitrile (1.065 g, 80%) as an off-white solid.

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 5.29 (s, 2H), 6.46 (s, 1H), 6.66 (d, 1H, $J=2.3$ Hz), 7.15 (d, 1H, $J=3.5$ Hz), 7.27 (s, 1H), 7.88 (s, 1H), 8.37 (d, 1H, $J=5.9$ Hz), 8.94 (s, 1H).

Mass: ES^+ 224.10.

Step 5: Methyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-carboximidate hydrochloride




Anhydrous hydrogen chloride gas was bubbled through a solution of 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-

129

3-carbonitrile (from step 4, 1.05 g, 4.70 mmol) in anhydrous methanol (30 mL) at room temperature for 45 minutes; the reaction mixture was stirred at room temperature for 20 h. Solvent was evaporated and dried under vacuum to give methyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-carboximidate hydrochloride (1.05 g, crude) as a brown gummy solid. The compound was used in the next step without further purification. ¹H NMR (400 MHz, CD₃OD): δ 4.28 (s, 3H), 5.82 (s, 2H), 7.16 (br s, 1H), 7.22 (br s, 1H), 7.98 (br s, 1H), 8.30 (br s, 1H), 8.46 (br s, 1H), 8.54 (s, 1H), 9.20 (s, 1H).

Mass: ES⁺ 256.10.

Step 6: 5-(1H-Pyrrolo[3,2-c]pyridin-1-ylmethyl) furan-3-carboximidamide




Anhydrous ammonia gas was bubbled through a solution of methyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-carboximidate hydrochloride (from step 5, 1.0 g crude, ~4.7 mmol) in anhydrous methanol (30 mL) at room temperature for 45 minutes. The reaction mixture was stirred at room temperature for 16 h. Solvent was evaporated and dried under vacuum to give 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-carboximidamide (1.90 g, crude) as a brown solid.

The compound was used in the next step without further purification.

Mass: ES⁺ 241.11.

130

Step 7: tert-Butyl {[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-yl]carboximidoyl}carbamate




To a solution of 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl) furan-3-carboximidamide (from step 6, 1.90 g crude, ~4.7 mmol) in 1,4-dioxane (5 mL) was added aqueous saturated sodium bicarbonate solution (5 mL) followed by di-tert-butyl dicarbonate (4.10 g, 18.8 mmol) and the reaction mixture was stirred at room temperature for 4 h. Solvent was evaporated, water (50 mL) was added and extracted with methylene chloride (2x50 mL). The organic phase was dried over Na₂SO₄. Solvent was evaporated and the crude compound was purified by flash chromatography using RediSep silica 40 g flash column (2-5% methanol in methylene chloride as eluent) to give tert-butyl {[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-yl]carboximidoyl}carbamate (116 mg, 7% over steps) as an off-white solid.

¹H NMR (400 MHz, CDCl₃): δ 1.51 (s, 9H), 5.26 (s, 2H), 6.59 (s, 1H), 6.63 (dd, 1H, J=3.1, 0.8 Hz), 7.15 (d, 1H, J=3.5 Hz), 7.22 (d, 1H, J=5.9 Hz), 8.01 (s, 1H), 8.22 (d, 1H, J=5.9 Hz), 8.85 (s, 1H).

Mass: ES⁺ 341.23.

Step 8: 5-[[[(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino]acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-({4-[N-(tert-butoxycarbonyl)carbamimidoyl]furan-2-yl}methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium iodide





131

A mixture of tert-butyl {[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)furan-3-yl]carbamimidoyl}carbamate (from step 7, 110 mg, 0.32 mmol), 4-methoxybenzyl (6R,7R)-7-[[{(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]}-2-[[{(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (252 mg, 0.32 mmol), and sodium iodide (96 mg, 0.64 mmol) was cooled to 0° C. and dimethylformamide (3 mL) was added, stirred for 45 minutes then the mixture was stirred at -16° C. for 16 h under nitrogen. Sodium bisulphite (0.6 g) in sodium chloride solution (75 mL) was added at 15° C., stirred for 10 minutes, and the solid was filtered, washed with water and dried under vacuum to give 5-[[{(6R,7R)-7-[[{(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[{(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-2-[[{(4-methoxybenzyl)oxy]carbonyl]}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl}-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]furan-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (328 mg, crude) as yellow solid which was used in the next step without further purification.

Mass: ES⁺ 1084.68.

Step 9: (6R,7R)-7-[[{(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[{(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-3-(1-[4-(4-carbamimidoyl)furan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



To a solution of 5-[[{(6R,7R)-7-[[{(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[{(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino]-2-[[{(4-methoxybenzyl)oxy]carbonyl]}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl}-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]furan-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (from step 8, 328 mg, crude, ~0.32 mmol) in anhydrous methylene chloride (3 mL) was added anisole (0.8 mL) followed by trifluoroacetic acid (2.0 mL). The reaction mixture was stirred at room temperature for 3 h and the solvent was evaporated. To the residue was added diisopro-

132

pyl ether (30 mL) and the solid separated was filtered off, washed with diisopropyl ether and dried under vacuum to get greenish yellow color solid (315 mg), which was dissolved in water (30 mL) and filtered. The aqueous phase was lyophilized to give yellow solid (160 mg), which was purified by preparative HPLC to give (6R,7R)-7-[[{(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[{(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-3-(1-[4-(4-carbamimidoyl)furan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (28 mg, 12% over two steps).

¹H NMR (400 MHz, D₂O): δ 1.27 (s, 3H), 1.28 (s, 3H), 2.97 (d, 1H, J=18.0 Hz), 3.44 (d, 1H, J=18.0 Hz), 5.04-5.16 (m, 2H), 5.42 (d, 1H, J=14.5 Hz), 5.66 (d, 1H, J=4.7 Hz), 6.72 (s, 1H), 6.84 (s, 1H), 6.96 (d, 1H, J=3.1 Hz), 7.74 (d, 1H, J=3.5 Hz), 7.96 (d, 1H, J=7.0 Hz), 8.14 (s, 1H), 8.43 (d, 1H, J=7.0 Hz), 9.14 (s, 1H). Seven protons were not observed in D₂O.

Mass: ES⁺ 708.27.

Example 7 (Table 1, Compound 3)

(6R,7R)-7-[[{(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[{(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-8-oxo-3-[[1-(4-[N-[(3R)-piperidin-3-yl]carbamimidoyl]benzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate


¹H NMR (400 MHz, CD₃OD): δ 1.23 (s, 3H), 1.29 (s, 3H), 1.81-2.02 (m, 4H), 2.89 (d, 1H, J=18.0 Hz), 2.99 (m, 1H), 3.16 (m, 1H), 3.40-3.51 (m, 2H), 3.63 (d, 1H, J=17.6 Hz), 4.21 (m, 1H), 4.95 (m, 1H), 5.18 (d, 1H, J=4.8 Hz), 5.66-5.83 (m, 3H), 5.93 (d, 1H, J=14.0 Hz), 6.78 (s, 1H), 7.14 (d, 1H, J=3.2 Hz), 7.44 (d, 2H, J=8.0 Hz), 7.70 (d, 1H, J=6.8 Hz), 7.85 (d, 1H, J=8.4 Hz), 8.07 (s, 1H), 8.34 (s, 1H), 9.29 (d, 1H, J=6.0 Hz), 9.43 (s, 1H). Seven protons were not observed in CD₃OD.

Mass: ES⁻ 799.2.

133

Example 8 (Table 1, Compound 14)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo
 3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-
 azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.23 (s, 6H), 2.93 (d, 1H, J=18.0 Hz), 3.38 (d, 1H, J=17.6 Hz), 5.08-5.12 (m, 2H), 5.34 (d, 1H, J=14.4 Hz), 5.57 (s, 2H), 5.64 (d, 1H, J=4.4 Hz), 6.75 (s, 1H), 6.92 (d, 1H, J=3.6 Hz), 7.15 (t, 1H, J=7.6 Hz), 7.36 (d, 1H, J=7.2 Hz), 7.41 (d, 1H, J=10.4 Hz), 7.67 (d, 1H, J=3.6 Hz), 7.77 (d, 1H, J=6.8 Hz), 8.27 (d, 1H, J=8.0 Hz), 9.05 (s, 1H). Seven protons were not observed in D₂O.

Mass: ES⁺ 736.41

Example 9 (Table 1, Compound 17)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-[[1-({3-[(diaminomethylidene)amino]-1,2-oxazol-
 5-yl}methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]
 methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-
 2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.25 (s, 6H), 2.93 (d, 1H, J=17.6 Hz), 3.42 (d, 1H, J=18.0 Hz), 5.07-5.10 (m, 2H), 5.39 (d, 1H, J=13.6 Hz), 5.61 (s, 2H), 5.64 (d, 1H, J=4.8 Hz), 6.10 (s, 1H), 6.78 (s, 1H), 6.94 (d, 1H, J=3.2 Hz), 7.68 (d, 1H, J=3.6 Hz), 7.85 (d, 1H, J=6.4 Hz), 8.38 (d, 1H, J=6.8 Hz), 9.09 (s, 1H). Eight protons were not observed in D₂O.

Mass: ES⁺ 724.35


135

Example 10 (Table 1, Compound 18)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-
 imidazo[1,2-b]pyrazol-5-ium-5-yl}methyl)-8-oxo-5-
 thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



Step 1: 5-(1H-Imidazo[1,2-b]pyrazol-1-ylmethyl)
 thiophene-3-carbonitrile




To a stirred solution of 1H-imidazo[1,2-b]pyrazole (0.25 g, 2.33 mmol) in anhydrous dimethylformamide (5 mL) was added sodium hydride (60% dispersion in mineral oil, 0.14 g, 3.50 mmol) in small portions at 0° C. The reaction mixture was stirred at 0° C. for 30 min and a solution of 5-(bromomethyl)thiophene-3-carbonitrile (0.495 g, 2.45 mmol) in anhydrous dimethylformamide (1.0 mL) was added slowly. The reaction mixture was stirred at 0° C. for 1 h then at room temperature for 2 h. The reaction was quenched using aqueous saturated ammonium chloride solution (2 mL) and the resulting mixture was partitioned between ethyl acetate (80 mL) and water (10 mL). The organic phase was separated, washed with water (10 mL), brine (10 mL), dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude mixture was purified by flash column chromatography (silica gel, 230-400 mesh) using a 50 to 100% gradient of ethyl acetate in hexane to afford 5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophene-3-carbonitrile (0.28 g, 53%) as an off-white solid.

¹H NMR (400 MHz, DMSO-d₆): δ 5.38 (s, 2H), 5.72 (d, 1H, J=1.9 Hz), 7.29-7.32 (m, 1H), 7.46-7.48 (m, 1H), 7.57 (m, 2H), 8.49 (d, 1H, J=1.2 Hz)

Mass: ES⁺ 229.09

Step 2: Ethyl 5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)
 thiophene-3-carboximidate hydrochloride



A solution of 5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophene-3-carbonitrile (from step 1, 0.39 g, 1.71 mmol) in anhydrous ethanol (30 mL) at 0° C. was purged with a stream of anhydrous hydrogen chloride gas for 15 min. The reaction flask was stoppered and the reaction mixture was stirred at room temperature for 24 h. The volatiles were removed under reduced pressure and the residue was further


136

dried under high vacuum to afford ethyl 5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophene-3-carboximidate hydrochloride (0.60 g, crude) as a yellow solid which was used directly in the next step without further purification.

¹H NMR (400 MHz, DMSO-d₆): δ 1.45 (t, 3H, J=7.0 Hz), 4.55 (q, 2H, J=7.0 Hz), 5.46 (s, 2H), 5.83 (d, 1H, J=1.9 Hz), 7.40 (s, 1H), 7.58 (s, 1H), 7.65 (d, 1H, J=1.6 Hz), 7.81 (s, 1H), 8.78 (d, 1H, J=1.6 Hz). One proton was not observed in DMSO

Mass: ES⁺ 275.16

Step 3: 5-(1H-Imidazo[1,2-b]pyrazol-1-ylmethyl)
 thiophene-3-carboximidamide




A solution of ethyl 5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophene-3-carboximidate hydrochloride (from step 2, crude, 0.6 g, 1.71 mmol) in anhydrous methanol (20 mL) at 0° C. was purged with a stream of anhydrous ammonia gas for 15 min. The reaction flask was sealed and the reaction mixture was allowed to warm to room temperature and stirred for 17 h. The volatiles were removed under reduced pressure and the residue was further dried under high vacuum to afford 5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophene-3-carboximidamide (1.0 g, crude) as a reddish solid which was used in the next step without further purification.

¹H NMR (400 MHz, DMSO-d₆): δ 5.40 (s, 2H), 5.69 (d, 1H, J=1.9 Hz), 7.29-7.32 (m, 1H), 7.44-7.48 (m, 1H), 7.59 (d, 1H, J=2.7 Hz), 7.69 (d, 1H, J=0.8 Hz), 8.50 (d, 1H, J=1.6 Hz). Three protons were not observed in DMSO.

Mass: ES⁺ 246.09

137

Step 4: tert-Butyl {[5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophen-3-yl](imino)methyl}carbamate




To a vigorously stirred solution of 5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophene-3-carboximidamide (from step 3, crude, 1.0 g, 1.71 mmol) in 1,4-dioxane (10 mL) was added aqueous saturated sodium bicarbonate solution (6 mL) followed by di-tert-butyl dicarbonate (1.5 g, 6.84 mmol). The reaction mixture was stirred at room temperature for 17 h. The majority of 1,4-dioxane was removed under reduced pressure and the mixture was partitioned between dichloromethane (50 mL) and water (30 mL). The phases were separated and the aqueous layer was re-extracted with dichloromethane (60 mL). The combined organic extracts were dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude mixture was purified by flash column chromatography (silica gel, 230-400 mesh) using a 30 to 100% gradient of ethyl acetate in hexane to afford tert-butyl {[5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophen-3-yl](imino)methyl}carbamate (0.215 g, 36% over 3 steps) as a yellowish solid.

¹H NMR (400 MHz, DMSO-d₆): δ 1.41 (s, 9H), 5.32 (s, 2H), 5.65 (d, 1H, J=1.9 Hz), 7.24-7.30 (m, 1H), 7.39-7.46 (m, 1H), 7.54 (d, 1H, J=1.9 Hz), 7.60 (d, 1H, J=1.8 Hz), 8.21 (d, 1H, J=1.2 Hz), 8.85 (br s, 2H)

Mass: ES⁺346.23, ES⁻ 344.18

Step 5: 5-{[(6R,7R)-7-{[(2Z)-2-{2-[(tert-Butoxycarbonyl)amino]-1,3-thiazol-4-yl}-2-[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl}-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl)methyl)-1H-imidazo[1,2-b]pyrazol-5-ium iodide




138

To a solution of tert-butyl {[5-(1H-imidazo[1,2-b]pyrazol-1-ylmethyl)thiophen-3-yl](imino)methyl}carbamate (from step 4, 0.118 g, 0.34 mmol) in anhydrous dimethylformamide (2 mL) was added (6R,7R)-7-{[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl}-2-[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino}-3-(iodomethyl)-2-[(4-methoxybenzyl)oxy]carbonyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-5-olate (0.303 g, 0.34 mmol) in one portion at room temperature. The reaction mixture was stirred at room temperature for 18 h. Anhydrous dimethylformamide (1 mL) was added and the reaction mixture was cooled to -40° C. Potassium iodide (0.317 g, 2.39 mmol) followed by acetyl chloride (0.107 g, 1.37 mmol) was then added at -40° C. and the reaction mixture was stirred at 0° C. for 1 h. Aqueous 5% sodium bisulphite solution (containing sodium metabisulphite, 10 mL) was added and the resulting mixture was stirred for 15 min at room temperature. The precipitated solid was collected by filtration, washed with water (3×10 mL) and air-dried. The crude product was further dried under high vacuum to afford 5-{[(6R,7R)-7-{[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl}-2-[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino}acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl}-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl)methyl)-1H-imidazo[1,2-b]pyrazol-5-ium iodide (0.37 g, crude) as a yellow solid which was used in the next step without further purification.

Mass: ES⁺ 1090.75

139

Step 6: (6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



140

To a stirred solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl]-2-[[[(1-tert-butoxy-2-methyl-1-oxopropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl)methyl]-1-({4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium iodide (0.37 g, 0.30 mmol) and anisole (1 mL) in anhydrous dichloromethane (7 mL) was added trifluoroacetic acid (3 mL) dropwise at 0° C. The reaction mixture was allowed to warm to room temperature and stirred for 3 h. The volatiles were then removed under reduced pressure and the residue was triturated using diisopropyl ether (15 mL). The precipitate was collected by filtration, washed with diisopropyl ether (5 mL) and dried under high vacuum to afford a yellow solid (0.33 g). The crude material was then taken in distilled water (20 mL) and the resulting suspension was stirred vigorously for 30 min at room temperature. The insoluble material was


filtered off, the filtrate was collected and lyophilized. The crude product (0.16 g) was purified by prep HPLC to afford the trifluoroacetic acid salt of (6R,7R)-7-[[[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (15 mg; 6% over 2 steps).

¹H NMR (400 MHz, D₂O): δ 1.27 (s, 3H), 1.29 (s, 3H), 2.92 (d, 1H, J=18.0 Hz), 3.25 (d, 1H, J=18.0 Hz), 5.03 (d, 1H, J=5.1 Hz), 5.13 (s, 2H), 5.44 (s, 2H), 5.61 (d, 1H, J=4.7 Hz), 6.19 (d, 1H, J=3.5 Hz), 6.83 (s, 1H), 7.38-7.42 (m, 2H), 7.85-7.87 (m, 1H), 7.88 (d, 1H, J=4.3 Hz), 8.14 (s, 1H). Seven protons were not observed in D₂O.

Mass: ES⁺ 712.37


Example 11 (Table 1, Compound 59)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




141

Step 1: 5-{[(6R,7R)-7-{[(2Z)-2-{2-[(tert-Butoxycarbonyl)amino]-1,3-thiazol-4-yl}-2-(methoxyimino)acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl)methyl]-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium iodide



To a solution tert-butyl {[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophen-3-yl]carbamimidoyl} carbamate (0.12 g, 0.30 mmol) in dimethylformamide (2 mL) was added 4-methoxybenzyl (6R,7R)-7-{[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl}-2-(methoxyimino)acetyl] amino}-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0] oct-2-ene-2-carboxylate (0.19 g, 0.30 mmol) at 0° C. The reaction mixture was then degassed under reduced pressure for 0.5 h, followed by adding NaI (0.09 g, 0.6 mmol) and stirred at room temperature overnight. Sodium bisulphite (0.6 g) in sodium chloride solution (75 mL) was added at 15° C. and stirred for 10 minutes. The solid was filtered, washed with water and dried under vacuum to a yellow solid (0.29 g), which was used in the next step without further purification.

Step 2: (6R,7R)-7-{[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-({1-[4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



142

To a solution of 5-{[(6R,7R)-7-{[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-1,3-thiazol-4-yl}-2-(methoxyimino)acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl)methyl]-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl)methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (0.29 g) in dry dichloromethane (2.5 mL) at 0° C. was added


trifluoroacetic acid (2.5 mL) and anisole (0.82 mL, 7.75 mmol) and then stirred at room temperature for 2 h. Solvent was evaporated and diisopropyl ether (30 mL) was added. The solid separated was filtered, washed with di-isopropyl ether and dried under vacuum. The greenish yellow solid (315 mg) was dissolved in water (30 mL), filtered and the aqueous phase was lyophilized to give yellow solid (160 mg) which was purified by preparative HPLC to give (6R,7R)-7-{[(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-({1-[4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (0.007 g, 4%).

¹H NMR (400 MHz, D₂O): δ 3.05 (d, 1H, J=17.6 Hz), 3.45 (d, 1H, J=18.0 Hz), 3.84 (s, 3H), 5.11 (d, 1H, J=4.8 Hz), 5.21 (d, 1H, J=14.8 Hz), 5.35 (d, 1H, J=14.4 Hz), 5.70 (m, 3H), 6.84 (s, 1H), 6.98 (d, 1H, J=3.2 Hz), 7.34 (s, 1H), 7.73 (d, 1H, J=3.6 Hz), 7.88 (d, 1H, J=7.2 Hz), 8.12 (s, 1H), 8.31 (d, 1H, J=8.4 Hz), 9.08 (s, 1H). Six protons were not observed in D₂O.


Mass: ES⁺ 652.15

Example 12 (Table 1, Compound 62)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-8-oxo-3-[[1-[[4-(N-phenylcarbamimidoyl)thiophen-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



Step 1: N-phenyl-5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidamide




To a solution of ethyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride (0.5 g, 1.55 mmol) and triethylamine (0.56 mL, 4.0 mmol) in ethanol (5 mL) was added aniline (0.233 mmol) at room temperature. The reaction mixture was stirred at room

temperature for 2 days and concentrated. The residue was purified by column chromatography using methanol: dichloromethane (15:85) as eluent to give N-phenyl-5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidamide (0.28 g, 54%) as a beige solid.


¹H NMR (400 MHz, CD₃OD): δ 5.78 (s, 2H), 6.81 (d, 1H, J=4.4 Hz), 7.38-7.47 (m, 3H), 7.53-7.57 (m, 2H), 7.62 (m, 2H), 7.70 (d, 1H, J=6.4 Hz), 8.23 (d, 1H, J=6.4 Hz), 8.35 (s, 1H), 8.87 (s, 1H). Two protons were not observed in CD₃OD.

Step 2: 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-Butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[[4-(methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[[4-(N-phenylcarbamimidoyl)thiophen-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide



To a solution N-phenyl-5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidamide (0.10 g, 0.30 mmol) in dimethylformamide (2 mL) was added 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]}-2-(methoxyimino)acetyl]amino}-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.21 g, 0.30 mmol) at 0° C. The reaction mixture was then degassed under reduced pressure for 0.5 h, followed by adding NaI (0.09 g, 0.6 mmol) and stirred at room temperature overnight. Sodium bisulphite (0.6 g) in sodium chloride solution (75 mL) was added at 15° C. and stirred for 10 minutes. The solid was filtered off, washed with water and dried under vacuum to a yellow solid (0.28 g), which was used in the next step without further purification.

Step 3: (6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[(1-{[4-(N-phenylcarbamimidoyl)thiophen-2-yl]methyl}-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



To a solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]}-2-(methoxyimino)acetyl]amino}-2-[[[4-(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[[4-(N-phenylcarbamimidoyl)thiophen-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (0.28 g) in dry dichloromethane (2.5 mL) at 0° C. was added trifluoroacetic acid (2.5 mL) and anisole (0.82 mL, 7.75 mmol) and then stirred at room temperature for 2 h. Solvent was evaporated and diisopropyl ether (30 mL) was added. The solid was filtered, washed with diisopropyl ether and dried under vacuum. The greenish yellow solid (315 mg) was dissolved in water (30 mL), filtered and the aqueous phase was lyophilized to give yellow solid (160 mg) which was purified by preparative HPLC to give (6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[(1-{[4-(N-phenylcarbamimidoyl)thiophen-2-yl]methyl}-1H-pyrrolo[3,2-c]pyridin-5-ium-5-


yl)methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (0.01 g, 5%).

¹H NMR (400 MHz, D₂O): δ 3.05 (d, 1H, J=18.0 Hz), 3.43 (d, 1H, J=18.0 Hz), 3.84 (s, 3H), 5.10 (d, 1H, J=4.8 Hz), 5.22 (d, 1H, J=15.2 Hz), 5.35 (d, 1H, J=14.4 Hz), 5.71 (d, 1H, J=4.8 Hz), 5.76 (s, 2H), 7.00 (d, 1H, J=3.2 Hz), 7.30 (d, 2H, J=7.2 Hz), 7.41-7.49 (m, 4H), 7.78 (d, 1H, J=3.2 Hz), 7.92 (d, 1H, J=7.6 Hz), 8.21 (s, 1H), 8.33 (d, 1H, J=8.0 Hz), 9.09 (s, 1H). Five protons were not observed in D₂O.

Mass: ES⁺ 662.09


Example 13 (Table 1, Compound 63)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[(1-{[4-(pyrrolidin-1-ylcarbamimidoyl)thiophen-2-yl]methyl}-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



147


Step 1: 1-(Pyrrolidin-1-yl)-1-[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophen-3-yl]methanimine



To a solution of ethyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride (0.5 g, 1.55 mmol) and triethylamine (0.56 mL, 4.0 mmol) in ethanol (5 mL) was added pyrrolidine (0.2 g, 2.81 mmol) at room temperature. The reaction mixture was stirred at room temperature for 2 days and concentrated. The residue was purified by column chromatography using methanol: dichloromethane (15:85) as eluent to give 1-(pyrrolidin-1-yl)-1-[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophen-3-yl]methanimine (0.26 g, 54%) as a beige solid.

¹H NMR (400 MHz, CD₃OD): δ 1.96 (m, 2H), 2.14 (m, 2H), 3.30 (m, 4H), 5.64 (s, 2H), 6.74 (d, 1H, J=4.0 Hz), 7.38 (s, 1H), 7.51 (d, 1H, J=3.6 Hz), 7.57 (d, 1H, J=6.0 Hz), 7.99 (s, 1H), 8.18 (d, 1H, J=5.6 Hz), 8.81 (s, 1H). One proton was not observed in CD₃OD.


Step 2: 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]}-2-(methoxyimino)acetyl]amino}-2-[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[4-(pyrrolidin-1-ylcarboximidoyl)thiophen-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide



To a solution 1-(pyrrolidin-1-yl)-1-[5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophen-3-yl]methanimine (0.093 g, 0.30 mmol) in dimethylformamide (2 mL) was added 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]}-2-(methoxyimino)acetyl]amino]-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.21 g, 0.30 mmol) at 0° C. The reaction mixture was then degassed under reduced pressure for 0.5 h, followed by adding NaI (0.09 g, 0.6 mmol) and stirred at room temperature overnight. Sodium bisulphite (0.6 g) in sodium chloride solution (75 mL) was added at 15° C. and stirred for 10 minutes. The solid separated was filtered, washed with water and dried under vacuum to a yellow solid (0.28 g), which was used in the next step without further purification.

149

Step 3: (6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[(1-{[4-(pyrrolidin-1-ylcarbamimidoyl)thiophen-2-yl]methyl}-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



To a solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxy-carbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[[4-(methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[[4-(pyrrolidin-1-ylcarbamimidoyl)thiophen-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (0.29 g) in dry dichloromethane (2.5 mL) at 0° C. was added trifluoroacetic acid (2.5 mL) and anisole (0.82 mL, 7.75


H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (0.011 g, 5.7%).

¹H NMR (200 MHz, D₂O): δ 1.79 (m, 2H), 1.97 (m, 2H), 3.03 (d, 1H, J=18.0 Hz), 3.42 (m, 5H), 3.84 (s, 3H), 5.10 (d, 1H, J=4.4 Hz), 5.21 (d, 1H, J=14.8 Hz), 5.35 (d, 1H, J=14.4 Hz), 5.70 (m, 3H), 6.98 (d, 1H, J=3.2 Hz), 7.22 (s, 1H), 7.74 (d, 1H, J=3.2 Hz), 7.85 (s, 1H), 7.89 (d, 1H, J=7.2 Hz), 8.32 (d, 1H, 7.2 Hz), 9.08 (s, 1H). Four protons were not observed in D₂O.

Mass: ES⁺ 740.13


Example 14 (Table 1, Compound 64)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[[1-({4-[N-(1,3-thiazol-2-yl)carbamimidoyl]thiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



151


Step 1: 5-(1H-Pyrrolo[3,2-c]pyridin-1-ylmethyl)-N-(1,3-thiazol-2-yl)thiophene-3-carboximidamide



To a solution of ethyl 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride (0.5 g, 1.55 mmol) and triethylamine (0.56 mL, 4.0 mmol) in ethanol (5 mL) was added 1,3-thiazol-2-amine (0.25 g, 2.33 mmol) at room temperature. The reaction mixture was stirred at room temperature for 2 days and concentrated. The residue was purified by column chromatography using methanol: dichloromethane (15:85) as eluent to give 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)-N-(1,3-thiazol-2-yl)thiophene-3-carboximidamide (0.22 g, 42%) as a beige solid.

$^1\text{H NMR}$ (400 MHz, CD_3OD): δ 5.43 (s, 2H), 6.65 (d, 1H, $J=4.0$ Hz), 7.16 (m, 2H), 7.27 (m, 2H), 7.44 (m, 1H, $J=6.0$ Hz), 7.64 (s, 1H), 8.32 (d, 1H, $J=6.0$ Hz), 8.92 (s, 1H).


Step 2: 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-(4-[N-(1,3-thiazol-2-yl)carbamimidoyl]thiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide



To a solution of 5-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)-N-(1,3-thiazol-2-yl)thiophene-3-carboximidamide (0.102 g, 0.30 mmol) in dimethylformamide (2 mL) was added 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.21 g, 0.30 mmol) at 0°C . The reaction mixture was then degassed under reduced pressure for 0.5 h, followed by adding NaI (0.09 g, 0.6 mmol) and stirred at room temperature overnight. Sodium bisulphite (0.6 g) in sodium chloride solution (75 mL) was added at 0°C ., stirred for 10 minutes and the solid separated was filtered, washed with water and dried under vacuum to a yellow solid (0.3 g), which was used in the next step without further purification.

153

Step 3: (6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[[1-({4-[N-(1,3-thiazol-2-yl)carbamimidoyl]thiophen-2-yl}methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



To a solution of 5-[[[(6R,7R)-7-[[[(2Z)-2-[(tert-butoxy-35 carbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino]-2-[[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-({4-[N-(1,3-thiazol-2-yl)carbamimidoyl]thiophen-2-yl}methyl)-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (0.3 g) in dry dichloromethane (2.5 mL) at 0° C. was added trifluoroacetic acid (2.5 mL) and anisole (0.82 mL, 7.75 mmol) and then stirred at room temperature for 2 h, then the solvent was evaporated and diisopropyl ether (30 mL) was added. The solid was filtered, washed with diisopropyl ether and dried under vacuum. The greenish yellow solid (315 mg) was dissolved in water (30 mL), filtered and the aqueous phase was lyophilized to give yellow solid (160 mg) which was purified by preparative HPLC to give (6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-40 (methoxyimino)acetyl]amino]-8-oxo-3-[[1-({4-[N-(1,3-thiazol-2-yl)carbamimidoyl]thiophen-2-yl}methyl)-1H-pyr-


rolo[3,2-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (0.005 g, 2.5%).

¹H NMR (400 MHz, D₂O): δ 3.03 (d, 1H, J=17.6 Hz), 3.43 (d, 1H, J=17.2 Hz), 3.84 (s, 3H), 5.11 (d, 1H, J=4.8 Hz), 5.21 (d, 1H, J=13.6 Hz), 5.36 (d, 1H, J=14.0 Hz), 5.72 (m, 3H), 7.01 (d, 1H, J=4.2 Hz), 7.25 (d, 1H, J=3.6 Hz), 7.41 (d, 1H, J=7.6 Hz), 7.49 (m, 1H), 7.75 (m, 1H), 7.92 (d, 1H, J=7.2 Hz), 8.20 (s, 1H), 8.33 (d, 1H, J=7.6 Hz), 9.06 (s, 1H). Five protons were not observed in D₂O.

Mass: ES⁺ 769.03


Example 15 (Table 1, Compound 60)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[(4-carbamimidoyl]thiophen-2-yl)methyl]-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl}methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



155


Step 1: 5-(1H-Pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carbonitrile



To a solution of 1H-pyrrolo[3,2-b]pyridine (0.59 g, 5.00 mmol) in anhydrous dimethylformamide (10 mL) was added sodium hydride (60% suspension in mineral oil, 0.29 g, 7.25 mmol) at 0° C. in small portions under nitrogen. After the addition, the mixture was stirred for 10 minutes at 0° C. and then at room temperature for 10 minutes, cooled to 0° C. and was added 5-(bromomethyl)thiophene-3-carbonitrile (1.11 g, 5.50 mmol) in anhydrous tetrahydrofuran (10 mL) dropwise. After the addition, the reaction mixture was stirred at 0° C. for 1.5 h, quenched with ice-water (10 mL) and extracted with ethyl acetate (20 mL×2). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and evaporated. The residue was purified by column chromatography using DCM: MeOH (92:8) as eluent to afford 5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carbonitrile (1.0 g, 80%) as a brown solid.

¹H NMR (400 MHz, CDCl₃): δ 5.47 (s, 2H), 6.79 (d, 1H, J=4.0 Hz), 7.10 (s, 1H), 7.12 (m, 1H), 7.36 (d, 1H, J=3.6 Hz), 7.58 (d, 1H, J=10.4 Hz), 7.82 (s, 1H), 8.50 (d, 1H, J=6.0 Hz).


Step 2: Methyl 5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride



Anhydrous hydrogen chloride gas was bubbled through a solution of 5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carbonitrile (1.0 g, 4.18 mmol) in anhydrous ethanol (30 mL) at 0° C. for 20 minutes. The reaction mixture was stirred at room temperature for 20 h and the solvent was evaporated and dried under vacuum to give methyl 5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride (0.97 g crude) as brown solid, which was used in next step without further purification.


156

Step 3: 5-(1H-Pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carboximidamide



Anhydrous ammonia gas was bubbled through a solution of methyl 5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride (0.97 g crude, ~4.1 mmol) in anhydrous methanol (40 mL) at 0° C. for 15 minutes. Then the reaction mixture was stirred at room temperature for 16 h. Solvent was evaporated and dried under vacuum to give 5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carboximidamide (1.2 g, crude) as a brown solid which was used in the next step without further purification.

Step 4: tert-Butyl {[5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophen-3-yl]carboximidoyl}carbamate




To a solution of 5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophene-3-carboximidamide (0.97 g, crude) in 1,4-dioxane (100 mL) was added aqueous saturated sodium carbonate solution (50 mL) followed by di-tert-butyl dicarbonate (3.30 g, 15.12 mmol) at 0° C. The reaction mixture was stirred at room temperature for 16 h, the solvent was evaporated and was added water (50 mL) was added then extracted with ethyl acetate (2×50 mL). The organic phases were combined, washed with brine (2×50 mL) and dried over Na₂SO₄. The solvent was evaporated and the crude compound was purified by flash chromatography using RediSep silica 40 g flash column (0-6% methanol in methylene chloride as eluent) to give tert-butyl {[5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophen-3-yl]carboximidoyl}carbamate (1.2 g, 61% over three steps) as an off-white solid.

¹H NMR (400 MHz, CDCl₃): δ 1.51 (s, 9H), 5.39 (s, 2H), 6.66 (d, 1H, J=4.00 Hz), 7.06 (m, 1H), 7.32 (d, 1H, J=3.2 Hz), 7.42 (s, 1H), 7.58 (d, 1H, J=10.4 Hz), 7.87 (s, 1H), 8.42 (d, 1H, J=5.6 Hz).


157

Step 5: 4-{[(6R,7R)-7-{[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl}-2-(methoxyimino)acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl}-1-({4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl}methyl)-1H-pyrrolo[3,2-b]pyridin-4-ium iodide



To a solution tert-butyl {[5-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)thiophen-3-yl]carboimidoyl} carbamate (0.12 g, 0.30 mmol) in dimethylformamide (2 mL) was added 4-methoxybenzyl (6R,7R)-7-{[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl}-2-(methoxyimino)acetyl]amino}-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.21 g, 0.30 mmol) at 0° C. The reaction mixture was then degassed under reduced pressure for 0.5 h, followed by adding NaI (0.09 g, 0.6 mmol) and stirred at room temperature overnight. Sodium bisulphite (0.6 g) in sodium chloride solution (75 mL) was added at 15° C. and stirred for 10 minutes. The solid was filtered, washed with water and dried under vacuum to a yellow solid (0.3 g), which was used in the next step without further purification.

Step 6: (6R,7R)-7-{[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl}methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



158

To a solution of 4-{[(6R,7R)-7-{[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl}-2-(methoxyimino)acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl}-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl}methyl}-1-({4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl}methyl)-1H-pyrrolo[3,2-b]pyridin-4-ium iodide (0.3 mg, crude) in dry dichloromethane (2.5 mL) at

0° C. was added trifluoroacetic acid (2.5 mL) and anisole (0.82 mL, 7.75 mmol) and then stirred at room temperature for 2 h. Solvent was evaporated, diisopropyl ether (30 mL) was added, and the solid separated was filtered, washed with diisopropyl ether and dried under vacuum. The greenish yellow solid (315 mg) was dissolved in water (30 mL), filtered and the aqueous phase was lyophilized to give yellow solid (165 mg), which was purified by preparative HPLC to give (6R,7R)-7-{[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl}methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (0.007 g, 3.9%).


¹H NMR (400 MHz, D₂O): δ 3.08 (d, 1H, J=18.0 Hz), 3.23 (d, 1H, J=17.6 Hz), 3.86 (s, 3H), 5.04 (d, 1H, J=4.8 Hz), 5.39 (d, 1H, J=15.2 Hz), 5.67 (d, 1H, J=14.8 Hz), 5.71 (d, 1H, J=4.4 Hz), 5.76 (s, 2H), 6.93 (d, 1H, J=3.6 Hz), 7.35 (s, 1H), 7.55 (m, 1H), 8.09 (d, 1H, J=3.2 Hz), 8.13 (s, 1H), 8.49 (m, 2H). Six protons were not observed in D₂O.

Mass: ES⁺ 686.14


159

Example 16 (Table 1, Compound 61)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({1-[4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl}methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




Step 1: 5-(1H-Pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-carbonitrile



To a solution of 1H-pyrrolo[2,3-c]pyridine (0.59 g, 5.00 mmol) in anhydrous dimethylformamide (10 mL) was added sodium hydride (60% suspension in mineral oil, 0.29 g, 7.25 mmol) at 0° C. in small portions under nitrogen. After the addition, reaction mixture was stirred for 10 minutes at 0° C., then at room temperature for 10 minutes, cooled to 0° C. then 5-(bromomethyl)thiophene-3-carbonitrile (1.11 g, 5.50 mmol) in anhydrous tetrahydrofuran (10 mL) was added dropwise. After the addition, reaction mixture was stirred at 0° C. for 1.5 h, quenched with ice-water (10 mL) and extracted with ethyl acetate (20 mLx2). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and evaporated. The residue was purified by column chromatography using DCM: MeOH (92:8) as eluent to afford 5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-carbonitrile (0.88 g, 73%) as a brown solid. ¹H NMR (400 MHz, CDCl₃): δ 5.56 (s, 2H), 6.59 (d, 1H, J=2.8 Hz), 7.13 (s, 1H), 7.27 (d, 1H, J=6.8 Hz), 7.54 (d, 1H, J=5.2 Hz), 7.82 (s, 1H), 8.29 (d, 1H, J=5.2 Hz), 8.73 (s, 1H).

Step 2: Methyl 5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride




160

Anhydrous hydrogen chloride gas was bubbled through a solution of 5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-carbonitrile (0.8 g, 3.34 mmol) in anhydrous ethanol (30 mL) at 0° C. for 20 minutes. Then the reaction mixture was stirred at room temperature for 20 h. Solvent was evaporated and dried under vacuum to give methyl 5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-car-


boximidate hydrochloride (1.0 g, crude) as brown solid, which was used in next step without further purification.

Step 3: 5-(1H-Pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-carbonitrile



Anhydrous ammonia gas was bubbled through a solution of methyl 5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-carboximidate hydrochloride (1.0 g crude, ~4.1 mmol) in anhydrous methanol (40 mL) at 0° C. for 15 minutes. The reaction mixture was stirred at room temperature for 16 h. Solvent was evaporated and dried under vacuum to give 5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-carboximidamide (0.73 g, crude) as a brown solid, which was used in the next step without further purification.


Step 4: tert-Butyl {[5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophen-3-yl]carbamimidoyl}carbamate



To a solution of 5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophene-3-carboximidamide (0.73 g crude) in 1,4-dioxane (100 mL) was added aqueous saturated sodium carbonate solution (50 mL) followed by di-tert-butylidicarbonate (2.9 g, 13.12 mmol) at 0° C. and stirred at room temperature for 16 h. The solvent was evaporated and was added water (50 mL) then extracted with ethyl acetate (2×50 mL). The organic phases were combined, washed with brine (2×50 mL), dried over Na₂SO₄, concentrated and the crude compound was purified by flash chromatography using RediSep silica 40 g flash column (0-6% methanol in methylene chloride as eluent) to give tert-butyl {[5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophen-3-yl]carboximidoyl}carbamate (1.0 g, 51% over three steps) as an off-white solid.

¹H NMR (400 MHz, CDCl₃): δ 1.53 (s, 9H), 5.50 (s, 2H), 6.54 (d, 1H, J=3.2 Hz), 6.88 (d, 1H, J=3.6 Hz), 7.26 (s, 1H), 7.37 (d, 1H, J=3.6 Hz), 7.43 (s, 1H), 7.52 (d, 1H, J=4.8 Hz), 7.84 (s, 1H), 8.25 (d, 1H, J=5.6 Hz), 8.78 (s, 1H).

Step 5: 6-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-(4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl)methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium iodide




To a solution tert-butyl {[5-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)thiophen-3-yl]carboximidoyl}carbamate (0.12 g, 0.30 mmol) in dimethylformamide (2 mL) was added 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.21 g, 0.30 mmol) at 0° C. The reaction mixture was then degassed under reduced pressure for 0.5 h, treated with NaI (0.09 g, 0.6 mmol) and stirred at room temperature overnight.

Sodium bisulphite (0.6 g) in sodium chloride solution (75 mL) was added to the reaction mixture at 15° C., stirred for

10 minutes and the solid separated was filtered, washed with water then dried under vacuum to a yellow solid (0.3 g), which was used in the next step without further purification.

Step 6: (6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate formic acid salt



163

To a solution of 6-[[[(6R,7R)-7-[[[(2Z)-2-[[[(tert-butoxy-carbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[[4-[N-(tert-butoxycarbonyl)carbamimidoyl]thiophen-2-yl]methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium iodide (0.3 mg, crude)] in dry dichloromethane (2.5 mL) at 0° C. was added trifluoroacetic acid (2.5 mL) and anisole (0.82 mL, 7.75 mmol) then stirred at room temperature for 2 h. Solvent was evaporated, diisopropyl ether (30 mL) was added and solid was filtered, washed with diisopropyl ether then dried under vacuum. The greenish yellow solid (315 mg) was dissolved in water (30 mL), filtered and the aqueous phase was lyophilized to give yellow solid, which was purified by preparative HPLC to get (6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-[[4-carbamimidoyl]thiophen-2-yl]methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-

164


oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate yellow solid (165 mg) as formic acid salt (0.02 g, 11.2%).

¹H NMR (400 MHz, D₂O): δ 2.99 (d, 1H, J=18.0 Hz), 3.38 (d, 1H, J=18.0 Hz), 3.85 (s, 3H), 5.06 (d, 1H, J=5.2 Hz), 5.13 (d, 1H, J=14.4 Hz), 5.34 (d, 1H, J=14.0 Hz), 5.69-5.82 (m, 3H), 6.88 (d, 1H, J=3.2 Hz), 7.49 (s, 1H), 7.96 (d, 1H, J=6.8 Hz), 8.14 (d, 1H, J=6.8 Hz), 8.32 (s, 2H), 9.06 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 686.23

Example 17 (Table 1, Compound 28)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-[[4-carbamimidoyl]thiophen-2-yl]methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O) δ 1.24 (s, 3H), 1.27 (s, 3H), 2.88 (d, 1H, J=18.0 Hz), 3.24 (d, 1H, J=18.0 Hz), 5.02 (d, 1H, J=4.7 Hz), 5.13 (m, 2H), 5.45 (s, 2H), 5.61 (d, 1H, J=4.7 Hz), 6.20 (d, 1H, J=3.9 Hz), 7.40 (s, 2H), 7.87-7.89 (m, 2H), 8.14 (s, 1H). Seven protons were not observed in D₂O.

Mass: ES⁺ 747.34 and 749.22

Example 18 (Table 1, Compound 4)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-[[4-carbamimidamido]benzyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




165

¹H NMR (400 MHz, D₂O): δ 1.27 (s, 3H), 1.28 (s, 3H), 2.92 (d, 1H, J=17.6 Hz), 3.38 (d, 1H, J=17.6 Hz), 5.08-5.12 (m, 2H), 5.33 (d, 1H, J=14.9 Hz), 5.44 (s, 2H), 5.64 (s, 1H), 6.80 (s, 1H), 6.91 (s, 1H), 7.11-7.17 (m, 4H), 7.67-7.71 (m, 2H), 8.16 (s, 0.4H, formic acid), 8.25 (m, 1H), 9.04 (s, 1H). 5
Eight protons were not observed in D₂O.

Mass: ES⁺ 733.32

Example 19 (Table 1, Compound 7)

(6R,7R)-3-[(1-{4-[N-(2-Aminoethyl)carbamimidoyl] 10
benzyl}-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)
methyl]-7-[[2Z]-2-(2-amino-1,3-thiazol-4-yl)-2-
{[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-
8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-car-
boxylate




¹H NMR (400 MHz, D₂O): δ 1.20 (s, 3H), 1.21 (s, 3H), 30
2.92 (d, 1H, J=18.0 Hz), 3.21 (t, 2H, J=6.3 Hz), 3.38 (d, 1H,
J=18.0 Hz), 3.63 (t, 2H, J=6.3 Hz), 5.06-5.11 (m, 2H), 5.32
(d, 1H, J=14.4 Hz), 5.52 (s, 2H), 5.59 (d, 1H, J=4.7 Hz), 6.73
(s, 1H), 6.92 (d, 1H, J=3.1 Hz), 7.21 (d, 2H, J=8.2 Hz), 7.52
(d, 2H, J=8.2 Hz), 7.66-7.69 (m, 2H), 8.23 (d, 1H, J=7.0 Hz), 35
9.05 (s, 1H). Eight protons were not observed in D₂O.

Mass: ES⁺ 761.13

Example 20 (Table 1, Compound 8)

(6R,7R)-7-[[2Z]-2-(2-Amino-1,3-thiazol-4-yl)-2- 40
{[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-
3-({1-[4-(4,5-dihydro-1H-imidazol-2-yl)benzyl]-1H-
pyrrolo[3,2-c]pyridin-5-ium-5-yl}methyl)-8-oxo-5-
thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.24 (s, 3H), 1.27 (s, 3H), 60
2.94 (d, 1H, J=17.5 Hz), 3.40 (d, 1H, J=17.5 Hz), 3.89 (s,
4H), 5.09-5.13 (m, 2H), 5.34 (d, 1H, J=14.8 Hz), 5.55 (s,
2H), 5.63 (d, 1H, J=4.7 Hz), 6.78 (s, 1H), 6.96 (s, 1H), 7.23
(d, 2H, J=7.1 Hz), 7.58 (d, 2H, J=7.1 Hz), 7.69-7.71 (m, 2H),
8.19 (s, 1H, formic acid), 8.25 (d, 1H, J=7.1 Hz), 9.08 (s, 65
1H). Five protons were not observed in D₂O.

Mass: ES⁺ 744.27

Example 21 (Table 1, Compound 12)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-({1-[2-(4-carbamimidamidophenyl)ethyl]-1H-
 pyrrolo[3,2-c]pyridin-5-ium-5-yl}methyl)-8-oxo-5-
 thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.30 (s, 6H), 2.80 (d, 1H, J=17.6 Hz), 3.00 (m, 2H), 3.23 (d, 1H, J=17.6 Hz), 4.44 (m, 2H), 5.03-5.07 (m, 2H), 5.21 (d, 1H, J=14.1 Hz), 5.56 (d, 1H, J=4.3 Hz), 6.75-6.86 (m, 6H), 7.02 (d, 1H, J=6.7 Hz), 7.60 (s, 1H), 7.85 (d, 1H, J=6.7 Hz), 8.19 (s, 0.5H, formic acid), 8.90 (s, 1H). Eight protons were not observed in D₂O.

Mass: ES⁺ 747.27

Example 22 (Table 1, Compound 16)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-({1-[4-(4-carbamimidoyl)-1,3-thiazol-2-yl]methyl]-
 1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl}methyl)-8-oxo-
 5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.29 (s, 3H), 1.32 (s, 3H), 2.98 (d, 1H, J=17.6 Hz), 3.40 (d, 1H, J=17.6 Hz), 5.09 (d, 1H, J=4.7 Hz), 5.14 (d, 1H, J=14.4 Hz), 5.34 (d, 1H, J=14.4 Hz), 5.67 (d, 1H, J=4.7 Hz), 5.85 (s, 2H), 6.91 (s, 1H), 6.97 (d, 1H, J=3.1 Hz), 7.74 (d, 1H, J=3.1 Hz), 7.86 (d, 1H, J=7.1 Hz), 8.09 (s, 0.4H, formic acid), 8.31 (d, 1H, J=7.1 Hz), 8.39 (s, 1H), 9.08 (s, 1H). Seven protons were not observed in D₂O.

Mass: ES⁺ 725.37

169

Example 23 (Table 1, Compound 19)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-({1-[4-(N'-methoxycarbamimidoyl)benzyl]-1H-
 pyrrolo[3,2-c]pyridin-5-ium-5-yl}methyl)-8-oxo-5-
 thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.25 (s, 3H), 1.26 (s, 3H),
 2.94 (d, 1H, J=18.0 Hz), 3.35 (d, 1H, J=18.0 Hz), 3.64 (s,
 3H), 5.07 (d, 1H, J=4.7 Hz), 5.09 (d, 1H, J=14.5 Hz), 5.30
 (d, 1H, J=14.5 Hz), 5.45 (s, 2H), 5.65 (d, 1H, J=4.7 Hz), 6.77
 (s, 1H), 6.91 (d, 1H, J=3.5 Hz), 7.11 (d, 2H, J=8.2 Hz), 7.38
 (d, 2H, J=8.2 Hz), 7.64 (d, 1H, J=3.5 Hz), 7.69 (d, 1H, J=6.7
 Hz), 8.19 (d, 1H, J=6.7 Hz), 8.26 (s, 3H, formic acid), 9.02
 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 748.45

Example 24 (Table 1, Compound 30)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-({1-[4-(4-carbamimidoyl-1,3-oxazol-2-yl)methyl]-
 1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl}methyl)-8-oxo-
 5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.30 (s, 3H), 1.32 (s, 3H),
 2.98 (d, 1H, J=18.0 Hz), 3.39 (d, 1H, J=18.0 Hz), 5.08 (d,
 1H, J=4.7 Hz), 5.14 (d, 1H, J=14.9 Hz), 5.33 (d, 1H, J=14.9
 Hz), 5.66 (d, 1H, J=4.7 Hz), 5.71 (s, 2H), 6.92 (s, 1H), 6.95
 (d, 1H, J=3.5 Hz), 7.69 (d, 1H, J=3.5 Hz), 7.85 (d, 1H, J=7.0
 Hz), 8.31 (d, 1H, J=7.0 Hz), 8.50 (s, 1H), 9.06 (s, 1H). Seven
 protons were not observed in D₂O.

Mass: ES⁺ 709.29

171

Example 25 (Table 1, Compound 31)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-
 3-({1-[(4-{N-[(Z)-(dimethylamino)methylidene]
 carbamimidoyl}-1,3-oxazol-2-yl)methyl]-1H-pyrrolo
 [3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-
 azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.29 (s, 3H), 1.3 (s, 3H),
 2.98 (d, 1H, J=18.0 Hz), 3.05 (s, 3H), 3.13 (s, 3H), 3.40 (d,
 1H, J=18.0 Hz), 5.08 (d, 1H, J=4.7 Hz), 5.13 (d, 1H, J=14.9
 Hz), 5.34 (d, 1H, J=14.9 Hz), 5.65 (d, 1H, J=4.7 Hz), 5.71
 (s, 2H), 6.87 (s, 1H), 6.95 (d, 1H, J=3.2 Hz), 7.70 (d, 1H,
 J=3.6 Hz), 7.86 (d, 1H, J=7.4 Hz), 8.23 (s, 1H), 8.32 (d, 1H,
 J=6.7 Hz), 8.45 (s, 1H), 9.08 (s, 1H). Five protons were not
 observed in D₂O.

Mass: ES⁺ 764.36

Example 26 (Table 1, Compound 29)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-
 3-[(1-[[3-(4,5-dihydro-1H-imidazol-2-yl)-1,2-ox-
 azol-5-yl]methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-
 5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-
 ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.25 (s, 6H), 2.94 (d, 1H,
 J=17.2 Hz), 3.40 (d, 1H, J=17.2 Hz), 4.59 (s, 4H), 5.08-5.13
 (m, 2H), 5.35 (d, 1H, J=14.4 Hz), 5.63 (d, 1H, J=5.1 Hz),
 5.74 (s, 2H), 6.72 (m, 2H), 6.94 (d, 1H, J=1.2 Hz), 7.68 (d,
 1H, J=3.1 Hz), 7.86 (d, 1H, J=7.1 Hz), 8.23 (s, 1H, formic
 acid), 8.34 (d, 1H, J=7.1 Hz), 9.07 (s, 1H). Five protons were
 not observed in D₂O.

Mass: ES⁺ 735.21

Example 27 (Table 1, Compound 15)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[2-carboxypropan-2-yl]oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

5



¹H NMR (400 MHz, D₂O): δ 1.29 (s, 3H), 1.31 (s, 3H), 2.95 (d, 1H, J=17.6 Hz), 3.24 (d, 1H, J=17.6 Hz), 5.04 (d, 1H, J=4.8 Hz), 5.14 (s, 2H), 5.37 (s, 2H), 5.62 (d, 1H, J=4.8 Hz), 6.23 (d, 1H, J=3.2 Hz), 6.92 (s, 1H), 7.40-7.48 (m, 4H), 7.84 (d, 1H, J=2.0 Hz), 7.88 (d, 1H, J=3.6 Hz). Seven protons were not observed in D₂O.


30

Mass: ES⁺ 725.42

Example 28 (Table 1, Compound 6)

35

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-[[[2-carboxypropan-2-yl]oxy]imino]acetyl]amino]-3-({1-[4-(2-carbamimidamidoethoxy)benzyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl}methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



¹H NMR (400 MHz, DMSO-d₆): δ 1.38 (s, 6H), 2.91 (d, 1H, J=18.4 Hz), 3.20-3.50 (m, 3H), 3.92 (s, 1H), 4.02 (s, 1H), 4.85 (d, 1H, J=13.2 Hz), 5.04 (d, 1H, J=5.2 Hz), 5.46 (s, 2H), 5.68 (d, 1H, J=13.2 Hz), 5.79 (d, 1H, J=4.8 Hz), 6.72 (s, 1H), 6.81 (d, 2H, J=8.8 Hz), 7.03 (d, 1H, J=3.2 Hz), 7.17 (s, 1H), 7.30 (d, 2H, J=8.0 Hz), 8.05 (d, 1H, J=6.0 Hz), 8.14 (s, 1H), 9.30 (br s, 1H), 9.44 (s, 1H). Eight protons were not observed in D₂O.


60

65

Mass: ES⁺ 777.25

Example 29 (Table 1, Compound 20)

(6R,7R)-7-[[[(2Z)-2-(5-Amino-1,2,4-thiadiazol-3-yl)-
2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]
amino]-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-
1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-
5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.25 (s, 6H), 2.93 (d, 1H, J=17.6 Hz), 3.37 (d, 1H, J=18.0 Hz), 5.08-5.12 (m, 2H), 5.33 (d, 1H, J=14.4 Hz), 5.57 (s, 2H), 5.65 (d, 1H, J=4.4 Hz), 6.92 (d, 1H, J=3.6 Hz), 7.14 (t, 1H, J=8.0 Hz), 7.36 (d, 1H, J=8.4 Hz), 7.41 (d, 1H, J=10.0 Hz), 7.66 (d, 1H, J=3.2 Hz), 7.77 (d, 1H, J=6.8 Hz), 8.26 (d, 1H, J=6.8 Hz), 9.05 (s, 1H). Seven protons were not observed in D₂O.

Mass: ES⁺ 737.36

Example 30 (Table 1, Compound 21)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



¹H NMR (400 MHz, D₂O): δ 1.22 (s, 6H), 2.92 (d, 1H, J=18.0 Hz), 3.37 (d, 1H, J=19.2 Hz), 5.07-5.11 (m, 2H), 5.34 (d, 1H, J=14.4 Hz), 5.58 (s, 2H), 5.64 (d, 1H, J=4.4 Hz), 6.93 (d, 1H, J=4.0 Hz), 7.17 (t, 1H, J=7.6 Hz), 7.36 (d, 1H, J=8.8 Hz), 7.42 (d, 1H, J=10.0 Hz), 7.67 (d, 1H, J=3.6 Hz), 7.78 (d, 1H, J=6.4 Hz), 8.25 (d, 1H, J=6.8 Hz), 9.05 (s, 1H). Seven protons were not observed in D₂O.


Mass: ES⁺ 770.36

177

Example 31 (Table 1, Compound 26)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-
 3-({1-[(4-carbamimidoyl-1,3-thiazol-2-yl)methyl]-
 1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl)-8-
 oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-
 carboxylate

5



25

¹H NMR (400 MHz, D₂O): δ 1.26 (s, 3H), 1.28 (s, 3H),
 2.91 (d, 1H, J=18 Hz), 3.28 (d, 1H, J=18 Hz), 5.08 (s, 1H),
 5.15 (s, 1H), 5.18 (s, 1H), 5.60-5.70 (m, 3H), 6.32 (s, 1H),
 6.79 (s, 1H), 7.48 (s, 1H), 7.92-8.00 (m, 2H), 8.44 (s, 1H).
 65 Seven protons were not observed in D₂O.


30

Mass: ES⁺ 714.27

Example 32 (Table 1, Compound 22)

(6R,7R)-7-({[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 [(carboxymethoxy)imino]acetyl]amino)-3-{{1-[(4-
 carbamimidoyl-2-fluorobenzyl)-1H-imidazo[1,2-b]
 pyrazol-5-ium-5-yl)methyl}]-8-oxo-5-thia-1-
 azabicyclo[4.2.0]oct-2-ene-2-carboxylate

35



60

¹H NMR (400 MHz, D₂O): δ 2.97 (d, 1H, J=18 Hz), 3.25
 (d, 1H, J=18 Hz), 5.04 (s, 1H), 5.15 (s, 2H), 5.38 (s, 2H),
 5.66 (s, 1H), 6.23 (s, 1H), 6.96 (s, 1H), 7.38-7.52 (m, 4H),
 7.83 (s, 1H), 7.88 (s, 1H). Seven protons were not observed
 in D₂O.

65


Mass: ES⁺ 697.34

179

Example 33 (Table 1, Compound 23)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

5



¹H NMR (400 MHz, D₂O): δ 1.27 (s, 3H), 1.30 (s, 3H), 2.90 (d, 1H, J=18 Hz), 3.20 (d, 1H, J=18 Hz), 5.00 (d, 1H, J=5.0 Hz), 5.13 (s, 2H), 5.32 (s, 2H), 5.61 (d, 1H, J=5.0 Hz), 6.24 (s, 1H), 7.40-7.55 (m, 4H), 7.90 (d, 2H, J=4.0 Hz), 8.20 (s, 0.2H). Six protons were not observed in D₂O.

30


Mass: ES⁺ 760.29

Example 34 (Table 1, Compound 24)

35

(6R,7R)-7-[[[(2Z)-2-(5-Amino-1,2,4-thiadiazol-3-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

40



60

¹H NMR (400 MHz, D₂O): δ 1.25 and 1.29 (2s, 6H), 2.90 (d, 1H, J=18 Hz), 3.24 (d, 1H, J=18 Hz), 5.06 (s, 1H), 5.16 (s, 2H), 5.38 (s, 2H), 5.64 (s, 1H), 6.25 (s, 1H), 7.40-7.50 (m, 4H), 7.90 (s, 2H), 8.27 (s, 2.75H). Six protons were not observed in D₂O.


65

Mass: ES⁺ 726.27

Example 35 (Table 1, Compound 25)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-({1-[(4-carbamimidoylfuran-2-yl)methyl]-1H-
 imidazo[1,2-b]pyrazol-5-ium-5-yl}methyl)-8-oxo-5-
 thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

5



¹H NMR (400 MHz, D₂O): δ 1.30 (s, 3H), 1.32 (s, 3H), 2.95 (d, 1H, J=18 Hz), 3.24 (d, 1H, J=18 Hz), 5.03 (d, 1H, J=4.8 Hz), 5.13 (d, 2H, J=4.8 Hz), 5.29 (s, 2H), 5.62 (d, 1H, J=4.4 Hz), 6.26 (d, 1H, J=3.6 Hz), 6.85 (s, 1H), 6.92 (s, 1H), 7.38 (s, 1H), 7.82 (s, 1H), 7.89 (d, 1H, J=3.6 Hz), 8.13 (s, 1H). Seven protons were not observed in D₂O.


30

Mass: ES⁺ 697.25

Example 36 (Table 1, Compound 10)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-
 3-({1-[4-(2-carbamimidamidoethoxy)benzyl]-2,3-
 dihydro-1H-imidazo[1,2-b]pyrazol-5-ium-5-
 yl}methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-
 ene-2-carboxylate

35



¹H NMR (400 MHz, DMSO-d₆): δ 1.30 (s, 3H), 1.32 (s, 3H), 3.03 (d, 1H, J=17.6 Hz), 3.25 (d, 1H, J=17.6 Hz), 3.43 (t, 2H, J=4.7 Hz), 3.79 (t, 2H, J=8.2 Hz), 4.02-4.08 (m, 3H), 4.26 (m, 2H), 4.78 (d, 1H, J=15.3 Hz), 4.87 (d, 1H, J=15.7 Hz), 5.03 (d, 1H, J=5.1 Hz), 5.61 (dd, 2H, J=5.05 and 1.9 Hz), 6.84-6.86 (m, 3H), 7.17 (d, 2H, J=8.9 Hz), 7.78 (d, 1H, J=3.1 Hz), 8.18 (s, 1H). Eight protons were not observed in D₂O.

60


65

Mass: ES⁺ 768.31

Example 37 (Table 1, Compound 49)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-
 {[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-
 3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-
 1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-
 5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

5



¹H NMR (400 MHz, D₂O): δ 1.21 (s, 6H), 2.90 (d, 1H, J=17.6 Hz), 3.36 (d, 1H, J=18.0 Hz), 5.05-5.09 (m, 2H), 5.32 (d, 1H, J=14.4 Hz), 5.59 (s, 3H), 6.69 (s, 1H), 6.84 (d, 1H, J=3.2 Hz), 7.40 (d, 1H, J=9.6 Hz), 7.58 (d, 2H, J=10.0 Hz), 7.87 (d, 1H, J=7.2 Hz), 8.28 (d, 1H, J=7.2 Hz), 9.00 (s, 1H). Seven protons were not observed in D₂O.


30

Mass: ES⁺ 770.28

Example 38 (Table 1, Compound 27)

(6R,7R)-7-[[[(2Z)-2-(5-Amino-1,2,4-thiadiazol-3-yl)-
 2-[[[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-
 3-[[1-(4-carbamimidoylthiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-
 yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-
 ene-2-carboxylate

35



¹H NMR (400 MHz, D₂O): δ 1.28 (s, 3H), 1.30 (s, 3H), 2.90 (d, 1H, J=18.0 Hz), 3.22 (d, 1H, J=18.0 Hz), 5.01 (d, 1H, J=4.7 Hz), 5.12 (s, 2H), 5.43 (s, 2H), 5.62 (d, 1H, J=4.7 Hz), 6.17 (d, 1H, J=3.5 Hz), 7.38-7.40 (m, 2H), 7.80-7.88 (m, 2H), 8.12 (s, 1H). Seven protons were not observed in D₂O.


60

Mass: ES⁺ 714.27

185

Example 39 (Table 1, Compound 51)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-(1-[4-carbamimidoylthiophen-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (D₂O): δ 2.90 (d, 1H, J=17.6 Hz), 3.15 (d, 1H, J=17.6 Hz), 3.78 (s, 3H), 4.96 (d, 1H, J=4.8 Hz), 5.10 (d, 2H, J=3.6 Hz), 5.39 (s, 2H), 5.63 (d, 1H, J=4.8 Hz), 6.15 (d, 1H, J=3.6 Hz), 7.37 (d, 2H, J=10.8 Hz), 7.80 (d, 1H, J=2.4 Hz), 7.85 (d, 1H, J=2.4 Hz), 7.86 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 676.10

Example 40 (Table 1, Compound 50)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-(1-[4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (D₂O): δ 3.04 (d, 1H, J=17.6 Hz), 3.42 (d, 1H, J=18.0 Hz), 3.84 (s, 3H), 5.10 (d, 1H, J=4.8 Hz), 5.21 (d, 1H, J=14.8 Hz), 5.34 (d, 1H, J=14.4 Hz), 5.70 (m, 3H), 6.98 (d, 1H, J=3.2 Hz), 7.35 (s, 1H), 7.74 (d, 1H, J=3.2 Hz), 7.89 (d, 1H, J=6.8 Hz), 8.12 (s, 1H), 8.31 (d, 1H, J=2.4 Hz), 9.08 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 686.17

Example 41 (Table 1, Compound 52)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 2.97 (d, 1H, J=18.0 Hz), 3.35 (d, 1H, J=17.6 Hz), 3.77 (s, 3H), 5.03 (d, 1H, J=4.8 Hz), 5.14 (d, 1H, J=15.2 Hz), 5.26 (d, 1H, J=14.0 Hz), 5.56 (s, 2H), 5.65 (d, 1H, J=4.8 Hz), 6.91 (d, 1H, J=3.6 Hz), 7.15 (t, 1H, J=8.0 Hz), 7.35 (d, 1H, J=7.6 Hz), 7.41 (d, 1H, J=10.4 Hz), 7.65 (d, 1H, J=3.2 Hz), 7.77 (d, 1H, J=7.2 Hz), 8.22 (d, 1H, J=8.4 Hz), 9.01 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 698.16

Example 42 (Table 1, Compound 54)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-8-oxo-3-[[1-(4-{N-[(3R)-piperidin-3-yl]carbamimidoyl}benzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 1.56-1.92 (m, 2H), 1.88 (d, 1H, J=15.2 Hz), 2.07 (d, 1H, J=10.4 Hz), 2.83-3.01 (m, 3H), 3.21 (d, 1H, J=12.8 Hz), 3.32 (d, 1H, J=18.0 Hz), 3.47 (d, 1H, J=12.4 Hz), 3.74 (s, 3H), 3.92-3.97 (m, 1H), 4.99 (d, 1H, J=4.8 Hz), 5.13 (d, 1H, J=14.8 Hz), 5.23 (d, 1H, J=14.4 Hz), 5.51 (s, 2H), 5.60 (d, 1H, J=4.4 Hz), 6.90 (d, 1H, J=3.6 Hz), 7.17 (d, 2H, J=8.0 Hz), 7.47 (d, 2H, J=8.0 Hz), 7.63 (d, 1H, J=3.2 Hz), 7.66 (d, 1H, J=7.2 Hz), 8.16 (d, 1H, J=7.2 Hz), 9.00 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 763.22

Example 43 (Table 1, Compound 53)


(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({-[4-carbamimidoyl-1,3-thiazol-2-yl)methyl]-1H-imidazo[1,2-b]pyrazol-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



¹H NMR (400 MHz, D₂O): δ 3.02 (d, 1H, J=16.4 Hz), 3.25 (d, 1H, J=18.4 Hz), 3.83 (s, 3H), 5.04 (s, 1H), 5.11 (d, 1H, J=16.5 Hz), 5.20 (d, 1H, J=15.4 Hz), 5.67 (s, 3H), 6.34 (m, 1H), 7.51 (m, 1H), 7.87 (m, 1H), 7.95 (m, 1H), 8.46 (s, 1H). Six protons were not observed in D₂O.
Mass: ES⁺ 677.13

Example 44 (Table 1, Compound 55)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-({[1-(4-carbamimidoyl-2-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 2.96 (d, 1H, J=17.6 Hz), 3.36 (d, 1H, J=18.0 Hz), 3.75 (s, 3H), 5.04 (d, 1H, J=4.8 Hz), 5.12 (d, 1H, J=14.8 Hz), 5.26 (d, 1H, J=14.8 Hz), 5.55 (s, 2H), 5.62 (d, 1H, J=4.8 Hz), 6.75 (s, 1H), 6.90 (d, 1H, J=2.8 Hz), 7.11 (t, 1H, J=8.0 Hz), 7.33 (d, 1H, J=8.0 Hz), 7.40 (d, 1H, J=10.8 Hz), 7.64 (d, 1H, J=3.2 Hz), 7.75 (d, 1H, J=6.8 Hz), 8.21 (d, 1H, J=7.2 Hz), 9.00 (s, 1H). Six protons were not observed in D₂O.
Mass: ES⁺ 664.23


191

Example 45 (Table 1, Compound 58)

(6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



Step 1: 3-Chloro-5-fluoro-4-formylbenzonitrile




To a solution of 3-chloro-5-fluorobenzonitrile (1.8 g, 11.57 mmol) in anhydrous tetrahydrofuran (40 mL) at -78°C . was added lithium diisopropylamide (2M, 6.4 mL, 12.73 mmol). After stirring at same temperature for 30 minutes was added dimethylformamide (1.02 g, 13.84 mmol). The reaction mixture was stirred at -78°C . for an additional 15 minutes, then was added acetic acid (2.7 mL) followed by water (40 mL) and the mixture was warmed to room temperature. After extraction with ethyl acetate (60 mL), the organic phase was washed with 1M HCl solution, brine, dried over Na_2SO_4 , filtered and evaporated to give 3-chloro-5-fluoro-4-formylbenzonitrile (1.08 g, 51%) as a gummy solid.

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 7.42 (m, 1H), 7.61 (d, 1H, $J=2.8$ Hz), 10.45 (s, 1H).

Step 2:

3-Chloro-5-fluoro-4-(hydroxymethyl)benzonitrile



To a solution of 3-chloro-5-fluoro-4-formylbenzonitrile (from step 1, 1.08 g, 5.88 mmol) in methanol (6 mL) was added sodium borohydride (0.25 g, 6.61 mmol) at 0°C . After 15 minutes, the mixture was quenched with water (1 mL) and concentrated to obtain a yellow oil residue. Dis-


192

solved the residue in ethyl acetate (35 mL), washed with water (30 mL) and the aqueous layer was extracted again with ethyl acetate (20 mL). The extracts were combined, washed with brine, dried over Na_2SO_4 , filtered and evaporated. The residue was purified by column chromatography (silica gel) eluting 1:1 ethyl acetate/hexanes as eluent to give 3-chloro-5-fluoro-4-(hydroxymethyl)benzonitrile (0.65 g, 60%) as an orange thick oil.

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 4.88 (s, 2H), 7.33 (m, 1H), 7.54 (d, 1H, $J=1.2$ Hz).

Step 3:


4-(Bromomethyl)-3-chloro-5-fluorobenzonitrile



To a solution of 3-chloro-5-fluoro-4-(hydroxymethyl)benzonitrile (from step 2, 0.65 g, 3.5 mmol) and carbon tetrabromide (1.25 g, 3.73 mmol) in dichloromethane (20 mL) was added triphenylphosphene (1.02 g, 2.73 mmol) slowly at 0°C . After stirring for 10 minutes at the same temperature, the solution was stirred overnight at room temperature. The solution was concentrated under reduced pressure and the residue was purified by column chromatography (silica gel) eluting with 1:20 ethyl acetate/hexanes as eluent to give 4-(bromomethyl)-3-chloro-5-fluorobenzonitrile (0.72 g, 83%) as a white solid.

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 4.59 (s, 2H), 7.32 (m, 1H), 7.54 (s, 1H, $J=1.2$ Hz).

Step 4: 3-Chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzonitrile




193

An ice-cold mixture of 5-azaindole (0.34 g, 2.88 mmol) in N,N-dimethylformamide (10 mL) was treated with sodium hydride (60% in mineral oil, 0.172 g, 4.32 mmol) in portions, then stirred at room temperature for 15 minutes. The mixture was cooled to 0° C., and then treated with 4-(bromomethyl)-3-chloro-5-fluorobenzonitrile (from step 3, 0.73 g, 2.93 mmol) in tetrahydrofuran (10 mL). The resulting mixture was stirred at 0° C. for 1 hour then quenched with saturated ammonium chloride solution. The mixture was extracted with ethyl acetate, and the extracts were washed with brine, dried (sodium sulphate), filtered and concentrated in vacuo to brown oil, which was purified by column chromatography (silica gel) eluting with 5:3:2 dichloromethane/ethyl acetate/methanol as eluent to afford 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzonitrile (0.53 g, 65%) as a white solid.


¹H NMR (400 MHz, CDCl₃): δ 5.49 (d, 2H, J=1.9 Hz), 6.62 (d, 1H, J=3.2 Hz), 7.19 (d, 1H, J=2.8 Hz), 7.37-7.41 (m, 2H), 7.59 (s, 1H), 8.35 (d, 1H, J=6.0 Hz), 8.91 (s, 1H).

Step 5: Ethyl 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride



Through a solution of 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzonitrile (from step 4, 0.53 g, 1.86 mmol) in anhydrous ethanol (20 mL) in a pressure vessel at 0° C. was bubbled a stream of anhydrous hydrogen chloride gas for 10 min. The reaction vessel was stoppered and stirred at room temperature for 18 h and the contents were transferred into a flask and evaporated under reduced pressure. The crude product was triturated with ether, then dried under vacuum to provide ethyl 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (0.64 g, 94%) as a white solid which was used in the next step without further purification.


Step 6: 3-Chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidamide



194

A pressure vessel containing a suspension of ethyl 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (from step 5, 0.64 g, 1.74 mmol) in methanol (20 mL) was saturated with ammonia gas and stoppered. The reaction mixture was stirred at room temperature for 24 h, then excess ammonia was vented out and the contents were concentrated under reduced pressure. The residue was triturated with hexanes and dried under vacuum to give 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidamide (0.69 g, 130%) as a grey solid which was used in the next step without further purification.

Step 7: tert-Butyl {[3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)phenyl]carboximidoyl}carbamate



A solution of 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)benzenecarboximidamide (from step 6, 0.69 g, 1.74 mmol) in 1,4-dioxane (50 mL) was treated with a saturated sodium carbonate solution (25 mL) followed by di-tert-butyl dicarbonate (2.51 g, 11.4 mmol) and stirred at room temperature for 40 h. The reaction mixture was concentrated under reduced pressure to remove the volatiles and the remaining solution was dissolved in ethyl acetate (60 mL), then washed with water (60 mL), brine solution (60 mL), dried and concentrated. The crude product was purified by silica gel column chromatography using ethyl acetate:methanol:ammonium hydroxide (60:39:1) as eluent to afford tert-butyl {[3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)phenyl]carboximidoyl}carbamate (0.3 g, 33%).


¹H NMR (400 MHz, CDCl₃): δ 1.52 (s, 9H), 5.43 (s, 2H), 6.56 (d, 1, J=3.2 Hz), 7.18 (d, 1H, J=2.8 Hz), 7.37 (d, 1H, J=6.0 Hz), 7.58 (d, 1H, J=9.6 Hz), 7.80 (s, 1H), 8.22 (d, 1H, J=5.6 Hz), 8.82 (s, 1H).

195

196

Step 8: 5-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-Butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[4-[N-(tert-butoxycarbonyl)carbamimidoyl]-2-chloro-6-fluorobenzyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide

5



To a solution of tert-butyl {[3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-c]pyridin-1-ylmethyl)phenyl]carbamimidoyl}carbamate (from step 7, 0.12 g, 0.30 mmol) in dimethylformamide (1 mL) was added (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-3-(iodomethyl)-2-[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thionia-1-azabicyclo[4.2.0]oct-2-en-5-olate (0.24 g, 0.30 mmol) at 0° C. The reaction mixture was then degassed under reduced pressure for 0.5 h, and stirred at room temperature for 16 h. Dimethyl formamide (2 mL) was added and cooled to -40° C., followed by adding KI (0.35 g, 2.1 mmol) and acetyl chloride (0.09 mL, 1.2 mmol). After stirring at 0° C.


40

45

50

for 1 h, ice water and ethyl acetate (30 mL) were poured to the reaction solution, and then NaHSO₃ solution (20 mL) was added. The aqueous solution was extracted with ethyl acetate (30 mL×2) and the organic layers were combined, washed with brine (3×50 mL), dried over Na₂SO₄, filtered and evaporated to give yellow solid (0.29 g), which was used in the next step without further purification.

Step 9: (6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



197

To a solution of 5-[(6R,7R)-7-[(2Z)-2-{2-[(tert-butoxy-carbonyl)amino]-5-chloro-1,3-thiazol-4-yl}-2-(methoxy-imino)acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbo-nyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[4-[N-(tert-butoxycarbonyl)carbamimidoyl]-2-chloro-6-fluorobenzyl]-1H-pyrrolo[3,2-c]pyridin-5-ium iodide (0.29 g) in dry dichloromethane (6 mL) at -40° C. was added anisole (0.52 mL, 4.92 mmol) and then 2M AlCl_3 in CH_3NO_2 (2.46 mL, 4.92 mmol). The liquid was stirred at 0° C. for 30 min. To the reaction mixture were added diisopropyl ether (10 mL) and water (0.5 mL), and the resultant was stirred to generate a precipitate. The supernatant was removed by decantation. To the insoluble matter adhering to the vessel were added diluted aqueous hydrochloric acid solution (5 mL) and acetonitrile (10 mL). The resultant was stirred to dissolve the matter completely. Thereto was added HP20 resin (1.0 g), and stirred for 30 min, and then filtered. The filtrate was concentrated and lyophilized to give a crude product which was purified by HPLC to obtain (6R,7R)-7-[(2Z)-2-(2-amino-5-chloro-1,3-

198


thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate as formic acid salt (0.0135 g, 7.5%).

^1H NMR (400 MHz, D_2O): δ 2.94 (d, 1H, $J=18.0$ Hz), 3.33 (d, 1H, $J=17.6$ Hz), 3.73 (s, 3H), 5.00 (d, 1H, $J=5.2$ Hz), 5.11 (d, 1H, $J=14.4$ Hz), 5.26 (d, 1H, $J=14.0$ Hz), 5.58-5.62 (m, 3H), 6.84 (s, 1H), 7.39 (d, 1H, $J=9.6$ Hz), 7.58 (s, 2H), 7.89 (d, 1H, $J=7.2$ Hz), 8.25 (d, 1H, $J=6.4$ Hz), 8.97 (s, 1H). Six protons were not observed in D_2O .

Mass: ES^+ 732.21

Example 46 (Table 1, Compound 56)

(6R,7R)-7-[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-([1-(4-carbamimidoylfuran-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl)methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




^1H NMR (400 MHz, D_2O): δ 2.91 (d, 1H, $J=18.0$ Hz), 3.34 (d, 1H, $J=18.0$ Hz), 3.77 (s, 3H), 5.03 (d, 1H, $J=4.8$ Hz), 5.15 (d, 1H, $J=15.2$ Hz), 5.26 (d, 1H, $J=14.0$ Hz), 5.47 (s, 2H), 5.65 (d, 1H, $J=4.8$ Hz), 6.74 (s, 1H), 6.88 (d, 1H, $J=3.2$ Hz), 7.64 (d, 1H, $J=3.6$ Hz), 7.86 (d, 1H, $J=7.6$ Hz), 8.04 (s, 1H), 8.25 (d, 1H, $J=4.0$ Hz), 8.99 (s, 1H). Six protons were not observed in D_2O .

Mass: ES^+ 670.06

Example 47 (Table 1, Compound 65)

(6R,7R)-7-[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino}-3-([1-(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl)methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




199

¹HNMR (400 MHz, D₂O): δ 1.33 (s, 3H), 1.35 (s, 3H), 2.95 (d, 1H, J=18.0 Hz), 3.38 (d, 1H, J=18.4 Hz), 5.07-5.11 (m, 2H), 5.39 (d, 1H, J=14.5 Hz), 5.67-5.78 (m, 3H), 6.88 (d, 1H, J=2.8 Hz), 7.47 (s, 1H), 7.96 (d, 1H, J=6.0 Hz), 8.12 (m, 3H), 9.10 (s, 1H). Seven protons were not observed in D₂O. 5
Mass: ES⁺ 758.17

Example 48 (Table 1, Compound 66)


(6R,7R)-7-[[[(2Z)-2-(5-Amino-1,2,4-thiadiazol-3-yl)-
2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[4-carbamimidoylthiophen-2-yl]
methyl]-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl}methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate 10



¹H NMR (400 MHz, D₂O): δ 1.37 (s, 3H), 1.39 (s, 3H), 2.97 (d, 1H, J=18.0 Hz), 3.38 (d, 1H, J=17.6 Hz), 5.08-5.13 (m, 2H), 5.37 (d, 1H, J=14.4 Hz), 5.69-5.80 (m, 3H), 6.85 (d, 1H, J=2.8 Hz), 7.46 (s, 1H), 7.94 (d, 1H, J=6.8 Hz), 8.09-8.16 (m, 3H), 9.08 (s, 1H). Seven protons were not observed in D₂O. 35
Mass: ES⁺ 725.16 40

Example 49 (Table 1, Compound 67)

(6R,7R)-7-[[[(2Z)-2-(5-Amino-1,2,4-thiadiazol-3-yl)-
2-(methoxyimino)acetyl]amino]-3-({1-[4-carbamimidoylthiophen-2-yl]methyl]-1H-pyrrolo[2,3-c]
pyridin-6-ium-6-yl}methyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate 45



201

202

¹H NMR (400 MHz, D₂O): δ 2.96 (d, 1H, J=17.6 Hz), 3.38 (d, 1H, J=17.6 Hz), 3.95 (s, 3H), 5.06 (d, 1H, J=4.8 Hz), 5.12 (d, 1H, J=14.8 Hz), 5.31 (d, 1H, J=14.4 Hz), 5.53 (d, 1H, J=4.8 Hz), 5.72 (m, 2H), 6.88 (d, 1H, J=3.2 Hz), 7.50 (s, 1H), 7.95 (d, 1H, J=6.4 Hz), 8.11-8.30 (m, 3H), 8.99 (s, 1H). Six protons were not observed in D₂O.


Mass: ES⁺ 653.19

Example 50 (Table 1, Compound 68)

10

(6R,7R)-7-[[[(2Z)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino]acetyl]amino]-3-({1-[(4-carbamimidoylthiophen-2-yl)methyl]-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl)methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

15



35

¹H NMR (400 MHz, D₂O): δ 1.40 (s, 6H), 3.03 (d, 1H, J=17.6 Hz), 3.24 (d, 1H, J=17.6 Hz), 5.06 (d, 1H, J=4.8 Hz), 5.42 (d, 1H, J=15.2 Hz), 5.58 (d, 1H, J=15.2 Hz), 5.71 (d, 1H, J=4.8 Hz), 5.76 (s, 2H), 6.95 (d, 1H, J=3.2 Hz), 7.34 (s, 1H), 7.54-7.58 (m, 1H), 8.08 (d, 1H, J=3.2 Hz), 8.12 (s, 1H), 8.48-8.53 (m, 2H). Seven protons were not observed in D₂O.

40


Mass: ES⁺ 725.13

Example 51 (Table 1, Compound 71)

45


(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate

50



203


Step 1: 3-Chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzonitrile



An ice-cold mixture of 4-azaindole (0.47 g, 3.98 mmol) in N,N-dimethylformamide (10 mL) was treated with sodium hydride (60% in mineral oil, 0.23 g, 6.0 mmol) in portions, then stirred at room temperature for 15 minutes. The mixture was cooled to 0° C., then treated with 4-(bromomethyl)-3-chloro-5-fluorobenzonitrile (1.00 g, 4.00 mmol) in tetrahydrofuran (10 mL). The resulting mixture was stirred at 0° C. for 1 hour then quenched with saturated ammonium chloride solution. The mixture was extracted with ethyl acetate and the extracts were washed with brine, dried (sodium sulphate), filtered and concentrated in vacuo to brown oil, which was purified by column chromatography (silica gel) eluting with 5:3:2 dichloromethane/ethyl acetate/methanol as eluent to afford 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzonitrile (0.62 g, 54%) as brown solid.

¹H NMR (400 MHz, CDCl₃): δ 5.49 (d, 2H, J=1.9 Hz), 6.72 (d, 1H, J=3.4 Hz), 7.14 (dd, 1H, J=8.2, 4.7 Hz), 7.37 (d, 1H, J=8.7 Hz), 7.41 (d, 1H, J=3.4 Hz), 7.58 (br s, 1H), 7.78 (d, 1H, J=8.4 Hz), 8.47 (d, 1H, J=4.7 Hz).

Step 2: Ethyl 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride



Anhydrous hydrochloric acid was bubbled into an ice-cold brown suspension of 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzonitrile (0.65 g, 2.23 mmol) in anhydrous ethanol (30 mL) for 10 minutes. The mixture was sealed and stirred at room temperature overnight. The mixture was concentrated in vacuo to afford ethyl 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (0.83 g) as a brown solid, which was used in the next step without further purification.

¹H NMR (400 MHz, CD₃OD): δ 1.58 (t, 3H, J=7.2 Hz), 4.59 (q, 2H, J=7.0 Hz), 6.09 (s, 2H), 7.15 (d, 1H, J=3.2 Hz),

204


7.92-8.04 (m, 2H), 8.16 (s, 1H), 8.34 (d, 1H, J=2.9 Hz), 8.83 (d, 1H, J=5.6 Hz), 9.03 (d, 1H, J=8.2 Hz).

Step 3: 3-Chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzenecarboximidamide

5

10

15



Anhydrous ammonia was bubbled into an ice-cold solution of 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (0.83 g, 2.23 mmol) in anhydrous ethanol (30 mL) for 10 minutes. The mixture was sealed and stirred at room temperature overnight. The mixture was concentrated in vacuo to afford 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzenecarboximidamide as a brown solid (0.72 g), which was used in the next step without further purification.


¹H NMR (400 MHz, CD₃OD): δ 5.59 (s, 2H), 6.62 (br s, 1H), 7.22 (br s, 1H), 7.54-7.60 (m, 2H), 7.64-7.69 (m, 1H), 7.82-7.87 (m, 1H), 8.23-8.43 (br s, 1H).

Step 4: tert-Butyl {[3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)phenyl]carboximidoyl}carbamate

35

40

45



50

55

60

65


An ice-cold solution of 3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)benzenecarboximidamide (0.72 g, 2.38 mmol) in dioxane (40 mL) and saturated sodium carbonate solution (20 mL) was treated with di-tert-butyl dicarbonate (2.08 g, 9.53 mmol). The mixture was stirred at room temperature overnight, then concentrated in vacuo. The residue was diluted with ethyl acetate and washed with water and brine, then dried (sodium sulphate), filtered and concentrated to brown solid, which was purified by column chromatography (silica gel) eluting with 2-4% methanol in dichloromethane to afford tert-butyl {[3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)phenyl]carboximidoyl}carbamate (0.20 g, 21%) as a yellow solid.

¹H NMR (400 MHz, CDCl₃): δ 1.55 (s, 9H), 5.47 (d, 2H, J=1.8 Hz), 6.60 (d, 1H, J=3.2 Hz), 7.19 (d, 1H, J=3.1 Hz), 7.41 (d, 1H, J=6.0 Hz), 7.55-7.60 (m, 1H), 7.78 (s, 1H), 8.31 (d, 1H, J=6.0 Hz), 8.88 (s, 1H).

205

Step 5: 4-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-Butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-{4-[N-(tert-butoxycarbonyl)carbamimidoyl]-2-chloro-6-fluorobenzyl]-1H-pyrrolo[3,2-b]pyridin-4-ium iodide

5




An ice-cold mixture of tert-butyl [[3-chloro-5-fluoro-4-(1H-pyrrolo[3,2-b]pyridin-1-ylmethyl)phenyl]carbamimidoyl]carbamate (0.20 g, 0.50 mmol) in anhydrous N,N-dimethylformamide (2.0 mL) was degassed under reduced pressure for 5 minutes, then treated with 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.34 g, 0.50 mmol). The resulting mixture was degassed for 30 minutes, and treated with sodium iodide (0.15 g, 1.00 mmol). The mixture was stirred at room temperature overnight, cooled to 0° C. and quenched with a 5% aqueous solution of sodium chloride and sodium thiosulfate to give an orange suspension. The

35

solid was filtered, washed with water and dried in vacuo to afford 4-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[[4-methoxybenzyl]oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-{4-[N-(tert-butoxycarbonyl)carbamimidoyl]-2-chloro-6-fluorobenzyl]-1H-pyrrolo[3,2-b]pyridin-4-ium iodide, which was used in the next step without further purification.

45

Step 6: (6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



207

A solution of 4-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxy-carbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxy-imino)acetyl]amino}-2-[[4-methoxybenzyl]oxy]carbo-nyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-{4-[N-(tert-butoxycarbonyl)carbamimidoyl]-2-chloro-6-fluorobenzyl]-1H-pyrrolo[3,2-b]pyridin-4-ium iodide (0.50 mmol) in dichloromethane (2.5 mL) and anisole (0.85 mL, 7.82 mmol) was cooled to 0° C., then treated with trifluoroacetic acid (2.5 mL, 32.6 mmol). The resulting brown solution was stirred at 0° C. for 10 minutes, then at room temperature for 2 hours. The mixture was concentrated in vacuo to a brown residue, which was washed with diethyl ether and the brown solid was purified by preparative HPLC (0.1% formic acid in water/acetonitrile) to afford (6R,7R)-7-[[[(2Z)-2-(2-amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-b]pyridin-4-ium-4-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.050 g, 14% over 2 steps) as formic acid salt.

208


¹H NMR (400 MHz, D₂O): δ 3.18 (d, 1H, J=17.9 Hz), 3.32 (d, 1H, J=17.9 Hz), 3.95 (s, 3H), 5.12 (d, 1H, J=5.0 Hz), 5.50 (d, 1H, J=14.7 Hz), 5.71 (d, 1H, J=15.0 Hz), 5.80 (d, 1H, J=4.7 Hz), 5.85 (s, 2H), 7.01 (d, 1H, J=3.5 Hz), 7.62 (dd, 1H, J=9.7, 1.7 Hz), 7.70 (dd, 1H, J=8.2, 6.2 Hz), 7.80 (s, 1H), 8.15 (d, 1H, J=3.2 Hz), 8.62 (d, 1H, J=6.2 Hz), 8.69 (d, 1H, J=8.5 Hz).

Six protons were not observed in D₂O.

Mass: ES⁺ 732.15.

Example 52 (Table 1, Compound 69)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[3,2-c]pyridin-5-ium-5-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




¹H NMR (400 MHz, D₂O): δ 3.14 (d, 1H, J=17.9 Hz), 3.58 (d, 1H, J=17.9 Hz), 3.93 (s, 3H), 5.22 (d, 1H, J=5.0 Hz), 5.29 (d, 1H, J=14.7 Hz), 5.48 (d, 1H, J=14.4 Hz), 5.74-5.79 (m, 3H), 6.82 (s, 1H), 7.02 (d, 1H, J=3.2 Hz), 7.60 (dd, 1H, J=9.7, 1.5 Hz), 7.73-7.81 (m, 2H), 8.06 (d, 1H, J=7.3 Hz), 8.44 (d, 1H, J=7.0 Hz), 9.17 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 698.19.


Example 53 (Table 1, Compound 70)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate formic acid salt



209


Step 1: 3-Chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzonitrile



An ice-cold mixture of 6-azaindole (0.98 g, 7.96 mmol) in N,N-dimethylformamide (10 mL) was treated with sodium hydride (60% in mineral oil, 0.46 g, 12.0 mmol) in portions, then stirred at room temperature for 15 minutes. The mixture was cooled to 0° C., then treated with 4-(bromomethyl)-3-chloro-5-fluorobenzonitrile (2.00 g, 8.05 mmol) in tetrahydrofuran (10 mL). The resulting mixture was stirred at 0° C. for 1 hour then quenched with saturated ammonium chloride solution. The mixture was extracted with ethyl acetate, and the extracts were washed with brine, dried (sodium sulphate), filtered and concentrated in vacuo to brown oil, which was purified by column chromatography (silica gel) eluting with 5:3:2 dichloromethane/ethyl acetate/methanol as eluent to afford 3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzonitrile (1.14 g, 50%) as a brown solid.

¹H NMR (400 MHz, CDCl₃): δ 5.59 (d, 2H, J=1.9 Hz), 6.54 (dd, 1H), 7.31 (dd, 1H, J=3.15, 1.25 Hz), 7.38 (dd, 1H, J=8.7, 1.6 Hz), 7.51 (dd, 1H, J=5.4, 1.0 Hz), 7.59 (t, 1H, J=1.4 Hz), 8.27 (d, 1H, J=5.4 Hz), 8.92 (s, 1H).


Step 2: Ethyl 3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride



Anhydrous hydrochloric acid was bubbled into an ice-cold brown suspension of 3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzonitrile (1.14 g, 3.99 mmol) in anhydrous ethanol (30 mL) for 10 minutes. The mixture was sealed and stirred at room temperature overnight. The mixture was concentrated in vacuo to afford ethyl 3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (1.16 g) as a brown solid, which was used in the next step without further purification.

210


Step 3: 3-Chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzenecarboximidamide



Anhydrous ammonia was bubbled into an ice-cold solution of ethyl 3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzenecarboximidate hydrochloride (1.6 g, 4.34 mmol) in anhydrous ethanol (30 mL) for 10 minutes. The mixture was sealed and stirred at room temperature overnight. The mixture was concentrated in vacuo to afford 3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzenecarboximidamide as a brown solid (0.59 g), which was used in the next step without further purification.

¹H NMR (400 MHz, CD₃OD): δ 5.80 (s, 2H), 6.64-6.71 (m, 1H), 7.55-7.74 (m, 3H), 7.84-7.91 (m, 1H), 8.12 (s, 1H), 8.92 (s, 1H).

Step 4: tert-Butyl {[3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)phenyl]carboximidoyl}carbamate



An ice-cold solution of 3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)benzenecarboximidamide (0.59 g, 1.95 mmol) in dioxane (40 mL) and saturated sodium carbonate solution (20 mL) was treated with di-tert-butyl dicarbonate (1.70 g, 7.78 mmol). The mixture was stirred at room temperature overnight and concentrated in vacuo. The residue was diluted with ethyl acetate, washed with water and brine, then dried (sodium sulphate), filtered and concentrated to a brown solid. The solid was purified by column chromatography (silica gel) eluting with 2-4% methanol in dichloromethane to afford tert-butyl {[3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)phenyl]carboximidoyl}carbamate (0.64 g, 82%) as a yellow solid.


¹H NMR (400 MHz, CDCl₃): δ 1.55 (s, 9H), 5.53 (d, 2H, J=1.5 Hz), 6.49 (d, 1H, J=3.2 Hz), 7.30 (d, 1H, J=3.2 Hz), 7.48 (d, 1H, J=5.4 Hz), 7.57-7.63 (m, 1H), 7.81 (s, 1H), 8.21 (d, 1H, J=5.4 Hz), 8.91 (s, 1H).

211

212

Step 5: 6-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-Butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-{4-[N-(tert-butoxycarbonyl)carbamimidoyl]-2-chloro-6-fluorobenzyl]-1H-pyrrolo[2,3-c]pyridin-6-ium iodide


5



An ice-cold mixture of tert-butyl [[3-chloro-5-fluoro-4-(1H-pyrrolo[2,3-c]pyridin-1-ylmethyl)phenyl]carbamimidoyl]carbamate (0.20 g, 0.50 mmol) in anhydrous N,N-dimethylformamide (2.0 mL) was degassed under reduced pressure for 5 minutes, then treated with 4-methoxybenzyl (6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-3-(chloromethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.34 g, 0.50 mmol). The resulting mixture was degassed for 30 minutes, then treated with sodium iodide (0.15 g, 1.00 mmol). The mixture was stirred at room temperature overnight, then cooled to 0° C. and quenched with a 5% aqueous solution of sodium chloride and sodium thiosulfate to give an orange suspension.

The solid was filtered, washed with water and dried in vacuo to afford 6-[[[(6R,7R)-7-[[[(2Z)-2-{2-[(tert-butoxycarbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxyimino)acetyl]amino}-2-[(4-methoxybenzyl)oxy]carbonyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-{4-[N-(tert-butoxycarbonyl)carbamimidoyl]-2-chloro-6-fluorobenzyl]-1H-pyrrolo[2,3-c]pyridin-6-ium iodide, which was used in the next step without further purification.

Step 6: (6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino}-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



213

A solution of 6-[[[(6R,7R)-7-[[[(2Z)-2-[[[(tert-butoxy-carbonyl)amino]-5-chloro-1,3-thiazol-4-yl]-2-(methoxy-imino)acetyl]amino]-2-[[[(4-methoxybenzyl)oxy]carbo-nyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-[4-[N-(tert-butoxycarbonyl)carbamimidoyl]-2-chloro-6-fluorobenzyl]-1H-pyrrolo[2,3-c]pyridin-6-ium iodide (0.50 mmol) in dichloromethane (2.5 mL) and anisole (0.85 mL, 7.82 mmol) was cooled to 0° C., then treated with trifluoroacetic acid (2.5 mL, 32.6 mmol). The resulting brown solution was stirred at 0° C. for 10 minutes, then at room temperature for 2 hours. The mixture was concentrated in vacuo to a brown residue, which was washed with diethyl ether to afford the crude product as brown solid. The solid was purified by preparative HPLC (0.1% formic acid in water/acetonitrile) to afford (6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate (0.025 g, 7% over 2 steps) as formic acid salt.


214

¹H NMR (400 MHz, D₂O): δ 3.18 (d, 1H, J=18.2 Hz), 3.59 (d, 1H, J=17.9 Hz), 3.95 (s, 3H), 5.21-5.32 (m, 2H), 5.56 (d, 1H, J=14.1 Hz), 5.79-5.90 (m, 3H), 6.96 (d, 1H, J=3.2 Hz), 7.62 (dd, 1H, J=9.5, 1.60 Hz), 7.78 (s, 1H), 8.05 (d, 1H, J=6.8 Hz), 8.20 (d, 1H, J=3.2 Hz), 8.24 (d, 1H, J=6.8 Hz), 9.09 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 732.04.

Example 54 (Table 1, Compound 72)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-5-chloro-1,3-thiazol-4-yl)-2-[[[(2-carboxypropan-2-yl)oxy]imino}acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate



¹H NMR (400 MHz, D₂O): δ 1.46 (s, 3H), 1.47 (s, 3H), 3.16 (d, 1H, J=18.2 Hz), 3.61 (d, 1H, J=17.9 Hz), 5.21 (d, 1H, J=14.1 Hz), 5.28 (d, 1H, J=5.0 Hz), 5.61 (d, 1H, J=14.4 Hz), 5.79-5.91 (m, 3H), 6.95 (d, 1H, J=2.6 Hz), 7.61 (d, 1H, J=10.0 Hz), 7.78 (s, 1H), 8.04 (d, 1H, J=6.8 Hz), 8.21 (d, 1H, J=2.4 Hz), 8.24 (d, 1H, J=6.8 Hz), 9.13 (s, 1H). Six protons were not observed in D₂O.

Mass: ES⁺ 804.13

Example 55 (Table 1, Compound 73)

(6R,7R)-7-[[[(2Z)-2-(2-Amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[1-(4-carbamimidoyl-2-chloro-6-fluorobenzyl)-1H-pyrrolo[2,3-c]pyridin-6-ium-6-yl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate




TABLE 2-continued

MIC values of selected compounds alone and in combination with NXL-104 against G(+) and G(-) isolates											
Isolate	Resistance Mechanism	CAZ:		Ex-1	Ex-1: NXL104	EX-2	EX-2: NXL104	Ex-5	EX-5: NXL104	Ex-6	Ex-6: NXL104
		CAZ	NXL104								
S. pneumoniae 2404	Pen S	32.00	32.00	0.25	0.12	0.25	0.25	≤0.06	≤0.06	≤0.06	≤0.06
S. pneumoniae 2405	Pen S	0.25	0.25	≤0.06	≤0.06	0.12	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06
S. pneumoniae 2406	Pen S	0.12	0.12	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06
S. pneumoniae 2407	Pen S	4.00	32.00	0.12	0.12	0.12	0.12	≤0.06	≤0.06	≤0.06	≤0.06
E. cloacae 2704	P99	16.00	4.00	16.00	16.00	4.00	1.00	16.00	2.00	32.00	1.00
E. cloacae 551	Caz R, IMP S	16.00	4.00	32.00	1.00	8.00	1.00	32.00	1.00	16.00	1.00
E. cloacae 552	Caz S, IMP S	0.50	0.25	0.12	0.12	0.12	0.12	0.12	≤0.06	0.12	0.25
E. cloacae 566	Caz R, IMP S	>32	2.00	4.00	0.50	0.50	0.25	2.00	0.50	4.00	0.50
E. cloacae 596	Caz R, IMP S	4.00	1.00	4.00	0.50	0.50	0.25	8.00	0.25	8.00	0.50
E. cloacae 2044	Caz S, IMP S	1.00	0.25	0.25	0.12	0.25	0.12	0.50	0.12	0.12	≤0.06
C. freundii 15	Caz R, IMP S	8.00	2.00	16.00	0.50	4.00	0.50	16.00	0.50	8.00	0.50
C. freundii 579	Caz R, IMP S	4.00	2.00	4.00	0.25	2.00	0.50	4.00	0.25	2.00	0.25
C. freundii 580	Caz R, IMP S	8.00	2.00	16.00	0.50	4.00	0.50	8.00	0.25	4.00	0.25
C. freundii 2003	Caz R, IMP S	4.00	1.00	8.00	0.25	2.00	0.50	8.00	0.25	4.00	0.50
C. freundii 2487	Caz R, IMP S	16.00	8.00	32.00	1.00	8.00	1.00	32.00	1.00	8.00	1.00
K. pneumoniae 4104	KPC-3, TEM-1	32.00	8.00	16.00	2.00	8.00	2.00	32.00	2.00	32.00	2.00
K. pneumoniae 4105	KPC-3, TEM-1, SHV-11	32.00	4.00	16.00	2.00	32.00	2.00	32.00	1.00	32.00	1.00
K. pneumoniae 4106	KPC-3, TEM-1, SHV-12, SHV-141	16.00	4.00	8.00	1.00	8.00	2.00	16.00	1.00	16.00	1.00
K. pneumoniae 4107	KPC-2, TEM-1, SHV-11	32.00	8.00	16.00	2.00	16.00	4.00	32.00	2.00	32.00	2.00
K. pneumoniae 4108	KPC-3, TEM-166, SHV-12, SHV-141	16.00	4.00	4.00	0.50	4.00	1.00	8.00	0.50	16.00	0.50
K. pneumoniae 4109	SHV-1, SHV-12	2.00	0.50	2.00	0.12	0.50	0.12	8.00	0.12	8.00	0.12
M. morgani 2409	High level cephalase (ACC 43: 769-776, 1999), β-Lactamase producing	16.00	4.00	8.00	1.00	8.00	1.00	16.00	0.50	16.00	1.00
K. pneumoniae 2689	KPC-2	32.00	4.00	8.00	2.00	8.00	2.00	8.00	1.00	16.00	1.00
E. coli 2671	VIM-19	32.00	32.00	16.00	16.00	4.00	4.00	16.00	16.00	32.00	32.00
K. pneumoniae 2697	IMP-4	8.00	2.00	4.00	0.50	4.00	1.00	16.00	0.50	16.00	0.50
E. coli 4080	TEM-10	16.00	4.00	4.00	1.00	8.00	1.00	16.00	0.50	16.00	0.50
E. coli 4098	CTX-M-15, TEM-1	2.00	0.50	4.00	0.25	1.00	0.25	>32	0.12	32.00	0.12
E. coli 4101	CTX-M-14, TEM-1	0.50	0.25	0.50	0.12	0.50	0.12	0.50	0.12	0.50	≤0.06
E. coli 4102	CTX-M-14	1.00	0.25	1.00	0.12	1.00	0.12	1.00	≤0.06	2.00	0.12
E. coli 102	TEM-1	0.12	0.12	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06	≤0.06
K. baumannii 2579	OXA-25	>32	>32	32.00	8.00	>32	32.00	>32	8.00	>32	16.00
K. baumannii 4091	OXA-23	>32	16.00	16.00	4.00	8.00	2.00	8.00	2.00	16.00	4.00
K. baumannii 4094	OXA-51	>32	32.00	16.00	32.00	32.00	8.00	16.00	4.00	16.00	8.00
K. baumannii 4095	OXA-48	0.50	0.50	0.50	0.25	0.25	0.25	0.25	0.25	0.12	0.25
E. coli 2668	OXA-48	1.00	0.50	1.00	0.12	0.50	0.25	1.00	≤0.06	0.50	0.12
P. aeruginosa 2573	OXA-10 (Class D)	8.00	8.00	4.00	2.00	2.00	4.00	4.00	4.00	2.00	2.00

TABLE 3

MIC values of selected compounds alone and in combination with NXL-104 against KPC-producing organisms									
Isolate	Resistance Mechanism	CAZ	CAZ:		Ex-1: NXL104	Ex-12	Ex-12: NXL104	Ex-4	Ex-4: NXL104
			CAZ	NXL104					
E. cloacae 2689	KPC-2	>32	2	32	1	32	1	4	0.5
E. cloacae 2705	VIM-1, KPC-3type	>32	>32	>32	>32	>32	>32	>32	>32
E. cloacae 2706	VIM-1, KPC-3	>32	32	>32	32	>32	>32	>32	32
E. cloacae 2709	KPC-3, VIM-1	>32	8	32	4	32	4	8	4
E. cloacae 2710	KPC-3, VIM-1	>32	>32	>32	>32	>32	>32	>32	>32
E. coli 2666	KPC-2	>32	4	32	1	16	1	32	2
E. coli 4103	KPC-2, TEM-1, CMY-type	>32	16	>32	4	>32	4	>32	8
K. pneumoniae 2689	KPC-2	>32	4	16	1	16	1	16	1
K. pneumoniae 2690	KPC-2	>32	4	32	1	8	1	16	2
K. pneumoniae 2711	KPC-3, VIM-1	>32	2	4	0.25	4	0.5	8	1


TABLE 3-continued

MIC values of selected compounds alone and in combination with NXL-104 against KPC-producing organisms									
Isolate	Ex-14	Ex-14: NXL104	Ex-2	Ex-2: NXL104	Ex-5	Ex-5: NXL104	Ex-6	Ex-6: NXL104	
K. pneumoniae 2712	KPC-3, VIM-1	>32	16	>32	4	>32	4	>32	4
K. pneumoniae 2713	KPC-3, VIM-1	>32	8	>32	1	>32	2	>32	2
K. pneumoniae 4088	KPC-3	>32	8	>32	4	>32	2	>32	4
K. pneumoniae 4104	KPC-3, TEM-1	>32	2	32	1	16	1	16	1
K. pneumoniae 4105	KPC-3, TEM-1, SHV-11	>32	4	32	2	32	2	>32	2
K. pneumoniae 4106	KPC-3, TEM-1, SHV-12, SHV-141	>32	8	32	2	32	2	32	4
K. pneumoniae 4107	KPC-2, TEM-1, SHV-11	>32	4	32	1	16	1	32	2
K. pneumoniae 4108	KPC-3, TEM-166, SHV-12, SHV-141	>32	4	32	2	32	1	>32	2
P. aeruginosa 2686	KPC-2	>32	16	>32	8	32	32	32	8
E. cloacae 2689		16	0.5	8	0.5	16	0.5	16	1
E. cloacae 2705		>32	>32	>32	>32	>32	>32	>32	>32
E. cloacae 2706		>32	32	>32	32	>32	32	>32	32
E. cloacae 2709		32	4	16	4	32	4	32	4
E. cloacae 2710		>32	>32	>32	>32	>32	>32	>32	>32
E. coli 2666		32	1	32	2	16	1	16	1
E. coli 4103		>32	8	>32	8	>32	4	>32	4
K. pneumoniae 2689		32	1	32	2	8	1	16	1
K. pneumoniae 2690		32	1	32	2	8	.5	16	1
K. pneumoniae 2711		4	0.5	4	0.5	2	0.25	4	0.5
K. pneumoniae 2712		>32	4	>32	4	>32	4	>32	4
K. pneumoniae 2713		>32	2	>32	2	>32	2	>32	2
K. pneumoniae 4088		>32	4	>32	4	32	2	>32	2
K. pneumoniae 4104		32	1	32	2	16	0.5	32	0.5
K. pneumoniae 4105		>32	2	>32	2	32	1	32	2
K. pneumoniae 4106		>32	4	>32	4	32	2	32	2
K. pneumoniae 4107		16	2	>32	2	16	1	16	1
K. pneumoniae 4108		>32	2	>32	2	32	1	32	1
P. aeruginosa 2686		>32	16	>32	16	32	8	32	8

221


We claim:

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:




wherein in the formula (I),

(i) A is defined by the formula (Ia):



[Where] where X is N, C(H), C(F) or C(Cl);

(ii) B is defined as hydrogen, methyl, ethyl or represented by the formula (Ib)





wherein, R¹ and R² is independently hydrogen or lower alkyl; or

wherein R¹ and R² together may form a 3 to 6-membered spiro ring system; and









m is 0 or 1;

(iii) C represents a quaternized bicyclic nitrogen containing aromatic heterocyclic ring represented by the formulae (Ic) to (Iz)




222

-continued




223

-continued




224

-continued




- (iv) D represents CH₂, CH₂CH₂ or CH₂CO;
- (v) E represents a substituted or unsubstituted benzene ring or a substituted or unsubstituted 5- or 6-membered aromatic heterocyclic ring having at least one heteroatom selected from the group consisting of O, S and N, wherein the heteroaromatic ring [includes] is pyrrolyl, imidazolyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazolyl, isoxazolyl, oxazolyl, oxadiazolyl, isothiazolyl, thiazolyl, thiadiazolyl, furyl, [and] or thienyl;
- (vi) F is optionally substituted amidine or optionally substituted guanidine; and
- (vii) G is hydrogen, methyl, ethyl, C₃₋₆ alkyl, C₃₋₆ cycloalkyl or [a] an optionally substituted 5- or 6-membered aliphatic or [a] an optionally substituted 5- or 6-membered aromatic heterocyclic ring, wherein the heterocyclic ring is substituted with at least 1-2 heteroatoms selected from the group consisting of N, O, and S ([α or β] α or β).

2. A compound as recited in claim 1, where E is selected from [a substituted aryl or a 5- and] the group consisting of an optionally substituted benzene ring and an optionally substituted 5- or 6-membered aromatic heterocyclic [rings] ring, comprising:








225

-continued











3. A compound as recited in claim 2, wherein the optional substituents [include] on the benzene ring and the 5- or 6-membered aromatic heterocyclic rings of E are selected from the group consisting of chloro, fluoro, cyano, hydroxy, amino, carboxy, acetyl, methoxy, ethoxy, trifluoromethyl, pyrrolidinyloxy, and piperidinyloxy.

4. A compound as recited in claim 1, wherein [the preferred examples of] “-C-D-E-F” [include] comprises one of the following organic residues



226








-continued



227

-continued









[



228


-continued

21




229

-continued










230

-continued



231








-continued



232

-continued

49



50

51

52

53

54

55

42

43

44

45

46

47

48

49

50








51

52

53

233


-continued



234

-continued

56



5

10

57

15

20

58

25

30

59

35

40

60

45

61






50

55

62







60

65









235

-continued



236


-continued



65


237

-continued












238

-continued



239

-continued




240

-continued


96

5




97

10




98

15




99

20




100

25




101

30




102

35




103

40




104

45




105

50




106

55



107

60










108

65










241

-continued









242

-continued









243

-continued




244

-continued




245

-continued











246

-continued



247

-continued



248

-continued

154

5

155

10

156

15

157

20

158

25

159

30

160

35

161

40

162

45

163

50

164

55

165

60

166

65

162







163

164

165








166

167










249

-continued




250

-continued




251

-continued









252

-continued









253

-continued




254

-continued




255

-continued










256

-continued











257

-continued








258

-continued



259

-continued



260

-continued

236

5

10

237

15

20

238

25

30

35

239

40

45

50








55

240

60








65

241










261

-continued










262

-continued









263

-continued





264

-continued




265

-continued






266






-continued







20 **5.** The compound of formula (I) as recited in claim 1, [which is] selected from the [following] group [of compounds] consisting of:








-continued




-continued







-continued








-continued









-continued








-continued








-continued








-continued








-continued




-continued







-continued







-continued




-continued








-continued







-continued







-continued








-continued








-continued








-continued








-continued








-continued







-continued








-continued








-continued








-continued




-continued








-continued








-continued








-continued








-continued








-continued








-continued








-continued








-continued



-continued




-continued



343

344


-continued




6. A method of treating a bacterial infection comprising administering to a mammal in need thereof an antibacterially effective amount of a compound as recited in claim 1.

7. A pharmaceutical composition containing as an active ingredient, at least one compound as recited in claim 1 and at least one pharmaceutically acceptable carrier or diluent.


8. A method of treating a bacterial infection comprising administering to a mammal in need thereof a combination of (i) an antibacterially effective amount of a compound as recited in claim 1, and (ii) a therapeutically effective amount of a β -lactamase inhibitor selected from the group consisting of:




(IIa) 30




(IIIa) 35




(IIIb) 40




(IVa) 45




(Va)




(VIa)




(VIb)



(VIc)



RPX 7381




(VIId)


RPX 7400

345






-continued



NXL-104




MK-7655



346

(VIIa)


5



(IIa)

(VIIb)


15



(IIIa)

(VIIc)


25



(IIIb)

(VIId)


30



(IVa)

(VIIe)


40



(Va)

(VIIf)

45




(VIa)

(VIIg)

55

RPX 7262



(VIb)


RPX 7282

9. A pharmaceutical composition containing as an active ingredient at least (i) one compound as recited in claim 1 and (ii) a therapeutically effective amount of a β -lactamase inhibitor selected from the group consisting of:


65

347


-continued




RPX 7381






RPX 7400



(VIIa, NXL-104)



(VIIb, MK-7655)




348

-continued

(VIc)


5



10

(VIId)

15



20

(VIIIf)

(VIIg)


10. The method as recited in claim 8, wherein (i) and (ii) are administered simultaneously, sequentially, or separated in time.

25 11. The pharmaceutical composition as recited in claim 9, wherein the ratio of the weight of (i) to the weight of (ii) is in the range of from about 1:20 to about 20:1.

30 12. A process for preparing a compound of formula (I) as recited in claim 1, the process comprising one of the following processes:


35 Process I wherein the intermediate VIII ($q=0$, $Y=\text{chloride}$) is coupled with intermediate (IX) to provide the intermediate (X) followed by removal of protecting groups to provide the compound of formula (I)

(VIIc) 40




(VIIId)

50




(VIIe) 60



65


349

-continued



or

Process II wherein the intermediate VIII ($q=1$, Y =iodide) is coupled with intermediate (IX) and subsequently reducing the sulfoxide to sulfide providing the intermediate (XI) by removal of the protecting groups provides the compound of formula (I)




wherein in the formula VIII, P^1 is a suitable carboxy protecting group.

13. A compound of formula (I) or a pharmaceutically acceptable salt thereof:

350


(I)



wherein in the formula (I),


(i) A is defined by the formula (Ia)

(Ia)



(ii) B is defined as hydrogen, methyl, ethyl or represented by the formula (Ib):


(Ib)



wherein, R^1 and R^2 are independently hydrogen or lower alkyl; or wherein R^1 and R^2 together may form a 3 to 6 membered spiro ring system; and m is 0 or 1; and


(iii) wherein “-C-D-E-F” is selected from the group consisting of one of the following organic residues:

14




24

14. A compound of formula (I) or a pharmaceutically acceptable salt thereof:



15
 wherein the compound is selected from the group consisting of the following compounds:



* * * * *