



US00RE49423E

(19) **United States**
 (12) **Reissued Patent**
Williams

(10) **Patent Number: US RE49,423 E**
 (45) **Date of Reissued Patent: Feb. 21, 2023**

(54) **DNA PLASMIDS WITH IMPROVED EXPRESSION**

WO 2006029449 A1 3/2006
 WO 2008153733 A2 12/2008
 WO 2009025690 A2 2/2009

(71) Applicant: **NATURE TECHNOLOGY CORPORATION**, Lincoln, NE (US)

OTHER PUBLICATIONS

(72) Inventor: **James A. Williams**, Lincoln, NE (US)

A. Duilio et al.; Molecular Characterization of a Recombinant Replication Protein (Rep) from the Antarctic *Bacterium psychrobacter* sp. TA144; FEMS Microbiology, V 198, p. 49-55.

(73) Assignee: **Nature Technology Corporation**, Lincoln, NE (US)

Nature Technology Corporation: "Targeting RIG-I activating Expression Vectors", Instruction Manual, vol. version 1, Sep. 1, 2011 (Sep. 1, 2011), XP002713690.

(21) Appl. No.: **17/112,918**

Abhyankar et al., Biochemical Investigations of Control of Replication Initiation of Plasmid R6K*, Journal of Biological Chemistry, vol. 279 No. 8, pp. 6711-6719, 2004.

(22) Filed: **Dec. 4, 2020**

Abhyankar et al., "Reconstitution of R6K replication in Vitro Using 22 Purified Proteins", The Journal of Biological Chemistry, vol. 278, No. 46, pp. 45476-45484, 2003.

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **10,144,935**
 Issued: **Dec. 4, 2018**
 Appl. No.: **15/375,215**
 Filed: **Dec. 12, 2016**

U.S. Applications:

(62) Division of application No. 14/422,865, filed as application No. PCT/US2013/000068 on Mar. 14, 2013, now Pat. No. 9,550,998.

Angulo et al., "Identification of a Boundary Domain Adjacent to the Potent Human cytomegalovirus Enhancer That Represses Transcription of the Divergent UL127 Promoter" Journal of Virology, Mar. 2000, p. 2826-2839.

(60) Provisional application No. 61/743,219, filed on Aug. 29, 2012.

Ashfield et al. "MAZ-dependent termination between closely spaced human complement genes" The EMBO Journal vol. 13 No. 23 pp. 5656-5667, 1994.

(51) **Int. Cl.**
C12N 15/63 (2006.01)
C12N 15/85 (2006.01)
C12N 15/70 (2006.01)
A61K 39/00 (2006.01)
A61K 48/00 (2006.01)

Barouch et al. "A Human T-Cell Leukemia Virus Type 1 Regulatory Element Enhances the Immunogenicity of Human Immunodeficiency Virus Type 1 DNA Vaccines in Mice and Nonhuman Primates" Journal of Virology, Jul. 2005, pp. 8828-8834 2005.

(52) **U.S. Cl.**
 CPC **C12N 15/635** (2013.01); **C12N 15/70** (2013.01); **C12N 15/85** (2013.01); **A61K 39/00** (2013.01); **A61K 48/00** (2013.01); **C12N 2800/107** (2013.01); **C12N 2820/55** (2013.01)

Carnes "Fermentation Design for the Manufacture of Therapeutic Plasmid DNA" Bioprocess International, pp. 2-7, Oct. 2005.

(58) **Field of Classification Search**
 CPC **C12N 15/635**; **C12N 15/70**
 See application file for complete search history.

Chenna et al. "Multiple sequence alignment with the Clustal series of programs" Nucleic Acids Research, vol. 31, No. 13, pp. 3497-3500, 2003.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,922,583 A 7/1999 Morsey
 6,977,174 B2 12/2005 Cruzel et al.
 7,244,609 B2 7/2007 Drocourt et al.
 7,279,313 B2 10/2007 Soubrier
 7,611,883 B2 11/2009 Cranenburgh
 7,807,444 B2 10/2010 Soubrier
 7,943,377 B2 5/2011 Carnes et al.
 2003/0161844 A1 8/2003 Soubrier
 2006/0063232 A1 3/2006 Grabherr et al.
 2007/0015282 A1 1/2007 Soubrier
 2007/0141708 A2 6/2007 Soubrier
 2009/0075357 A1 3/2009 Snyder
 2009/0252713 A1 10/2009 Soubrier
 2010/0184158 A1 7/2010 Williams
 2010/0303859 A1 12/2010 Williams

Del Solar et al., "Replication and Control of Circular Bacterial Plasmids", Microbiology and Molecular Biology Reviews, vol. 62, No. 2, pp. 434-464, Jun. 1998.

Franch and Gerdes "U-turns and regulatory RNAs" Current Opinion in Microbiology, 3:159-164, 2000.

(Continued)

Primary Examiner — Bruce R Campell

Assistant Examiner — Dwayne C Jones

(74) *Attorney, Agent, or Firm* — Barnes & Thornburg LLP; Matthew S. Gibson

(57) **ABSTRACT**

The present invention relates to the production and use of covalently closed circular (ccc) recombinant plasmids, and more particularly to vector modifications that improve expression of said DNA molecules in the target organism.

FOREIGN PATENT DOCUMENTS

GB 2214508 A 6/1989
 WO 2004033664 A2 4/2004

14 Claims, 12 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

- Garg et al. "The Hybrid Cytomegalovirus Enhancer/Chicken b-Actin Promoter along with Woodchuck Hepatitis Virus Posttranscriptional Regulatory element Enhances the Protective Efficacy of DNA Vaccines" *The Journal of Immunology* pp. 550-558, 2004.
- Hebel et al. "Challenges in the process development of a novel Zero-CpG CFTR plasmid for human clinical use" VGX Pharmaceuticals, Inc., (1 page), 2008.
- Hiraga et al., "Comparative analysis of the replicon regions of eleven ColE2-related plasmids" *Journal of Bacteriology*, vol. 17, No. 23, pp. 7233-7243, 1994.
- Kim et al. "Improved Mammalian Expression Systems by Manipulating Transcriptional Termination Regions" *Biotechnol. Prog.* vol. 19, No. 5. p. 1620-1622, 2003.
- Lu et al. "The Extragenic Spacer Length Between the 5' and 3' Ends of the Transgene Expression Cassette Affects Transgene Silencing from Plasmid-based Vectors" *Molecular Therapy*, pp. 1-9, 2012.
- Luke et al. "Improved antibiotic-free DNA vaccine vectors utilizing a novel RNA based plasmid selection system" *Vaccine* 27, pp. 6454-6459, 2009.
- Luke et al. "Vector Insert-Targeted Integrative Antisense Expression System for Plasmid Stabilization" *Mol. Biotechnol.*, (7 pages), 2010.
- Manoj et al. "Approaches to Enhance the Efficacy of DNA Vaccines" *Critical Reviews in Clinical Laboratory Sciences*, 41(1), pp. 1-39, 2004.
- Marians et al. "Maximal Limits of the *Escherichia coli* Replication Factor Y Effector Site Sequences in pBR322 DNA" *The Journal of Biological Chemistry*, vol. 247, No. 10, pp. 5656-5662, May 1982.
- Masai et al. "The ABC-Primosome—A Novel Priming System Employing dnaA, dnaB, dnaC, and Primase on a Hairpin Containing a dnaA Box Sequence" *The Journal of Biological Chemistry*, vol. 265, No. 25, pp. 15134-15144, Sep. 1990.
- Masai et al. "Roles of ϕ X174 Type Primosome- and G4 Type Primase-dependent Primings in Initiation of Lagging and Leading Strand Syntheses of DNA replication" *The Journal of Biological Chemistry*, vol. 265, No. 25, pp. 15124-15133, Sep. 1990.
- Mutalik et al. "Rationally designed families of orthogonal RNA regulators of translation" *Nature of Chemical Biology*, pp. 1-8, Mar. 2012.
- Na et al. "Metabolic engineering of *Escherichia coli* using synthetic small regulatory RNAs" *Nature Biotechnology*, pp. 1-7, 2013.
- Nomura et al. "Identification of eleven single-strand initiation sequences (ssi) for priming of DNA replication in the F, R6K, R100 and ColE2 plasmids" *Gene*, 108 (1991) 15-22.
- Sato et al. "A specific DNA Sequence Controls Termination of Transcription in the Gastrin Gene" *Molecular and Cellular Biology*, vol. 6, No. 4, pp. 1032-1043, Apr. 1986.
- Soubrier et al. "pCOR: a new design of plasmids vectors for nonviral gene therapy" *Gene Therapy*, (1999)6, pp. 1482-1489.
- Takechi et al. "Control of ColE2 plasmid replication: regulation of Rep expression by a plasmid-coded antisense RNA", *Mol Gen Genet* (1994) 244: pp. 49-56.
- Wagner et al. "Antisense RNAs in Bacteria and Their Genetic Elements" *Advances in Genetics*, vol. 46, pp. 361-398, 2002.
- Wilson et al. "Importance of Structural Differences between Complementary RNA Molecules to Control of Replication of and IncB Plasmid" *Journal of Bacteriology*, vol. 179, No. 3, pp. 742-753, Feb. 1997.
- Wu et al. "A DNA Segment conferring Stable Maintenance on R6K γ -Origin Core Replicons" *Journal of Bacteriology*, vol. 177, No. 22, pp. 6338-6345, Nov. 1995.
- Yagura et al. "The Rep protein binding elements of the plasmid ColE2-P9 replication origin" *Biochemical and Biophysical Research Communications*, vol. 345, pp. 872-877, 2006.
- Yagura et al. "Anatomy of the Replication Origin of Plasmid ColE2-P9" *Journal of Bacteriology*, vol. 188, No. 3, pp. 999-1010, Feb. 2006.
- Bailone et al. "Nucleotide sequence of the operators of I ultravirulent mutants" *Nucleic Acids Research* vol. 8, No. 10, 1980.
- Datsenko et al. "One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products" *PNAS*, vol. 97, No. 12, pp. 6640-6645, 2000.
- Nagase et al. "Essential elements in the coding region of mRNA for translation of ColE2 Rep Protein" *Plasmid* 59, pp. 36-44, 2008.
- Haldimann et al. "Conditional-Replication, Integration, Excision, and Retrieval Plasmid-Host Systems for Gene Structure-Function Studies of Bacteria" *Journal of Bacteriology*, vol. 183, No. 21, pp. 6384-6393, 2001.
- Han et al. "The ColE2-P9 Rep protein binds to the origin DNA as a monomer" *Biochemical and Biophysical Research Communications* 353, pp. 306-310, 2007.
- Kay et al. "A robust system for production of minicircle DNA vectors" *Nature Biotechnology* 28, pp. 1287-1289, 2010.
- Lieb M. "Heat-Sensitive Lambda Repressors Retain Partial Activity During Bacteriophage Induction" *Journal of Virology*, vol. 32, No. 1, pp. 162-166, 1979.
- Lanza et al. "Global Strain Engineering by Mutant Transcription Factors" *Methods in Molecular Biology*, vol. 765, pp. 253-274, 2011.
- Lissemore et al. "Green Fluorescent Protein as a Quantitative Reporter of Relative Promoter Activity in *E. coli*" *Biotechniques*, vol. 28, No. 1, 2000.
- Luke et al. "Coexpressed RIG-1 Agonist Enhances Humoral Immune Response to Influenza Virus DNA Vaccine" *Journal of Virology*, vol. 85, No. 3, pp. 1370-1383, 2011.
- Metcalf et al. "Use of the rep technique for allele replacement to construct new *Escherichia coli* hosts for maintenance of R6K γ origin plasmids at different copy numbers" *Gene Therapy* vol. 138, pp. 1-7, 1994.
- Wang et al. "Construction and analysis of compact muscle-specific promoters for AAV vectors" *Gene Therapy* 15, pp. 1489-1499, 2008.
- Yasueda et al. "Structural and functional organization of ColE2 and ColE3 replicons" *Mol Gen Genet* 215, pp. 209-216, 1989.
- Yasueda et al. "Control of ColE2 plasmid replication: negative regulation of the expression of the plasmid-specified initiator protein, Rep, at a posttranscriptional step" *Mol Gen Genet* 244, pp. 41-48, 1994.
- Zhao et al. "Transfection of shRNA-encoding Minivector DNA of a few hundred base pairs to regulate gene expression in lymphoma cells" *Gene Therapy* 18, pp. 220-224, 2011.
- Benson et al., "DNA Sequence Determinants of λ Repressor Binding in Vivo", *Genetics*, vol. 188, Jan. 1988, pp. 21-29.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6





FIGURE 7



SEQ ID # 10 - WILD TYPE
 11 - OL1-G
 12 - OL1-G to T

FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11





FIGURE 12



B)



DNA PLASMIDS WITH IMPROVED EXPRESSION

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue; a claim printed with strikethrough indicates that the claim was canceled, disclaimed, or held invalid by a prior post-patent action or proceeding.

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a division of U.S. application Ser. No. 14/422,865, filed Feb. 20, 2015, which is a 371 U.S. National Phase application of PCT/US2013/000068, filed Mar. 14, 2013 which claim the benefit of U.S. Provisional Application Ser. No. 61/743,219, filed Aug. 29, 2012 entitled "DNA Plasmids With Improved Expression". The entire disclosures of each of the above-identified applications are incorporated herein by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

This invention was supported in part with government support under Grant No. R44GM080768, awarded by the National Institutes of Health. The government has certain rights in this invention.

Applicants submit herewith a computer readable form (CRF) of the Sequence Listing for entry into this application. The content of the Sequence Listing information recorded in computer readable form includes no new matter beyond the scope of the published International Application.

FIELD OF THE INVENTION

The present invention relates to a family of eukaryotic expression plasmids useful for gene therapy, obtaining improved genetic immunization, natural interferon production, and more particularly, for improving the expression of plasmid encoded antigens, therapeutic proteins and RNAs.

The present invention also relates to the production of covalently closed circular (ccc) recombinant DNA molecules such as plasmids, cosmids, bacterial artificial chromosomes (BACs), bacteriophages, viral vectors and hybrids thereof, and more particularly to strain modifications that improve production yield of said DNA molecules in fermentation culture.

Such recombinant DNA molecules are useful in biotechnology, transgenic organisms, gene therapy, therapeutic vaccination, agriculture and DNA vaccines.

BACKGROUND OF THE INVENTION

E. coli plasmids have long been an important source of recombinant DNA molecules used by researchers and by industry. Today, plasmid DNA is becoming increasingly important as the next generation of biotechnology products (e.g., gene medicines and DNA vaccines) make their way into clinical trials, and eventually into the pharmaceutical marketplace. Plasmid DNA vaccines may find application as preventive vaccines for viral, bacterial, or parasitic diseases; immunizing agents for the preparation of hyper immune globulin products; therapeutic vaccines for infectious diseases; or as cancer vaccines. Plasmids are also utilized in

gene therapy or gene replacement applications, wherein the desired gene product is expressed from the plasmid after administration to the patient.

Therapeutic plasmids often contain a pMB1, ColE1 or pBR322 derived replication origin. Common high copy number derivatives have mutations affecting copy number regulation, such as ROP (Repressor of primer gene) deletion, with a second site mutation that increases copy number (e.g. pMB1 pUC G to A point mutation, or ColE1 pMM1). Higher temperature (42° C.) can be employed to induce selective plasmid amplification with pUC and pMM1 replication origins.

U.S. Pat. No. 7,943,377 (Carnes, A E and Williams, J A, 2011) disclose methods for fed-batch fermentation, in which plasmid-containing *E. coli* cells were grown at a reduced temperature during part of the fed-batch phase, during which growth rate was restricted, followed by a temperature upshift and continued growth at elevated temperature in order to accumulate plasmid; the temperature shift at restricted growth rate improved yield and purity of plasmid. Other fermentation processes for plasmid production are described in Carnes A. E. 2005 *BioProcess Intl*; 3:36-44, which is incorporated herein by reference in its entirety.

The art teaches that one of the limitations of application of plasmid therapies and plasmid vaccines is regulatory agency (e.g. Food and Drug Administration, EMEC) safety concerns regarding 1) plasmid transfer and replication in endogenous bacterial flora, or 2) plasmid encoded selection marker expression in human cells, or endogenous bacterial flora. Additionally, regulatory agency guidances recommend removal of all non essential sequences in a vector. Plasmids containing a pMB1, ColE1 or pBR322 derived replication origin can replicate promiscuously in *E. coli* hosts. This presents a safety concern that a plasmid therapeutic gene or antigen will be transferred and replicated to a patient's endogenous flora. Ideally, a therapeutic or vaccine plasmid would be replication incompetent in endogenous *E. coli* strains. This requires replacement of the pMB1, ColE1 or pBR322 derived replication origin with a conditional replication origin that requires a specialized cell line for propagation. As well, regulatory agencies such as the EMEA and FDA are concerned with utilization of antibiotic resistance or alternative protein markers in gene therapy and gene vaccine vectors, due to concerns that the gene (antibiotic resistance marker or protein marker) may be expressed in a patient's cells. Ideally, plasmid therapies and plasmid vaccines would be 1) replication incompetent in endogenous *E. coli* strains, 2) would not encode a protein based selection marker and 3) be minimalized to eliminate all non essential sequences.

The art further teaches that one of the limitations of application of plasmid therapies and vaccines is that antigen expression is generally very low. Vector modifications that improve antigen expression (e.g. codon optimization of the gene, inclusion of an intron, use of the strong constitutive CMV or CAGG promoters versus weaker or cell line specific promoter) are highly correlative with improved in vivo expression and, where applicable, immune responses (reviewed in Manoj S, Babiuk L A, van Drunen Little-van den Hurk S. 2004 *Crit Rev Clin Lab Sci* 41: 1-39). A hybrid CMV promoter (CMV/R), which increased antigen expression, also improved cellular immune responses to HIV DNA vaccines in mice and nonhuman primates (Barouch D H, Yang Z Y, Kong W P, Koriath-Schmitz B, Sumida S M, Truitt D M, Kishko M G, Arthur J C, Miura A, Mascola J R, Letvin N L, Nabel G J. 2005 *J Virol*. 79: 8828-8834). A plasmid containing the woodchuck hepatitis virus posttran-

scriptional regulatory element (a 600 bp element that increases stability and extranuclear transport of RNA resulting in enhanced levels of mRNA for translation) enhanced antigen expression and protective immunity to influenza hemagglutinin (HA) in mice (Garg S, Oran A E, Hon H, Jacob J. 2004 J Immunol. 173: 550-558). These studies teach that improvement in expression beyond that of current CMV based vectors may generally improve immunogenicity and, in the case of gene therapeutics, efficacy.

SUMMARY OF THE INVENTION

The present invention relates to a family of minimalized eukaryotic expression plasmids that are replication incompetent in endogenous flora and have dramatically improved in vivo expression. These vectors are useful for gene therapy, genetic immunization and or interferon therapy.

The present invention also relates generally to methods of increasing production yield of covalently closed supercoiled plasmid DNA.

Improved vectors that utilize novel replication origins that unexpectedly improve antigen expression are disclosed.

One object of the invention is to provide improved expression plasmid vectors. Yet another object of the invention is to provide methods for improving plasmid copy number.

According to one object of the invention, a method of increasing expression from an expression plasmid vector comprises modifying the plasmid DNA to replace the pMB1, ColE1 or pBR322 derived replication origin and selectable marker with an alternative replication origin selected from the group consisting of an minimal pUC origin, a R6K gamma replication origin, a ColE2-P9 replication origin, and a ColE2-P9 related replication origin and an RNA selectable marker; transforming the modified plasmid DNA into a bacterial cell line rendered competent for transformation; and isolating the resultant transformed bacterial cells. The resultant plasmid surprisingly has higher in vivo expression levels than the parent pMB1, ColE1 or pBR322 derived replication origin expression plasmid vector.

According to one object of the invention, a composition for construction of a eukaryotic expression vector comprises an R6K origin with at least 90% sequence identity to the sequence set forth as SEQ ID NO: 1, and a RNA selectable marker, wherein said R6K origin is operably linked to said RNA selectable marker and a eukaryotic region. According to another object of the invention, said R6K origin-RNA selectable marker improves said vector expression in vivo compared to a corresponding vector containing a pMB1, ColE1 or pBR322 derived replication origin. According to still another object of the invention, said vector has at least 95% sequence identity to a sequence selected from the group consisting of: SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO:41.

According to one object of the invention, a composition for construction of a eukaryotic expression vector comprises a ColE2-P9 origin with at least 90% sequence identity to the sequence set forth as SEQ ID NO: 4, 5, 6, or 7, and a RNA selectable marker, wherein said ColE2-P9 origin—a RNA selectable marker is operably linked to a eukaryotic region. According to another object of the invention, said ColE2-P9 origin-RNA selectable marker improves said vector expression in vivo compared to a corresponding vector containing a pMB1, ColE1 or pBR322 derived replication origin. According to still another object of the invention, said vector has at least 95% sequence identity to a sequence selected

from the group consisting of: SEQ ID NO: 8, SEQ ID NO: 9. According to still another object of the invention, a primosomal assembly site (ssiA) is optionally incorporated into the vector adjacent to the ColE2-P9 origin.

According to another object of the invention, production cell lines are disclosed that improve plasmid yield in shake flask and or fermentation culture with said R6K gamma replication origin, ColE2-P9 replication origin, or ColE2-P9 related replication origin plasmid vectors of the current invention.

According to another object of the invention, production cell lines providing heat inducible induction of R6K gamma replication origin, ColE2-P9 replication origin, or ColE2-P9 related replication origin plasmid vectors for DNA production are disclosed. These cell lines contain one or more copies of the corresponding R6K gamma replication origin, ColE2-P9 replication origin, or ColE2-P9 related replication protein integrated into the genome and expressed from the group consisting of: the heat inducible P_L promoter (SEQ ID NO: 10), the heat inducible P_L promoter incorporating the OL1-G deletion (SEQ ID NO: 11), the heat inducible P_L promoter incorporating the OL1-G to T substitution (SEQ ID NO: 12).

According to another object of the invention, mutant R6K replication proteins that improve heat inducible induction of R6K gamma replication origin vectors are disclosed. These cell lines contain one or more copies of the mutant R6K gamma replication origin replication protein integrated into the genome and expressed from the group consisting of: the heat inducible P_L promoter (SEQ ID NO: 10), the heat inducible P_L promoter incorporating the OL1-G deletion (SEQ ID NO: 11), the heat inducible P_L promoter incorporating the OL1-G to T substitution (SEQ ID NO: 12). The mutant R6K gamma replication origin replication protein are selected from the group consisting of: P42L-P113S (SEQ ID NO: 13), P42L-P106L-F107S (SEQ ID NO: 14).

According to another object of the invention, a mutant ColE2-P9 replication protein that improve heat inducible induction of ColE2-P9 replication origin vectors is disclosed. These cell lines contain one or more copies of the mutant ColE2-P9 replication origin replication protein integrated into the genome and expressed from the group consisting of: the heat inducible P_L promoter (SEQ ID NO: 10), the heat inducible P_L promoter incorporating the OL1-G deletion (SEQ ID NO: 11), the heat inducible P_L promoter incorporating the OL1-G to T substitution (SEQ ID NO: 12). The mutant ColE2-P9 replication origin replication protein is ColE2-P9 Rep mut G194D (SEQ ID NO: 16).

Further objects and advantages of the invention will become apparent from a consideration of the drawings and ensuing description.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 depicts the NTC8685 pUC origin expression vector;

FIG. 2 depicts the NTC9685C ColE2 origin expression vector;

FIG. 3 depicts the NTC9685R R6K origin expression vector;

FIG. 4 depicts the NTC9385C ColE2 origin expression vector;

FIG. 5 depicts the NTC9385R R6K origin expression vector;

FIG. 6 depicts the pINT pR pL R6K Rep pi P42L-P106L-F107S (P3-) integration vector;

5

FIG. 7 depicts SEQ ID NO: 44, which includes the P_L promoter with OL1 mutations OL1-G and OL1-G to T;

FIG. 8 depicts the pINT pR pL ColE2 Rep protein integration vector;

FIG. 9 shows Nanoplasmid expression in vitro after lipofectamine transfection of HEK293 cell line;

FIG. 10 shows Nanoplasmid expression in vivo after intramuscular injection with EP;

FIG. 11 depicts a ColE2 origin Nanoplasmid shRNA expression vector; and

FIG. 12 depicts an IncB RNAI based RNA selectable marker.

Table 1: P_L promoter with OL1 mutations OL1-G and OL1-G to T improve plasmid yields in HyperGRO fermentation

Table 2: NTC9385R-EGFP LB media shake flask production yields in R6K production strains

Table 3: NTC9385C-Luc plasmid performance in different processes and production cell lines

Table 4: ColE2 Origin EGFP vector production in NTC701131 ColE2 production cell line

Table 5: NTC9382C, NTC9385C, NTC9382R, NTC9385R, NTC9682C, NTC9685C, NTC9682R, and NTC9685R vectors

Table 6: gWIZ and NTC9385C Nanoplasmid expression compared to NTC8685

Table 7: SR vector expression in vitro and in vivo

Table 8: RNA Pol III Nanoplasmid vector expression

Table 9: High level expression is obtained with pMB1 RNAI or RNA-OUT antisense RNA vectors

SEQ ID NO:1: R6K gamma Origin

SEQ ID NO:2: NTC9385R vector backbone

SEQ ID NO:3: NTC9685R vector backbone

SEQ ID NO:4: ColE2 Origin (+7)

SEQ ID NO:5: ColE2 Origin (+7, CpG free)

SEQ ID NO:6: ColE2 Origin (Min)

SEQ ID NO:7: ColE2 Origin (+16)

SEQ ID NO:8: NTC9385C vector backbone

SEQ ID NO:9: NTC9685C vector backbone

SEQ ID NO:10: P_L Promoter (-35 to -10)

SEQ ID NO:11: P_L Promoter OL1-G (-35 to -10)

SEQ ID NO:12: P_L Promoter OL1-G to T(-35 to -10)

SEQ ID NO:13: R6K Rep protein P42L-P113S

SEQ ID NO:14: R6K Rep protein P42L-P106L-F107S

SEQ ID NO:15: ColE2 Rep protein (wild type)

SEQ ID NO:16: ColE2 Rep protein mut (G194D)

SEQ ID NO:17: pINT pR pL R6K Rep piP42L-P106L-F107S (P3-)

SEQ ID NO:18: pINT pR pL ColE2 Rep protein mut (G194D)

SEQ ID NO:19: NTC9385R and NTC9685R Bacterial region. [NheI site-trpA terminator-R6K Origin-RNA-OUT-KpnI site]

SEQ ID NO:20: NTC9385C and NTC9685C Bacterial region. [NheI site-ssiA-ColE2 Origin (+7)-RNA-OUT-KpnI site]

SEQ ID NO:21: NTC9385C and NTC9685C CpG free ssiA [from plasmid R6K]

SEQ ID NO:22: CpG free R6K origin

SEQ ID NO:23: RNA-OUT selectable marker from NTC9385C, NTC9685C, NTC9385R, and NTC9685R

SEQ ID NO:24: RNA-OUT Sense strand RNA from NTC9385C, NTC9685C, NTC9385R, NTC9685R, and NTC9385Ra

SEQ ID NO:25: TPA secretion sequence

SEQ ID NO:26: PCR primer 15061101

SEQ ID NO:27: PCR primer 15061102

6

SEQ ID NO:28: ColE2 core replication origin

SEQ ID NO:29: +7(CpG free)-ssiA ColE2 origin

SEQ ID NO:30: HTLV-IR-Rabbit β globin hybrid intron

SEQ ID NO:31: pMB1 RNAI antisense repressor RNA (origin antisense partner of RNAI)

SEQ ID NO:32: pMB1 RNAI selectable Marker, RNAI RNA (Sense strand)

SEQ ID NO:33: IncB RNAI antisense repressor RNA (IncB plasmid origin RNAI antisense partner)

SEQ ID NO:34: IncB RNAI selectable Marker. DraIII-KpnI restriction fragment

SEQ ID NO:35: IncB RNAI-SacB. PstI-MamI restriction fragment

SEQ ID NO:36: CpG free RNA-OUT selection marker—flanked by KpnI and BglII-EcoRI sites

SEQ ID NO:37: CpG free R6K gamma—RNA-OUT bacterial region (CpG free R6K origin-CpG free RNA-OUT selection marker)—flanked by EcoRI-SphI and BglII-EcoRI sites

SEQ ID NO:38: CpG free ColE2 bacterial region (CpG free ssiA-CpG free ColE2 origin-CpG free RNA-OUT selection marker)-flanked by EcoRI-SphI and BglII-EcoRI sites

SEQ ID NO:39: NTC9385Ra-O2 vector backbone

SEQ ID NO:40: NTC9385Ra-O1 vector backbone

SEQ ID NO:41: NTC9385R-BE vector backbone

SEQ ID NO:42: P_{min} minimal pUC replication origin

SEQ ID NO:43: pUC (0.85) Bacterial region [NheI site—trpA terminator- P_{min} pUC replication origin (minimal)-RNA-OUT-KpnI site]

SEQ ID NO:44: artificial sequence including a P_L promoter

Definition of Terms

A_{405} : Absorbance at 405 nanometers

35 AF: Antibiotic-free

APC: Antigen Processing Cell, for example, langerhans cells, plasmacytoid or conventional dendritic cells

Approximately: As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is the same or similar to a stated reference value

40 BAC: Bacterial artificial chromosome

Bacterial region: Region of a plasmid vector required for propagation and selection in the bacterial host

45 BE: Boundary element: Eukaryotic sequence that that blocks the interaction between enhancers and promoters. Also referred to as insulator element. An example is the AT-rich unique region upstream of the CMV enhancer SpeI site that can function as an insulator/boundary element (Angulo A, Kerry D, Huang H, Borst E M, Razinsky A, Wu J et al. 2000 J Virol 74: 2826-2839)

bp: basepairs

ccc: Covalently Closed Circular

cI: Lambda repressor

55 cITs857: Lambda repressor further incorporating a C to T (Ala to Thr) mutation that confers temperature sensitivity. cITs857 is a functional repressor at 28-30° C., but is mostly inactive at 37-42° C. Also called cI857

Cm^R : Chloramphenicol resistance

cmv: Cytomegalovirus

CMV promoter boundary element: AT-rich region of the human cytomegalovirus (CMV) genome between the UL127 open reading frame and the major immediate-early (MIE) enhancer. Also referred to as unique region (Angulo et al. Supra, 2000)

65 ColE2-P9 replication origin: a region which is specifically recognized by the plasmid-specified Rep protein to initiate

DNA replication. Includes but not limited to ColE2-P9 replication origin sequences disclosed in SEQ ID NO:4: ColE2 Origin (+7), SEQ ID NO:5: ColE2 Origin (+7, CpG free), SEQ ID NO:6: ColE2 Origin (Min) and SEQ ID NO:7: ColE2 Origin (+16) and replication functional mutations as disclosed in Yagura et al 2006, J Bacteriol 188:999-1010 included herein by reference

ColE2 related replication origin: The ColE2-P9 origin is highly conserved across the ColE2-related plasmid family. Fifteen ColE2 related plasmid members including ColE3 are compared in Hiraga et al 1994, J Bacteria 176:7233 and 53 ColE2 related plasmid members including ColE3 are compared in Yagura et al Supra, 2006. These sequences are included herein by reference

ColE2-P9 plasmid: a circular duplex DNA molecule of about 7 kb that is maintained at about 10 to 15 copies per host chromosome. The plasmid encodes an initiator protein (Rep protein), which is the only plasmid-specified transacting factor essential for ColE2-P9 plasmid replication

ColE2-P9 replication origin RNA-OUT bacterial region: Contains a ColE2-P9 replication origin for propagation and the RNA-OUT selection marker. Optionally includes a PAS, for example, the R6K plasmid CpG free ssiA primosomal assembly site (SEQ ID NO:21) or alternative \emptyset X174 type or ABC type primosomal assembly sites, such as those disclosed in Nomura et al 1991 Gene 108:15

ColE2 plasmid: NTC9385C and NTC9685C vectors disclosed herein, as well as modifications and alternative vectors containing a ColE2-P9 replication origin

delivery methods: Methods to deliver gene vectors [e.g. poly(lactide-co-glycolide) (PLGA), ISCOMs, liposomes, niosomes, virosomes, chitosan, and other biodegradable polymers, electroporation, piezoelectric permeabilization, sonoporation, ultrasound, corona plasma, plasma facilitated delivery, tissue tolerable plasma, laser microporation, shock wave energy, magnetic fields, contactless magneto-permeabilization, gene gun, microneedles, naked DNA injection, hydrodynamic delivery, high pressure tail vein injection, needle free biojector, liposomes, microparticles, microspheres, nanoparticles, virosomes, bacterial ghosts, bacteria, attenuated bacteria, etc] as known in the art and included herein by reference

DNA replicon: A genetic element that can replicate under its own control; examples include plasmids, cosmids, bacterial artificial chromosomes (BACs), bacteriophages, viral vectors and hybrids thereof

E. coli: Escherichia coli, a gram negative bacteria

EGFP: Enhanced green fluorescent protein

EP: Electroporation

Eukaryotic expression vector: A vector for expression of mRNA, protein antigens, protein therapeutics, shRNA, RNA or microRNA genes in a target organism

Eukaryotic region: The region of a plasmid that encodes eukaryotic sequences and/or sequences required for plasmid function in the target organism. This includes the region of a plasmid vector required for expression of one or more transgenes in the target organism including RNA Pol II enhancers, promoters, transgenes and polyA sequences. A eukaryotic region may express protein or RNA genes using one or more RNA Pol II promoters, or express RNA genes using one or more RNA Pol III promoters or encode both RNA Pol II and RNA Pol III expressed genes. Additional functional eukaryotic region sequences include RNA Pol I or RNA Pol III promoters, RNA Pol I or RNA Pol III expressed transgenes or RNAs, transcriptional terminators, S/MARs, boundary elements, etc

FU: Fluorescence units

g: Gram, kg for kilogram

Hr(s): Hour(s)

HTLV-I R: HTLV-I R 5' untranslated region (UTR).

Sequences and compositions were disclosed in Williams, J A 2008 World Patent Application WO2008153733 and included herein by reference

IM: Intramuscular

immune response: Antigen reactive cellular (e.g. antigen reactive T cells) or antibody (e.g. antigen reactive IgG) responses

IncB RNAI: plasmid pMU720 origin encoded RNAI (SEQ ID NO: 33) that represses RNA II regulated targets (Wilson I W, Siemering K R, Praszkiar J, Pittard A J. 1997. J Bacteriol 179:742)

kan: Kanamycin

kanR: Kanamycin Resistance gene

Kd: Kilodalton

kozak sequence: Optimized sequence of consensus DNA sequence gccRccATG (R=G or A) immediately upstream of an ATG start codon that ensures efficient translation initiation.

A Sall site (GTCGAC) immediately upstream of the ATG start codon (GTCGACATG) is an effective Kozak sequence

Minicircle: Covalently closed circular plasmid derivatives in which the bacterial region has been removed from the parent plasmid by in vivo or in vitro intramolecular (cis-) site specific recombination or in vitro restriction digestion/ligation

mSEAP: Murine secreted alkaline phosphatase

Nanoplasmid vector: Vector combining an RNA selection marker with a R6K or ColE2 related replication origin. For example, NTC9385C, NTC9685C, NTC9385R, NTC9685R, NTC9385R-BE, NTC9385Ra-O1 and NTC9385Ra-O2 vectors described herein and modifications thereof

NTC7382 promoter: A chimeric promoter comprising the CMV enhancer-CMV promoter-HTLV R-synthetic rabbit β globin 3' intron acceptor-exon 2-SRF protein binding site-kozak sequence, with or without an upstream SV40 enhancer. The creation and application of this chimeric promoter is disclosed in Williams J A Supra, 2008 and included herein by reference

NTC8385: NTC8385 and NTC8685 plasmids are antibiotic-free vectors that contain a short RNA (RNA-OUT) selection marker in place of the antibiotic resistance marker (kanR) The creation and application of these RNA-OUT based antibiotic-free vectors are disclosed in Williams, J A Supra, 2008 and included herein by reference

NTC8685: NTC8685 (FIG. 1) is an antibiotic-free vector that contains a short RNA (RNA-OUT) selection marker in place of the antibiotic resistance marker (kanR) The creation and application of NTC8685 is disclosed in Williams, J A 2010 US Patent Application 20100184158 and included herein by reference

OL1: Lambda repressor binding site in the P_L promoter (FIG. 7). Repressor binding to OL1 is altered by mutations in OL1, such as OL1-G (FIG. 7; this is a single base deletion that also reduces the distance between the P_L promoter -35 and -10 boxes from optimal 17 bp to 16 bp) and OL1-G to T (FIG. 7; this is a G to T substitution that maintains the distance between the P_L promoter -35 and -10 boxes at the optimal 17 bp; this is the V2 mutation described by Bailone A and Galibert F, 1980. Nucleic Acids Research 8:2147)

OD₆₀₀: optical density at 600 nm

PAS: Primosomal assembly site. Priming of DNA synthesis on a single stranded DNA ssi site. \emptyset X174 type PAS: DNA hairpin sequence that binds priA, which, in turn, recruits the remaining proteins to form the preprimosome [priB, dnaT,

recruits dnaB (delivered by dnaC)], which then also recruits primase (dnaG), which then, finally, makes a short RNA substrate for DNA polymerase I. ABC type PAS: DNA hairpin binds dnaA, recruits dnaB (delivered by dnaC) which then also recruits primase (dnaG), which then, finally, makes a short RNA substrate for DNA polymerase I. See Masai et al, 1990 J Biol Chem 265:15134. For example, the R6K plasmid CpG free ssiA primosomal assembly site (SEQ ID NO:21) or alternative \emptyset X174 type or ABC type primosomal assembly sites, such as those disclosed in Nomura et al Supra, 1991

PAS-BH: Primosomal assembly site on the heavy (leading) strand

PAS-BH region: pBR322 origin region between ROP and PAS-BL (approximately pBR322 2067-2351)

PAS-BL: Primosomal assembly site on the light (lagging) strand

PBS: Phosphate buffered Saline

PCR: Polymerase Chain Reaction

pDNA: Plasmid DNA

pINT pR pL vector: The pINT pR pL integration expression vector is disclosed in Luke et al 2011 Mol Biotechnol 47:43 and included herein by reference. The target gene to be expressed is cloned downstream of the pL1 promoter (FIG. 7). The vector encodes the temperature inducible cI857 repressor, allowing heat inducible target gene expression.

P_L promoter: Lambda promoter left (FIG. 7). P_L is a strong promoter that is repressed by the cI repressor binding to OL1, OL2 and OL3 repressor binding sites. The temperature sensitive cI857 repressor allows control of gene expression by heat induction since at 30° C. the cI857 repressor is functional and it represses gene expression, but at 37-42° C. the repressor is inactivated so expression of the gene ensues
Plasmid: An extra chromosomal DNA molecule separate from the chromosomal DNA which is capable of replicating independently from the chromosomal DNA

pMB1 RNAI: pMB1 plasmid origin encoded RNAI that represses RNAII regulated targets (SEQ ID NO: 31; SEQ ID NO:32) that represses RNAII regulated targets such as those described in Grabherr R, Pfaffenzeller I. 2006 US patent application US20060063232 and Cranenburgh R M. 2009; U.S. Pat. No. 7,611,883

P_{min} : Minimal 678 bp pUC replication origin SEQ ID NO:42 and functional variants with base substitutions and/or base deletions. Vectors described herein incorporating P_{min} include NTC8385-Min and NTC8885MP-U6

Pol: Polymerase

polyA: Polyadenylation signal or site. Polyadenylation is the addition of a poly(A) tail to an RNA molecule. The polyadenylation signal is the sequence motif recognized by the RNA cleavage complex. Most human polyadenylation sites contain an AAUAAA motif and conserved sequences 5' and 3' to it. Commonly utilized polyA sites are derived from the rabbit β globin (NTC8685; FIG. 1), bovine growth hormone (gWIZ; pVAX1), SV40 early, or 5V40 late polyA signals
pUC replication origin: pBR322-derived replication origin, with G to A transition that increases copy number at elevated temperature and deletion of the ROP negative regulator

pUC plasmid: Plasmid containing the pUC origin

R6K plasmid: NTC9385R, NTC9685R, NTC9385Ra-O1 and RNA9385Ra-O2 vectors disclosed herein, as well as modifications, and alternative R6K vectors known in the art including but not limited to pCOR vectors (Gencell), pCpG-free vectors (Invivogen), and CpG free University of Oxford vectors including pGM169

R6K replication origin: a region which is specifically recognized by the plasmid-specified Rep protein to initiate

DNA replication. Includes but not limited to R6K replication origin sequence disclosed as SEQ ID NO:1: R6K Origin, and CpG free versions (SEQ ID NO:22) as disclosed in Drocourt et al U.S. Pat. No. 7,244,609 and incorporated herein by reference

R6K replication origin-RNA-OUT bacterial region: Contains a R6K replication origin for propagation and the RNA-OUT selection marker

Rep: Replication

Rep protein dependent plasmid: A plasmid in which replication is dependent on a replication (Rep) protein provided in Trans. For example, R6K replication origin, ColE2-P9 replication origin and ColE2 related replication origin plasmids in which the Rep protein is expressed from the host strain genome. Numerous additional Rep protein dependent plasmids are known in the art, many of which are summarized in del Solar et al 1998 Microbiol. Mol. Biol. Rev 62:434-464 which is included herein by reference

RNA-IN: Insertion sequence 10 (IS10) encoded RNA-IN, an RNA complementary and antisense to RNA-OUT. When RNA-IN is cloned in the untranslated leader of a mRNA, annealing of RNA-IN to RNA-OUT reduces translation of the gene encoded downstream of RNA-IN

RNA-IN regulated selection marker: A genomically expressed RNA-IN regulated selectable marker. In the presence of plasmid borne RNA-OUT, expression of a protein encoded downstream of RNA-IN is repressed. An RNA-IN regulated selection marker is configured such that RNA-IN regulates either 1) a protein that is lethal or toxic to said cell per se or by generating a toxic substance (e.g. SacB), or 2) a repressor protein that is lethal or toxic to said bacterial cell by repressing the transcription of a gene that is essential for growth of said cell (e.g. murA essential gene regulated by RNA-IN tetR repressor gene). For example, genomically expressed RNA-IN-SacB cell lines for RNA-OUT plasmid propagation are disclosed in Williams, J A Supra, 2008 (SEQ ID NO:23) and included herein by reference. Alternative selection markers described in the art may be substituted for SacB

RNA-OUT: Insertion sequence 10 (IS10) encoded RNA-OUT (SEQ ID NO:24), an antisense RNA that hybridizes to, and reduces translation of, the transposon gene expressed downstream of RNA-IN. The sequence of the core RNA-OUT sequence (SEQ ID NO:24) and complementary RNA-IN SacB genomically expressed RNA-IN-SacB cell lines can be modified to incorporate alternative functional RNA-IN/RNA-OUT binding pairs such as those disclosed in Mutalik et al. 2012 Nat Chem Biol 8:447, including, but not limited to, the RNA-OUT A08/RNA-IN S49 pair, the RNA45 OUT A08/RNA-IN S08 pair, and CpG free modifications of RNA-OUT A08 that modify the CG in the RNA-OUT 5' TT CGCT sequence to a non-CpG sequence

RNA-OUT Selectable marker: An RNA-OUT selectable marker DNA fragment including E. coli transcription promoter and terminator sequences flanking an RNA-OUT RNA. An RNA-OUT selectable marker, utilizing the RNA-OUT promoter and terminator sequences, that is flanked by DraIII and KpnI restriction enzyme sites, and designer genomically expressed RNA-IN-SacB cell lines for RNA-OUT plasmid propagation, are disclosed in Williams, J A Supra, 2008 (SEQ ID NO:23) and included herein by reference. The RNA-OUT promoter and terminator sequences flanking the RNA-OUT RNA may be replaced with heterologous promoter and terminator sequences. For example, the RNA-OUT promoter may be substituted with a CpG free promoter known in the art, for example the

I-EC2K promoter or the P5/6 5/6 or P5/6 6/6 promoters disclosed in Williams, J A Supra, 2008 and included herein by reference

RNA selectable marker: also RNA selection marker. An RNA selectable marker is a plasmid borne expressed non translated RNA that regulates a chromosomally expressed target gene to afford selection. This may be a plasmid borne nonsense suppressing tRNA that regulates a nonsense suppressible selectable chromosomal target as described by Crouzet J and Soubrier F 2005 U.S. Pat. No. 6,977,174 included herein by reference. This may also be a plasmid borne antisense repressor RNA, a non limiting list included herein by reference includes RNA-OUT that represses RNA-IN regulated targets, pMB1 plasmid origin encoded RNAI that represses RNAII regulated targets (SEQ ID NO: 31; SEQ ID NO:32; Grabherr and, Pfaffenzeller Supra, 2006; Cranenburgh R M. Supra, 2009), IncB plasmid pMU720 origin encoded RNAI that represses RNA II regulated targets (SEQ ID NO: 33; SEQ ID NO:34; Wilson et al Supra, 1997), ParB locus Sok of plasmid R1 that represses Hok regulated targets, Flm locus FlmB of F plasmid that represses flmA regulated targets (Morse M A, 1999 U.S. Pat. No. 5,922,583). An RNA selectable marker may be another natural antisense repressor RNAs known in the art such as those described in Wagner E G H, Altuvia S, Romby P. 2002. Adv Genet 46:361 and Franch T, and Gerdes K. 2000. Current Opin Microbiol 3:159. An RNA selectable marker may also be an engineered repressor RNAs such as synthetic small RNAs expressed SgrS, MicC or MicF scaffolds as described in Na D, Yoo S M, Chung H, Park H, Park J H, Lee S Y. 2013. Nat Biotechnol 31:170

ROP: Repressor of primer

sacB: Structural gene encoding Bacillus subtilis levansucrase. Expression of sacB in gram negative bacteria is toxic in the presence of sucrose

SEAP: Secreted alkaline phosphatase

shRNA: Short hairpin RNA

SR: Spacer region. As used herein, spacer region is the region linking the 5' and 3' ends of the eukaryotic region sequences. The eukaryotic region 5' and 3' ends are typically separated by the replication origin and selection marker. In simple single RNA Pol II transcription vectors this will be between the RNA Pol II promoter region (5' to a promoter, enhancer, boundary element, S/MAR) and the RNA Pol II polyA region (3' to a polyA sequence, eukaryotic transcriptional terminator sequence, boundary element, S/MAR). For example, in NTC9385R (FIG. 5) the spacer region is region between NheI site at 1663 and KpnI site at 460. In dual RNA Pol II transcription vectors, the eukaryotic sequences separated by the spacer will depend on the orientation of the two transcription elements. For example, with divergent or convergent RNA Pol II transcription units, the spacer region may separate two polyA sequences or two enhancers respectively. In RNA Pol II, RNA Pol III dual expression vectors, the spacer region may separate an RNA Pol II enhancer and a RNA Pol III promoter. The spacer region may encode bacterial or eukaryotic selectable markers, bacterial transcription terminators, eukaryotic transcription terminators, boundary elements, S/MARs, RNA Pol I or RNA Pol III expressed sequences or other functionalities

ssi: Single stranded initiation sequences

SV40 enhancer: Region containing the 72 bp and optionally the 21 bp repeats

target antigen: Immunogenic protein or peptide epitope, or combination of proteins and epitopes, against which an immune response can be mounted. Target antigens may be derived from a pathogen for infectious disease applications,

or derived from a host organism for applications such as cancer, allergy, or autoimmune diseases. Target antigens are well defined in the art. Some examples are disclosed in Williams, Supra, 2008 and are included herein by reference
TE buffer: A solution containing approximately 10 mM Tris pH 8 and 1 mM EDTA

Transcription terminator: Bacterial: A DNA sequence that marks the end of a gene or operon for transcription. This may be an intrinsic transcription terminator or a Rho-dependent transcriptional terminator. For an intrinsic terminator, such as the trpA terminator, a hairpin structure forms within the transcript that disrupts the mRNA-DNA-RNA polymerase ternary complex. Alternatively, Rho-dependent transcriptional terminators require Rho factor, an RNA helicase protein complex, to disrupt the nascent mRNA-DNA-RNA polymerase ternary complex. Eukaryotic: PolyA sites are not 'terminators', instead internal cleavage at PolyA sites leaves an uncapped 5' end on the 3'UTR RNA for nuclease digestion. Nuclease catches up to RNA Pol II and causes termination. Termination can be promoted within a short region of the poly A site by introduction of RNA Pol II pause sites (eukaryotic transcription terminator). Pausing of RNA Pol II allows the nuclease introduced into the 3' UTR mRNA after PolyA cleavage to catch up to RNA Pol II at the pause site. A nonlimiting list of eukaryotic transcription terminators known in the art include the C2x4 terminator (Ashfield R, Patel A J, Bossone S A, Brown H, Campbell R D, Marcu K B, Proudfoot N J. 1994. EMBO J 13:5656) and the gastrin terminator (Sato K, Ito R, Baek K H, Agarwal K, 1986. Mol. Cell. Biol. 6:1032). Terminator element may stabilize mRNA by enhancing proper 3'-end processing of mRNA (Kim D, Kim J D, Baek K, Yoon Y, Yoon J. 2003. Biotechnol Prog 19:1620)

Transgene: Target antigen or protein or RNA gene that is cloned into a vector

ts: Temperature sensitive

µg: microgram

µl: microliter

UTR: Untranslated region of a mRNA (5' or 3' to the coding region)

VARNA: Adenoviral virus associated RNA, including VARNAI (VAI or VA1) and or VARNAII (VAII or VA2) from any Adenovirus serotype, for example, serotype 2, serotype 5 or hybrids thereof

VARNAI: Adenoviral virus associated RNAI, also referred to as VAI, or VA1, from any Adenovirus serotype, for example, serotype 2, serotype 5 or hybrids thereof

Vector: A gene delivery vehicle, including viral (e.g. alphavirus, poxvirus, lentivirus, retrovirus, adenovirus, adenovirus related virus, etc) and nonviral (e.g. plasmid, midge, transcriptionally active PCR fragment, minicircles, bacteriophage, etc) vectors. These are well known in the art and are included herein by reference

Vector backbone: Eukaryotic region and bacterial region of a vector, without the transgene or target antigen coding region

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The invention relates generally to plasmid DNA compositions and methods to improve plasmid expression and plasmid production. The invention can be practiced to improve expression of vectors such as eukaryotic expression plasmids useful for gene therapy, genetic immunization and or interferon therapy. The invention can be practiced to improve the copy number of vectors such as eukaryotic

expression plasmids useful for gene therapy, genetic immunization and or interferon therapy. It is to be understood that all references cited herein are incorporated by reference in their entirety.

According to one preferred embodiment, the present invention provides for method of increasing in vivo expression of transgene from covalently closed super-coiled plasmid DNA, which comprises modifying the plasmid DNA to replace the pMB1, ColE1 or pBR322 derived replication origin and selectable marker with a replication origin selected from the group consisting of an P_{min} minimal pUC replication origin, ColE2-P9 replication origin, ColE2 related replication origin, and R6K replication origin and a RNA selectable marker; transforming the modified plasmid DNA into a Rep protein producing bacterial cell line rendered competent for transformation; and isolating the resultant transformed bacterial cells. The modified plasmid produced from these cells has increased transgene expression in the target organism.

In one preferred embodiment, the spacer region encoded pMB1, ColE1 or pBR322 derived replication origin is replaced with a CpG free ColE2 origin. In another preferred embodiment, a primosome assembly site is incorporated into a ColE2 plasmid DNA backbone to improve plasmid copy number. In yet another preferred embodiment, the pMB1, ColE1 or pBR322 derived replication origin is replaced with a CpG free R6K origin.

The methods of plasmid modification of the present invention have been surprisingly found to improve plasmid expression in the target organism. Increased expression vectors may find application to improve the magnitude of DNA vaccination mediated antigen reactive B or T cell responses for preventative or therapeutic vaccination, increase RNA and or protein transgene levels to improve gene replacement therapy or gene knockdown therapy, increase plasmid based expression levels of DNA vector expressed therapeutic antibodies that neutralize infectious diseases such as influenza, HIV, malaria, hepatitis C virus, tuberculosis, etc.

Plasmid encoded transgene expression in the target organism is preferably increased by employing specific constructs or compositions incorporated in a vector. According to one preferred embodiment, the present invention provides a composition for construction of a vector, comprising a ColE2 origin with at least 90% sequence identity to the sequences set forth as SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, and a RNA selectable marker and a eukaryotic region, wherein the ColE2 origin is operably linked to the RNA selectable marker and eukaryotic region. It has been surprisingly found that this ColE2 origin-RNA selectable marker improves plasmid encoded transgene expression in the target organism. According to another preferred embodiment, the resultant vector of the invention has at least 95% sequence identity to a sequence selected from the group consisting of: SEQ ID NO: 8, SEQ ID NO: 9.

According to another preferred embodiment, the present invention provides a composition for construction of a vector, comprising an R6K origin with at least 90% sequence identity to the sequences set forth as SEQ ID NO: 1, SEQ ID NO: 22, and a RNA selectable marker and a eukaryotic region, wherein the R6K origin is operably linked to the RNA selectable marker and eukaryotic region. It has been surprisingly found that this R6K origin-RNA selectable marker improves plasmid encoded transgene expression in the target organism. According to another preferred embodiment, the resultant vector of the invention

has at least 95% sequence identity to a sequence selected from the group consisting of: SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 39, SEQ ID NO: 40, SEQ ID NO: 41.

As used herein, the term "sequence identity" refers to the degree of identity between any given query sequence, e.g., SEQ ID NO: 2, and a subject sequence. A subject sequence may, for example, have at least 90 percent, at least 95 percent, or at least 99 percent sequence identity to a given query sequence. To determine percent sequence identity, a query sequence (e.g., a nucleic acid sequence) is aligned to one or more subject sequences using any suitable sequence alignment program that is well known in the art, for instance, the computer program ClustalW (version 1.83, default parameters), which allows alignments of nucleic acid sequences to be carried out across their entire length (global alignment). Chema et al., 2003 Nucleic Acids Res., 31:3497-500. In a preferred method, the sequence alignment program (e.g., ClustalW) calculates the best match between a query and one or more subject sequences, and aligns them so that identities, similarities, and differences can be determined. Gaps of one or more nucleotides can be inserted into a query sequence, a subject sequence, or both, to maximize sequence alignments. For fast pair-wise alignments of nucleic acid sequences, suitable default parameters can be selected that are appropriate for the particular alignment program. The output is a sequence alignment that reflects the relationship between sequences. To further determine percent identity of a subject nucleic acid sequence to a query sequence, the sequences are aligned using the alignment program, the number of identical matches in the alignment is divided by the length of the query sequence, and the result is multiplied by 100. It is noted that the percent identity value can be rounded to the nearest tenth. For example, 78.11, 78.12, 78.13, and 78.14 are rounded down to 78.1, while 78.15, 78.16, 78.17, 78.18, and 78.19 are rounded up to 78.2.

According to another preferred embodiment, the present invention provides methods and compositions for production of a Rep protein dependent plasmid vector. Production cell lines providing improved heat inducible P_L promoter expression of a Rep protein integrated into the genome and expressed from the heat inducible P_L promoter incorporating the OL1-G deletion (SEQ ID NO: 11), or the heat inducible P_L promoter incorporating the OL1-G to T substitution (SEQ ID NO: 12). It has been surprisingly found that these promoter modifications improves Rep protein dependent plasmid vector copy number in shake flask and fermentation cultures.

Turning now to the drawings, FIG. 1. shows an annotated map of the antibiotic free NTC8685 pUC origin expression vector with the locations of the pUC origin, PAS-BH primosomal assembly site, SV40 enhancer and other key elements indicated. The replication origin (PAS-BH and pUC origin) is from the AgeI (230) site to the DraIII (1548) site (1318 bp total). The antibiotic free RNA-OUT selection marker is between the DraIII (1548) and KpnI (1695) sites (147 bp total). The spacer region encoded bacterial region (replication and selection) of this vector is 1465 bp.

FIG. 2 shows an annotated map of the antibiotic-free NTC9685C ColE2 origin expression vector with the locations of the primosomal assembly site, ColE2 Replication origin (Replication origin) and other key elements indicated. The spacer region encoded bacterial region (replication and selection) of this vector is 281 bp [NheI site-ssiA-ColE2 Origin (+7)-RNA-OUT-KpnI site] (SEQ ID NO:20).

FIG. 3 shows an annotated map of the antibiotic-free NTC9685R R6K origin expression vector with the locations of the primosomal assembly site, R6K Replication origin

(R6K mini-origin) and other key elements indicated. The spacer region encoded bacterial region (replication and selection) of this vector is 466 bp [NheI site-trpA terminator-R6K Origin-RNA-OUT-KpnI site] (SEQ ID NO:19).

FIG. 4 shows an annotated map of the antibiotic-free NTC9385C ColE2 origin expression vector with the locations of the primosomal assembly site, ColE2 Replication origin (Replication origin) and other key elements indicated. The spacer region encoded bacterial region (replication and selection) of this vector is 281 bp [NheI site-ssiA-ColE2 Origin (+7)-RNA-OUT-KpnI site] (SEQ ID NO:20). This vector differs from NTC9685C in that the VA1 RNA and SV40 enhancer are not present.

FIG. 5 shows an annotated map of the antibiotic-free NTC9385R R6K origin expression vector with the locations of the primosomal assembly site, R6K Replication origin (R6K mini-origin) and other key elements indicated. The spacer region encoded bacterial region (replication and selection) of this vector is 466 bp [NheI site-trpA terminator-R6K Origin-RNA-OUT-KpnI site] (SEQ ID NO:19). This vector differs from NTC9685R in that the VA1 RNA and SV40 enhancer are not present.

FIG. 6 shows an annotated map of the pINT pR pL R6K Rep pi P42L-P106L-F107S (P3-) integration vector; key features such as the c1857 is repressor, P_L promoter, R6K Rep protein, HK022 phage attachment site for site specific integration into the E. coli genome, R6K replication origin and spectinomycin/streptomycin resistance marker (SpecR StrepR) are shown.

FIG. 7 show an annotated sequence of the P_L promoter with locations of the P_L promoter OL1, OL2 and OL3 repressor binding sites, -10 and -35 promoter elements for P_{L1} and P_{L2} promoters. The OL1 mutations OL1-G and OL1-G to T alterations are shown.

FIG. 8 shows an annotated map of the pINT pR pL ColE2 Rep protein integration vector; key features such as the c1857 is repressor, P_L promoter, ColE2 Rep protein, HK022 phage attachment site for site specific integration into the E. coli genome, R6K replication origin and spectinomycin/streptomycin resistance marker (SpecR StrepR) are shown.

FIG. 9 shows Nanoplasmid expression in vitro after lipofectamine transfection of HEK293 cell line of various EGFP transgene encoding vectors.

FIG. 10 shows Nanoplasmid expression in vivo after intramuscular injection with EP of various muSEAP transgene encoding vectors.

FIG. 11 shows a ColE2 origin Nanoplasmid shRNA expression vector. In this vector, a 22 bp shRNA is expressed from the RNA Polymerase III H1 promoter, with a TTTTTT terminator. The bacterial region is the NTC9385C and NTC9685C Bacterial region (SEQ ID NO:20).

FIG. 12 shows an IncB RNAI based RNA selection marker. A) Genomically expressed target of IncB RNAI RNA selection marker (SEQ ID NO: 35). Plasmid expressed RNAI binding to the pseudoknot in the complementary genomically expressed RNAII target prevents translation of the downstream SacB gene, conferring sucrose resistance. The RNAI -10 and -35 promoter elements are mutated to prevent RNAI expression. B) Structure of plasmid expressed IncB RNAI RNA selection marker (SEQ ID NO: 34) encoding the IncB RNAI antisense repressor (SEQ ID NO: 33).

The invention also relates to compositions and methods for producing high expression level plasmids. The present invention provides sequences that, when introduced into a vector backbone, increase plasmid expression.

The surprising observation that a ColE2 replication origin-RNA selection marker or R6K replication origin-RNA selection marker can be utilized as a plasmid expression enhancer is disclosed.

As described herein, plasmid expression is increased by replacement of the pMB1, ColE1 or pBR322 derived origin-selection marker bacterial region with an R6K origin-RNA selection marker in the plasmid backbone. In yet another preferred embodiment, the R6K origin is CpG free. In yet another preferred embodiment, the R6K origin is included with an RNA-OUT selection marker. In yet another preferred embodiment, the R6K origin is included with an pMB1 RNAI selection marker. In yet another preferred embodiment, the R6K origin is included with an IncB RNAI selection marker.

In yet another preferred embodiment, plasmid expression is increased by replacement of the pMB1, ColE1 or pBR322 derived origin-selection marker bacterial region with a ColE2 origin-RNA selection marker in the plasmid backbone. In yet another preferred embodiment, the ColE2 origin is CpG free. In yet another preferred embodiment, the ColE2 origin is included with an RNA-OUT selection marker. In yet another preferred embodiment, the ColE2 origin is included with an pMB1 RNAI selection marker. In yet another preferred embodiment, the ColE2 origin is included with an IncB RNAI selection marker. In yet another preferred embodiment, the ColE2 origin is included with a primosome assembly site.

In yet another preferred embodiment, plasmid expression is increased by replacement of the pMB1, ColE1 or pBR322 derived origin-selection marker with a P_{min} minimal pUC, ColE2 or a R6K origin in the plasmid backbone spacer region and an RNA selection marker in an intron. In yet another preferred embodiment, the R6K or ColE2 origin is CpG free. In yet another preferred embodiment, the RNA selection marker is the RNA-OUT selection marker. In yet another preferred embodiment, the RNA selection marker is the pMB1 RNAI selection marker. In yet another preferred embodiment, the RNA selection marker is the IncB RNAI selection marker.

EXAMPLES

The methods of the invention are further illustrated by the following examples. These are provided by way of illustration and are not intended in any way to limit the scope of the invention.

Example 1: Heat Inducible R6K Replication Origin Plasmid Production

Fermentation:

Fermentations were performed using proprietary fed-batch media (NTC3019, HyperGRO media) in New Brunswick BioFlo 110 bioreactors as described (Carnes and Williams, Supra, 2011). The seed cultures were started from glycerol stocks or colonies and streaked onto LB medium agar plates containing 6% sucrose. The plates were grown at 30-32° C.; cells were resuspended in media, and used to provide approximately 0.1% inoculums for the fermentations that contained 0.5% sucrose to select for RNA-OUT plasmids.

Antibiotic-free RNA-OUT plasmid fermentations were performed in E. coli strain XL1Blue [recA1 endA1 gyrA96 thi-1 hsdR17 supE44 relA1 lac [F' proAB lacIqZAM15 Tn10 (Tet^r)] (Stratagene, La Jolla, Calif.)] or GT115 [F—mcrA Δ(mrr-hsdRMS-mcrBC) φ801 acZAM15 ΔlacX74 recA1

rspL (StrA) endA1 Δdcm uidA(ΔMluI)::pir-116 ΔsbcC-sbcD (Invivogen, San Diego)] strains containing chromosomally integrated pCAH63-CAT RNA-IN-SacB (P5/6 6/6) at the phage lambda integration site as disclosed in Williams, J A Supra, 2008. SacB (Bacillus subtilis levansucrase) is a counterselectable marker which is lethal to E. coli cells in the presence of sucrose. Translation of SacB from the RNA-IN-SacB transcript is inhibited by plasmid encoded RNA-OUT. This facilitates plasmid selection in the presence of sucrose, by inhibition of SacB mediated lethality.

Analytical Methods:

Culture samples were taken at key points and at regular intervals during all fermentations. Samples were analyzed immediately for biomass (OD₆₀₀) and for plasmid yield. Plasmid yield was determined by quantification of plasmid obtained from Qiagen Spin Miniprep Kit preparations as described (Carnes and Williams, Supra, 2011). Briefly, cells were alkaline lysed, clarified, plasmid was column purified, and eluted prior to quantification. Agarose gel electrophoresis analysis (AGE) was performed on 0.8-1% Tris/acetate/EDTA (TAE) gels as described in Carnes and Williams J A, Supra, 2011.

R6K Background:

The R6K gamma plasmid replication origin requires a single plasmid replication protein π that binds as a monomer to multiple repeated 'iteron' sites (seven core repeats containing TGAGNG consensus) and as a dimer to repressive sites [TGAGNG (dimer repress) as well as to iterons with reduced affinity]. Various host factors are used including IHF, DnaA, and primosomal assembly proteins DnaB, DnaC, DnaG (Abhyankar et al 2003 J Biol Chem 278: 45476-45484). The R6K core origin contains binding sites for DnaA and IHF that affect plasmid replication (π , IHF and DnaA interact to initiate replication).

Different versions of the R6K gamma replication origin have been utilized in various eukaryotic expression vectors, for example pCOR vectors (Soubrier et al 1999, Gene Therapy 6:1482) and a CpG free version in pCpGfree vectors (Invivogen, San Diego Calif.), and pGM169 (University of Oxford). Incorporation of the R6K replication origin does not improve expression levels compared to an optimized pUC origin vector (Soubrier et al Supra, 1999). However, use of a conditional replication origin such as R6K gamma that requires a specialized cell line for propagation adds a safety margin since the vector will not replicate if transferred to a patients endogenous flora.

A highly minimalized R6K gamma derived replication origin that contains core sequences required for replication (including the DnaA box and stb 1-3 sites; Wu et al, 1995. J Bacteriol. 177: 6338-6345), but with the upstream π dimer repressor binding sites and downstream π promoter deleted (by removing one copy of the iterons, as with pCpG; see map below) was designed (SEQ ID NO:1) and NTC9685R and NTC9385R expression vectors incorporating it constructed (see Example 3).

Typical R6K production strains incorporate the π protein derivative PIR116 that contains a P106L substitution that increases copy number (by reducing π dimerization; π monomers activate while π dimers repress). Fermentation

results with pCOR (Soubrier et al., Supra, 1999) and pCpG plasmids (Hebel H L, Cai Y, Davies L A, Hyde S C, Pringle I A, Gill D R. 2008. Mol Ther 16: S110) were low, around 100 mg/L in PIR116 cell lines.

As expected, fermentation yields of the R6K expression vector NTC9685R-EGFP in R6K plasmid production cell line NTC641642 (GT115-SacB; GT115 modified for RNA-OUT AF vector selection by insertion of pCAH63-CAT RNA-IN-SacB (P5/6 6/6) into the genome. The GT115 genome encoded endogenous π gene P3 promoter constitutively expresses R6K replication protein π containing the pir-116 mutation; Metcalf et al, 1994; Gene 138; 1-7) were low (Table 1). Mutagenesis of the pir-116 replication protein and selection for increased copy number has been used to make new production strains. For example, the TEX2pir42 strain contains a combination of P106L and P42L. The P42L mutation interferes with DNA looping replication repression. The TEX2pir42 cell line improved copy number and fermentation yields with pCOR plasmids with reported yields of 205 mg/L (Soubrier F. 2004. World Patent Application WO2004033664). Methods to improve R6K origin yields are needed.

Other combinations of π copy number mutants have been shown to improve copy number. This includes 'P42L and P113S' and 'P42L, P106L and F107S' (Abhyankar et al 2004. J Biol Chem 279:6711-6719).

Two cell lines using the endogenous π gene P3 promoter to express π mutants 'P42L and P113S' (SEQ ID NO:13) (NTC640722 cell line) and 'P42L, P106L and F107S' (SEQ ID NO:14) were constructed and tested for copy number improvement with NTC9685R-EGFP. Two additional cell lines using the P_L promoter in addition to the endogenous π gene P3 promoter to express π mutants 'P42L and P113S' (NTC641981 cell line) and 'P42L, P106L and F107S' (NTC641053 cell line) were made and tested for copy number improvement with NTC9685R-EGFP. R6K production cell lines were made in XL1-Blue SacB (XL1-Blue att λ :P5/6 6/6-RNAIN-SacB, CmR).

These cell lines were constructed as follows. The R6K replication proteins π were cloned into the pINT pR pL integration vectors as described in Luke et al Supra, 2011 and included herein by reference. Constructed R6K Rep protein vectors were integrated into the genome at the HK022 phage attachment site as described in Luke et al, Supra, 2011. Briefly the pINT pINT pR pL R6K Rep vectors were amplified by PCR to delete the R6K replication origin, ligated to form a circle, and integrated into the HK022 attachment site using the pAH69 helper plasmid as described.

The results (Table 1) demonstrated that constitutive expression of 'P42L and P113S' or 'P42L, P106L and F107S' resulted in much higher levels of NTC9685R-EGFP than the NTC641642 encoded P106L Rep protein. However, constitutive expression from P3 resulted in low overall biomass and plasmid multimerization with P42L, P106L and F107S (RF306, RF314), due to high plasmid levels and resultant metabolic burden during the growth phase.

TABLE 1

2.6 kb NTC9685R-EGFP R6K Nanoplasmid fermentation yields in R6K rep cell lines										
Ferm #	Cell line	Rep Gene	Promoter	Growth phase temp (C.)	Growth spec yield ^a	Induced phase temp (C.)	Induced spec yield ^a	Final OD ₆₀₀	Final plasmid yield (mg/L)	Plasmid multimerization
RF310	NTC641642	P106L	Const P3	37	1.1	NA, 37	1.2	43	53	
RF323	NTC641642	P106L	Const P3	37	ND	NA, 37	0.9	54	49	Monomer
RF305	NTC640722	P42L-P113S	Const P3	30	3.4	37	3.6	96	345	Monomer
RF321	NTC641981	P42L-P113S	pR PL & P3	32	4.4-5.0	42	2.8	93	259	Monomer ^b
RF306	NTC641053	P42L-P106L-F107S	pR PL & P3	30	3.1	37	6.6	86	567	Multimer
RF314	NTC641053	P42L-P106L-F107S	pR PL & P3	32	2.6-4.7	42	7.1	79	558	Multimer
RF351	NTC661135	P42L-P106L-F107S	pR PL only	32	0.4	42	1.45	88	128	Monomer
RF326	NTC661135-MUT	P42L-P106L-F107S	pR PL OL1 G	32	0.64	42	6.7	81	545	Monomer
RF358	NTC711055	P42L-P106L-F107S	pR PL OL1 G	32	1	42	5.9	118	690	Monomer
RF359	NTC711231	P42L-P106L-F107S	pR PL OL1 G to T	30	1.8	42	8.5	82	695	Monomer

NTC640721 = NTC5402-P42L-P106L-F107S

NTC640722 = NTC5402-P42L-P113S

NTC641053 = NTC5402-pR pL P42L-P106L-F107S

NTC641642 = GT115-SacB (relA+) pir116 = P106L

NTC641981 = NTC5402-pR pL P42L-P113S

NTC661135 = NTC54208-pR pL P42L-P106L-F107S (P3-)

NTC661135-MUT = NTG54208-pR pL (OL1-G) P42L-P106L-F107S (P3-)

NTC661134 = NTC5402-pR pL P42L-P113S (P3-)

NTC711055 = NTC54208-pR pL (OL1-G) P42L-P106L-F107S (P3-)

NTC711231 = NTC54208-pR pL (OL1-G to T) P42L-P106L-F107S (P3-)

ND = Not determined

NA = Not applicable

^aSpecific Yield = mg plasmid/L/OD₆₀₀

^bSome dimer present

NTC5402 = XL1Blue, SacB

NTC54208 = XL1Blue, SacB, dem-

35

Heat inducible versions were then made by deletion of the P3 promoter to determine if P_L promoter mediated replication protein induction in a temperature shift improved R6K plasmid production yields and quality by reduction of plasmid copy number and metabolic burden during the reduced temperature growth phase. A strain encoding a deletion of the P3 promoter expressing P42L, P106L and F107S (NTC661135, incorporating a single copy of the pINT pR pL R6K Rep pi P42L-P106L-F107S (P3-) integration vector; FIG. 6, SEQ ID NO:17) constructed as described above dramatically reduced copy number during the reduced temperature growth phase with copy number induction after temperature upshift (Table 1; RF351). However, the yield (128 mg/L) was overall lower than with the P3 promoter (567, 558 mg/L).

However, excellent results were obtained after fermentation with a second NTC661135 cell line (RF326) in which plasmid copy number was increased 10 fold by temperature shifting, resulting in excellent final plasmid yields of 545 mg/L. PCR amplification and sequencing of the P42L, P106L and F107S expression cassette from the RF326 cell line (NTC661135-MUT) and the RF351 cell line (NTC661135) demonstrated that NTC661135-MUT contained a mutation in the OL1 lambda repressor binding site in the P_L promoter (FIG. 7; OL1-G this is a single base deletion that also reduces the distance between the P_L1 promoter -35 and -10 boxes from optimal 17 bp to 16 bp).

This mutation was introduced into the pINT pR pL R6K Rep pi P42L-P106L-F107S (P3-) integration vector by PCR mutagenesis and a sequence verified clone incorporating the OL1-G mutation integrated into the genome (NTC711055)

as described above. Fermentation evaluation of this cell line with the NTC9685R-EGFP plasmid (Table 1; RF358) demonstrated similar dramatic 6 fold heat inducible plasmid copy number induction, resulting in excellent final plasmid yields of 690 mg/L.

Repressor binding to OL1 is altered by mutations in OL1, such as OL1-G (FIG. 7; SEQ ID NO:11) and V2 (OL1-G to T; FIG. 7; SEQ ID NO:12; this is a G to T substitution that maintains the distance between the P_L promoter -35 and -10 boxes at the optimal 17 bp; this is the V2 mutation described by Bailone and Galibert, Supra, 1980).

The OL1-G to T (V2) mutation was introduced into the pINT pR pL R6K Rep pi P42L-P106L-F107S (P3-) integration vector by PCR mutagenesis and a sequence verified clone incorporating the OL1-G mutation integrated into the genome as described above to create NTC711231. Fermentation evaluation of this cell line with the NTC9685R-EGFP plasmid (Table 1; RF359) demonstrated, similar to OL1G, a dramatic 5 fold heat inducible plasmid copy number induction, resulting in excellent final plasmid yields of 695 mg/L.

Cell lines incorporating the pINT pR pL R6K Rep pi P42L-P106L-F107S (P3-) integration vector containing either the wildtype P_L promoter (NTC661135, SEQ ID NO:10), the OL1-G mutation (NTC711055, SEQ ID NO:11) or the OL1-G to T mutation (NTC711231, SEQ ID NO:12) were transformed with the NTC9385R-EGFP plasmid and yields in shake flask determined (Table 2). The results demonstrated the OL1-G and OL1-G to T mutations dramatically improve temperature inducible R6K plasmid yields in shake flask culture. Improved yield with two different R6K plasmids (NTC9385R, Table 2; NTC9685R,

65

Table 1) in either LB shake flask media or HyperGRO fermentation media demonstrates improved temperature inducible R6K plasmid is generic, and is not plasmid or growth media specific. Thus the invention can be utilized with a plurality of R6K origin vectors, in various plasmid growth media described in the art and various temperature induction profiles.

Likewise the pINT pR pL R6K Rep plasmids can be integrated into alternative E. coli strains to create production hosts. Any strain that is acceptable for plasmid production, such as JM108, BL21, DH5, DH1, DH5 α , GT115, GT116, DH10B, EC100, can be converted to a high yield temperature inducible R6K plasmid production host by integration of a pINT pR pL R6K Rep plasmid into the genome. The pR pL R6K Rep expression cassette may alternatively be removed from the pINT vector backbone and directly integrated into the chromosome, for example, using Red Gam recombination cloning (for example, using the methods described in Datsenko and Wanner 2000 Proc Natl Acad Sci USA 97:6640-6645). The pR pL R6K Rep expression cassette may alternatively be transferred to a different vector backbone, such as integration vectors that target different phage attachment sites, for example, those described by Haldimann and Wanner 2001, J Bacteriol 183:6384-6393.

TABLE 2

NTC9385R-EGFP LB media shake flask production yields in R6K production strains					
Cell Line	Rep Gene	Rep Gene Promoter	30° C. Spec yield ^a	32° C. Spec yield ^a	37° C. Spec yield ^a
NTC661135	P42L-P106L-F107S	P _R P _L	1.2	2.1	0.6
NTC711055	P42L-P106L-F107S	P _R P _L (OL1-G)	0.5	1.3	9.1
NTC711231	P42L-P106L-F107S	P _R P _L (OL1-G to T)	1.3	7.0	9.3

^aSpecific yield = mg plasmid/L/OD₆₀₀

These results are surprising since the art teaches that P_L promoter mutations in the OL1 binding site such as V2 (OL1-G to T) are constitutively active due to an inability of the lambda repressor to stop expression from the P_L promoter (Bailone and Galibert, Supra, 1980). While not limiting the application of the invention, it is possible that the lambda repressor is able to repress the P_L promoter through binding to the OL2 and OL3 sites (FIG. 7) when the P_L promoter is integrated in the genome; the lambda repressor may not be able to bind multiple copies of the mutated P_L promoter as in a phage infection.

The application of two independent OL1 mutations (OL1-G and OL1-G to T) to create cell lines for high yield R6K plasmid production demonstrates the general utility of P_L promoters incorporating OL1 mutations to improve heat inducible chromosomal expression of a target protein. Any OL1 mutation is contemplated for use in the current invention. New OL1 mutations can be defined by standard methods known in the art, for example error prone mutagenesis of the OL1 region, with subsequent selection of beneficial OL1 mutations by screening for heat inducible target protein production. The target protein can be a Rep protein as described herein, or a fluorescent marker, or any target protein or RNA. Thus application of P_L promoters incorporating OL1 mutations is contemplated generally as a platform for improved heat inducible chromosomal expression

of any recombinant protein or RNA. This can be applied to improve heat inducible chromosomal expression of any recombinant protein or RNA using either shake flask (Table 2) or fermentation (Table 1) culture.

5 These cell lines may also be used to produce alternative R6K plasmids, such as CpGfree vectors, pCOR vectors, pGM169, etc. P_L promoter vectors with the OL1 mutations may be used to improve expression of alternative target proteins or mRNAs from the genome.

10 These cell lines may also be used to produce alternative Rep protein dependent plasmids, such as Cole2-P9 replication origin plasmids (Examples 2 and 3), Cole2 related replication origin plasmids, etc. Numerous additional Rep protein dependent plasmids known in the art may also be produced using the cell lines of the invention. Many Rep protein dependent plasmids are described in del Solar et al Supra, 1998 which is included herein by reference.

Heat inducible target protein production may be further improved, by further mutating OL1-G and OL1-G to T or an alternative OL1 mutation to incorporate a mutation in the P_L-10 GATACT sequence to make it more closely match the consensus TATAAT (-35 is already consensus TTGACA (FIG. 7).

Alternative temperature sensitive (ts) lambda repressors (cI) may be substituted for the cITs857 mutation utilized in the pINT vectors. Multiple alternative ts lambda repressors have been defined (for example, see Lieb M. 1979 J Virol 32:162 incorporated herein by reference) or new ts lambda repressors may be isolated by screening for temperature sensitive cI function.

Alternative integration methods rather than the described pINT pR pL integration vectors may be utilized such as integration of the pR pL expression cassette into the genome at defined sites using Red Gam recombination cloning (for example, using the methods described in Datsenko and Wanner Supra, 2000).

Example 2: Cole2-P9 Replication Origin Plasmid Production

40 Similar to plasmid R6K, the Cole2 replication origin is separate from the replication protein, so the Cole2 replication origin theoretically may be utilized to construct Rep protein dependent plasmids. Here application of the Cole2 replication origin, using Cole2-P9 as an example, to produce Cole2 Rep protein dependent plasmids is demonstrated (Example 3).

Cole2 Background:

The Cole2 replication origin (for example, Cole2-P9) is highly conserved across the Cole2-related plasmid family (15 members are compared in Hiraga et al Supra, 1994, and 53 Cole2 related plasmid members including Cole3 are compared in Yagura et al Supra, 2006, both references are included herein by reference). Plasmids containing this origin are normally 10 copies/cell (low copy #). For application in gene therapy or DNA vaccination vectors, the copy number of Cole2 replication origin vectors needs to be improved dramatically.

Expression of the Cole2-P9 replication (Rep) protein is regulated by antisense RNA (RNAI). Copy number mutations have been identified that interfere with this regulation and raise the copy number to 40/cell (Takechi et al 1994 Mol Gen Genet 244:49-56).

pINT pR pL Cole2 Rep Protein Cell Line NTC641711:

65 The Cole2 Rep protein (SEQ ID NO:15) was expressed using the heat inducible pINT pR pL vector as described in Example 1. The Cole2 RNAI region was removed and

replaced with an optimal kozak-ATG region. This modification deletes the RNAI -10 promoter box. The Rep internal RNAI -35 box (Yasueda et al 1994 Mol Gen Genet 244: 41-48) was mutagenized (from (opposite strand) TTGAAG to CTGAAG) to lower the consensus. A high copy mutation in the Rep coding region (C139T; Nagase et al 2008 Plasmid 59:36-44) was also incorporated.

These changes do not alter the Rep protein amino acid sequence (SEQ ID NO:15).

The ColE2 Rep gene was PCR amplified from CGSC Strain #8203 with following primers

15061101: (SEQ ID NO: 26)
 ggaacgggatccagaaggagatatacatatgagtgcctacttcagcgct
 tcagggga

15061102: (SEQ ID NO: 27)
 ggaacggaattccttatcattttgagagatctggatcacat

The 920 bp PCR product was digested with BamHI/EcoRI and cloned into BamHI/EcoRI digested pINT pR pL BamHI/EcoRI (3766, polylinker). Recombinant clones were selected by restriction digestion and sequence verified. The map of the resultant pINT pR pL CoE2 Rep integration vector is shown in FIG. 8. The integration plasmid was integrated into NTC54208 (XL1Blue-sacB [dcm-]) to create cell line NTC641711 as described in Example 1.

A kanR ColE2-P9 replication origin fluorescent reporter plasmid (pDNAVACCUltra5-C2-P5/6,4/6-T7RBS EGFP) was constructed to select for copy number improving mutations. The 1067 bp pUC replication origin was removed from the kanR pDNAVACCUltra5-P5/6,4/6-T7RBS EGFP vector (the pDNAVACCUltra5-EGFP vector disclosed in Williams J A, 2006 World patent application WO06078979, modified to express the EGFP reporter in E. coli utilizing the weak constitutive P5/6,4/6 promoter disclosed in Lissemore J L, Jankowski J T, Thomas C B, Mascotti D P, deHaseth P L. 2000. Biotechniques 28: 82-89 and included herein by reference) by NheI-DraIII digestion, and replaced with a 132 bp ColE2-P9 replication origin (+7-ssiA; see below). Recombinant clones were recovered in cell line NTC641711 and the ColE2 origin confirmed by restriction digestion and sequence verification. This demonstrates that the ColE2-P9 Rep protein cell line NTC641711 can be used to select and propagate ColE2 replication origin containing plasmids.

ColE2 Rep Protein Mutagenesis, Selection of Copy Number Increasing Mutants

Background:

The ColE2 Rep protein binds as a monomer to the ColE2 replication origin. However, Rep protein exists mostly as a dimer in solution; Rep dimerization will limit the amount of active monomeric Rep which is hypothesized will maintain ColE2 plasmid at a low copy number (Han M, Aoki K, Yagura M, Itoh T. 2007. Biochem Biophys Res Commun 353:306). Copy number autoregulation by Rep protein dimerization is a common copy number control mechanism. Significantly, R6K Rep protein mutations such as P106L (PIR116) utilized in Example 1 that interfere with dimer formation dramatically increase copy number (Abhyankar et al Supra, 2004). It was hypothesized that ColE2 plasmid copy number can also be increased with a dimerization deficient Rep mutation.

Mutagenesis:

ColE2 Rep protein functional domains have been mapped and a region responsible for dimerization defined (FIG. 8).

The dimerization region was mutagenized using the Genemorph II Random Mutagenesis Kit (Stratagene) as described (Lanza A M, Alper H S. 2011. Methods in Molecular Biology, Vol. 765, Strain Engineering: Methods and Protocols, Ed. J. A. Williams, Humana Press Inc., Totowa, N.J. pp 253-274). The Rep gene was error prone PCR amplified from the pINT pR pL ColE2 Rep vector with the kit enzyme. The mutagenized dimerization domain (359 bp BstB1/EcoRI fragment; FIG. 8) was cloned back into the pINT pR pL ColE2 Rep vector replacing the non mutagenized 359 bp BstB1/EcoRI fragment. An integrated pINT pR pL ColE2 Rep library was then made by mass genome integration without purification of the mutagenized plasmid pool into NTC54208 containing the pAH69 integration plasmid. The integrated Rep library was transformed with the kanR pDNAVACCUltra5-C2-P5/6,4/6-T7RBS EGFP fluorescent ColE2 reporter plasmid and transformants plated on LB+kanamycin agar plates and grown at 37° C. This EGFP reporter plasmid allows 1) visual selection of plasmid copy number improvement using a Dark Reader for agar plate illumination; and 2) quantitative copy number evaluation (fluorescence is linear with copy number) in liquid culture using a fluorometer (BioTek FLx800 microplate fluorescence reader).

Two colonies were isolated from 30,000 screened cells with significantly higher colony fluorescence. Both cell lines were verified to have improved pDNAVACCUltra5-C2-P5/6,4/6-T7RBS EGFP plasmid copy number in liquid culture demonstrating increased fluorescence corresponds to increased copy number.

The lambda repressor-P_L-ColE2 Rep regions from genomic DNA from these two cell lines were amplified by PCR and sequenced to determine the basis for improvement. One colony had a mutation in the lambda repressor which presumably reduces the activity of the repressor leading to Rep protein overexpression. This demonstrates that alternations to the vector backbone that increase P_L promoter activity improve ColE2 plasmid copy number. Thus ColE2 copy number, like R6K plasmids, will be improved by making a cell line with the ColE2 Rep protein (or Rep protein copy number improving mutations) expressed from pINT pR pL vectors incorporating the lambda repressor binding site OL1 mutations (OL1-G and OL1-G to T) identified in Example 1.

The second colony had a mutation in the Rep protein (G194D; SEQ ID NO:16). This mutation was introduced back into the pINT pR pL ColE2 Rep vector to create the pINT pR pL ColE2 Rep protein mutant (G194D) (SEQ ID NO:18). The integration plasmid was integrated into NTC54208 (XL1Blue-sacB [dcm-]) to create cell line NTC701131 as described in Example 1. ColE2 plasmid production yields were improved in the ColE2 Rep protein mutant cell line NTC701131, compared to the parental ColE2 Rep protein cell line NTC641711 in both shake flask and fermentation culture (Table 3). This demonstrates that the ColE2 Rep protein, like the R6K Rep protein, can be mutagenized to create copy number improving variants.

Combining the ColE2 Rep protein G194D mutant with pINT pR pL vector incorporating the lambda repressor binding site OL1 OL1-G to T mutation identified in Example 1 further increased copy number (cell line NTC710351=NTC54208-pR pL (OL1-G to T) ColE2 rep G194D) and fermentation production yields (Example 3).

TABLE 3

NTC9385C-Luc plasmid performance in different processes and production cell lines ^a									
ColE2 Plasmid	LB shake flask (37 C.)			Plasmid + Shake flask (37 C.) ^c			HyperGRO fermentation		
production cell line	OD ₆₀₀	mg/L	Spec yield ^b	OD ₆₀₀	mg/L	Spec yield ^b	OD ₆₀₀	mg/L	Spec yield ^b
NTC641711	3.4	1.4	0.4	13.0	12.3	0.93	148	61	0.4
NTC701131	3.4	3.1	0.9	16.6	17.9	1.1	113	110	1.0
Rep mutant							140	142	1.0

^aAll plasmid preparations at harvest were high quality monomer.

^b Specific yield = mg plasmid/L/OD₆₀₀

^c Plasmid + media from Thomson Instruments Company

Additional rounds of mutagenesis of the wild type Rep protein, or mutagenesis of mutant Rep protein such as G194D may be performed to further improve copy number. The entire Rep protein or subfragments can be mutagenized (e.g. BamH1-EcoRI fragment for entire Rep protein; FIG. 8). The ideal mutant will be similar to the R6K Rep protein mutants 'P42L and P113S' and 'P42L, P106L and F107S' (Example 1) with higher copy number at 37-42° C. (i.e. higher levels of replication inducing monomeric Rep protein are produced from the heat inducible P_R P_L promoters) to facilitate adaptation into NTC's inducible fermentation plasmid production process as in Example 1.

ColE2 Origin Vectors:

The following vectors containing the minimal ColE2-P9 origin (Yagura and Itoh 2006 Biochem Biophys Res Commun 345:872-877) and various origin region modifications were constructed.

+7-ssiA:

This combines the ColE2 origin (+7) (SEQ ID NO:4) with ssiA from plasmid R6K (SEQ ID NO:21). Thus ssiA vectors contain, in addition to the ColE2-P9 origin, a downstream primosome assembly site. Like most plasmid origins, the ColE2 origin contains a primosomal assembly site about 100 bp downstream of the origin (Nomura et al Supra, 1991). This site primes lagging strand DNA replication (Masai et al 1990 J Biol Chem 265:15124-15133) which may improve plasmid copy number or plasmid quality. The ColE2 PAS (ssiA) is similar to PAS-BH (ColE1 ssiA=PAS-BL Marians et al 1982 J Biol Chem 257:5656-5662) and both sites (and PAS-BH) are CpG rich ØX174 type PAS. A CpG free PAS (ssiA from R6K; Nomura et al Supra, 1991; SEQ ID NO:21) that acts as a dnaA, dnaB dnaC (ABC) primosome on a dnaA box hairpin sequence (Masai et al 1990 J Biol Chem 265:15134-15144) was selected for inclusion in the +7-ssiA vectors. Alternative ABC or ØX174 type PAS sequences are functionally equivalent to ssiA from R6K, and may be substituted for ssiA in these ColE2 replication origin vectors.

+7-ssiA vectors were constructed by replacing the pUC origin NheI-DraIII region (FIG. 1) with a NheI-DraIII compatible synthetic ssiA-+7 ColE2 origin restriction fragment (FIG. 2, FIG. 4). Plasmids were transformed into ColE2 plasmid production NTC641711. The correct ColE2 vectors were identified by restriction digestion and sequence verified.

+7 (No ssiA):

This deletes the ssiA sequence from +7-ssiA while retaining the ColE2 origin (+7) (SEQ ID NO:4). The ssiA sequence was removed by NheI-MfeI digestion, the sites blunted with Klenow and the vector religated to delete the 64 bp ssiA region. Plasmids were transformed into ColE2

plasmid production host NTC641711. The correct ColE2 vector was identified by restriction digestion and sequence verified.

+7 CpG-ssiA:

This combines the ColE2 replication origin (+7 CpG) (SEQ ID NO:5) with ssiA from plasmid R6K (SEQ ID NO:21). The single CpG in the ColE2 replication origin (Table 4) was removed from the vector by site directed mutagenesis. Plasmids were transformed into ColE2 plasmid production host NTC641711. The correct ColE2 vector was identified by restriction digestion and sequence verified.

+16-ssiA:

This combines the ColE2 replication origin (+16) (SEQ ID NO:7) with ssiA from plasmid R6K (SEQ ID NO:21). A 16 bp region of homology downstream of the ColE2-P9 replication origin is conserved with the ColE3 replication origin. This 16 bp region was added to the vector by site directed mutagenesis. Plasmids were transformed into ColE2 plasmid production host NTC641711. The correct ColE2 vector was identified by restriction digestion and sequence verified.

Min-ssiA:

This combines the ColE2 Min replication origin (SEQ ID NO:6) with ssiA from plasmid R6K (SEQ ID NO:21). This is the minimal 32 bp ColE2 sequence sufficient for replication defined by Yasueda et al 1989 Mol Gen Genet 215:209) (SEQ ID NO:28), extended by an additional 6 bp (Table 4). This vector was created by site directed mutagenesis of the +7-ssiA clone. Plasmids were transformed into ColE2 plasmid production host NTC641711. The correct ColE2 vector was identified by restriction digestion and sequence verified.

The series of plasmids were transformed into ColE2 plasmid production host NTC701131 (Rep mutant). The resultant cell lines were then used determine plasmid copy number and quality (Table 4). Two different backbones were evaluated with the +7-ssiA and +16-ssiA ColE2 replication origins to determine the effect of plasmid sequence alterations.

The results demonstrate that the four replication origin variants containing the ssiA sequence [+7-ssiA; +16-ssiA; +7 (CpG free) ssiA; Min-ssiA] are functional in NTC701131, replicating to a similar copy number (0.73-1x). All plasmids were high quality monomer. This demonstrates that any of these minimal ColE2 origin variants can function as a plasmid replication origin to produce high quality plasmid.

Yagura et al Supra, 2006 have demonstrated that the Min ColE2 Replication origin (SEQ ID NO:28, which is reverse complement of residues 7-38, in FIG. 1 of Yagura et al Supra, 2006) can be further deleted without eliminating replication function. Yagura et al, Supra, 2006, demonstrated

15

20

25

30

35

40

45

50

55

60

65

that the core sequence is residues 8-35, with residues 5-36 are required for full activity. The +7 ColE2 Replication origin (SEQ ID NO:4; which is the reverse complement of residues 0-44 in FIG. 1 of Yagura et al Supra, 2006) could therefore be reduced to span residues 8-35 or 5-36 of FIG. 1 of Yagura et al Supra, 2006. Such vectors should replicate similarly to the disclosed vectors. As well, a number of base changes can be made within the core ColE2 origin 8-34 region that do not affect ColE2 replication (see changes to residues that retain function in Table 2 in Yagura et al Supra, 2006).

A surprising observation that is contrary to the teachings of Yagura et al Supra, 2006 is that the +7(CpG free)-ssiA ColE2 origin is fully functional. This origin contains a change of a G to C in residue 36 (FIG. 1 Yagura et al Supra, 2006). This change is predicted to reduce origin activity (Relative transformation frequency 5 fold reduced with 36 G-C to C-G; Table 2 in Yagura et al Supra, 2006). This may be due to the different context in the +7(CpG free)-ssiA ColE2 origin, or the longer origin fragment (0-44). Regardless, the 121 bp +7(CpG free)-ssiA ColE2 origin (SEQ ID NO 29) or +7(CpG free) ColE2 origin (SEQ ID NO 5) are smaller CpG free replication origin alternatives to the 260 bp CpG free R6K replication origins (SEQ ID NO:22). CpG free ColE2 origins may be utilized to construct CpG free plasmid vectors, or to retrofit the replication origin in existing vectors with a CpG free alternative replication origin. Combinations of a CpG free ColE2 or R6K replication origin with a CpG free RNA-OUT selection marker may be utilized to construct antibiotic free CpG free plasmid vectors, or to retrofit the selection marker-replication origin region in existing vectors with an antibiotic free-CpG free alternative selection marker-replication origin.

The ssiA sequence was not necessary for plasmid replication, although removal of ssiA in +7 (no ssiA) reduced copy number to 55% of +7 (ssiA). Thus inclusion of a primosomal assembly site is beneficial to ColE2 plasmid copy number.

TABLE 4

ColE2 Origin EGFP vector production in NTC701131 ColE2 production cell line ^b					
Origin ^c	Cell line ID#	OD600	mg/L	Specific Yield (mg/L/OD ₆₀₀)	Relative copy number ^a
+7-ssiA (NTC9385C)	071-	13.5	20.7	2.3	1x
	020-	11.4	20.4	1.8	
	2D	12.5	27.2	2.2	
				(2.1 avg)	
+16-ssiA	071-	19.6	23.5	1.2	0.70x
	029-	16.5	20.9	1.3	
	1A	13.1	24.8	1.9	
				(1.5 avg)	
+7(CpG free)-ssiA	071-	10.5	25.1	2.4	0.83x
	020-	8.4	11.6	1.4	
	3D	10.1	14.1	1.4	
				(1.7 avg)	
Min-ssiA	071-	11.3	13.7	1.2	0.73x
	020-	15.9	37.2	2.3	
	4D	13.3	14.8	1.1	
				(1.5 avg)	
+7 (no ssiA)	071-	12.8	17.2	1.3	0.55x
	020-	13.5	16.4	1.2	
	5D	13.5	14.5	1.1	
				(1.2 avg)	

TABLE 4-continued

ColE2 Origin EGFP vector production in NTC701131 ColE2 production cell line ^b					
Origin ^c	Cell line ID#	OD600	mg/L	Specific Yield (mg/L/OD ₆₀₀)	Relative copy number ^a
+7-ssiA (NTC9685C)	071-	14.7	22.3	1.5	0.70x
	020-	13.2	21.3	1.6	
	6D	11.7	14.9	1.3	
				(1.5 avg)	
+16-ssiA (VA1-SV40)	071-	13.5	26.9	1.9	0.76x
	020-	11.9	17.7	1.5	
	7D	13.2	18.8	1.4	
				(1.6 avg)	

^aAverage specific yield/+7-ssiA average specific yield (specific yield = mg plasmid/L/OD₆₀₀)

^bPlasmid + media, 37 °C. throughout growth conditions. All plasmid preparations at harvest were high quality monomer NTC ColE2 origin sequences

^cNTC ColE2 origin sequences
 +7: caaaaggg**g**cgctg**ttatctgataaggcttatctggtctcattttg** (SEQ ID NO: 4) Min bold underlined
 +7 (CpG free): caaaaggg**g**cgctg**ttatctgataaggcttatctggtctcattttg** (SEQ ID NO: 5) (C to G change in bold underlined core to eliminate CpG is uppercase double underlined)
 Min: The 32 bp minimal origin defined by Yasueda et al Supra, 1989 (SEQ ID NO: 28) is Bold underlined):
 +16: CTGCTCAAAAAGACGCCaaaag
 ggcgctg**ttatctgataaggcttatctggtctcattttg** (SEQ ID NO: 7) Min bold underlined, additional 16 bp in +16 is uppercase

Example 3: NTC9382C, NTC9385C, NTC9382R, NTC9385R, NTC9682C, NTC9685C, NTC9682R, and NTC9685R Vectors

A series of AF eukaryotic expression vectors incorporating these novel ColE2-P9 derived vector origins were made. To replace the pUC origin, the +7 (ssiA) ColE2 origin from Example 2 was selected as well as the R6K origin (SEQ ID NO:1) from Example 1. The features of these vectors (NTC9382C, NTC9385C, NTC9382R, NTC9385R, NTC9682C, NTC9685C, NTC9682R, and NTC9685R) are summarized in Table 5.

NTC9682C, NTC9685C (FIG. 2), NTC9682R, NTC9685R (FIG. 3) are antibiotic-free RNA-OUT ColE2 origin (C) or R6K origin (R) versions of the pUC origin (C) or R6K origin (R) equivalents disclosed in Williams J A, Supra, 2010. These vectors contain the SV40 enhancer upstream of the CMV enhancer, and Adenoviral serotype 5 VA RNAI regulatory RNA (VARNAI).

NTC9382C, NTC9385C (FIG. 4), NTC9382R, NTC9385R (FIG. 5) are versions without the SV40 enhancer or VARNAI sequences.

NTC9682C, NTC9682R, NTC9382C, and NTC9382R all express the secreted transgene product as TPA fusion proteins while NTC9685C, NTC9685R, NTC9385C, and NTC9385R all express the native transgene product from a vector encoded ATG start codon.

The remainder of the vector sequences is identical between the different vectors, with the exception that the two R vectors NTC9682R and NTC9382R (FIG. 3, FIG. 5) contain the trpA bacterial terminator, which is absent in the two C vectors NTC9682C and NTC9382C (FIG. 2, FIG. 4).

An R6K gamma origin vector was constructed by swapping in the R6K gamma origin (SEQ ID NO:1) in a NotI-DraIII R6K origin synthetic gene for the corresponding NotI-DraIII pUC origin region in NTC8685. The NTC9682R, NTC9685R NTC9382R, NTC9385R vectors

were made by standard restriction digestion mediated fragment swaps. The ColE2 origin vectors were constructed in a similar fashion, by swapping in the +7 ssiA ColE2 origin in a NheI-DraIII synthetic gene for the corresponding NheI-DraIII pUC origin region. The NTC9682C, NTC9685C, NTC9382C, NTC9385C vectors were made by standard restriction digestion mediated fragment swaps. The 466 bp Bacterial region [NheI site-trpA terminator-R6K Origin-RNA-OUT-KpnI site] for NTC9385R and NTC9685R is shown in SEQ ID NO:19. The 281 bp Bacterial region [NheI site-ssiA-ColE2 Origin (+7)-RNA-OUT-KpnI site] for NTC9385C and NTC9685C is shown in SEQ ID NO:20.

High fermentation yields in HyperGRO media are obtained with these vectors. For example 695 mg/mL with NTC9685R-EGFP in R6K production cell line NTC711231 (Table 1) and 672 mg/L with NTC9385C-EGFP in ColE2 production cell line NTC710351.

These are just a few possible nonlimiting vector configurations. Many alternative vector configurations incorporating the novel R6K or ColE2 origin vector modifications may also be made, including but not limited to vectors with alternative selection markers, alternative promoters, alternative terminators, and different orientations of the various vector-encoded elements or alternative R6K or ColE2 origins as described in Examples 1 and 2.

TABLE 5

NTC9382C, NTC9385C, NTC9382R, NTC9385R, NTC9682C, NTC9685C, NTC9682R, and NTC9685R vectors				
Vector	Origin	VA RNAI present	SV40 enhancer	Transgene targeting
NTC9382C	ColE2-P9	No	No	Secretion (TPA)
NTC9382R	R6K	No	No	Secretion (TPA)
NTC9682C	ColE2-P9	Yes	Yes	Secretion (TPA)
NTC9682R	R6K	Yes	Yes	Secretion (TPA)
NTC9385C (SEQ ID NO: 8)	ColE2-P9	No	No	Native
NTC9385R (SEQ ID NO: 2)	R6K	No	No	Native
NTC9685C (SEQ ID NO: 9)	ColE2-P9	Yes	Yes	Native
NTC9685R (SEQ ID NO: 3)	R6K	Yes	Yes	Native

An example strategy for cloning into these vectors is outlined below.

GTCGACATG-----Gene of interest-----Stop codon
Sall

-----AGATCT
BglIII

For the NTC9385C, NTC9685C, NTC9385R, and NTC9685R vectors, the ATG start codon (double underlined) is immediately preceded by a unique Sall site. The Sall site is an effective Kozak sequence for translational initiation. In NTC9382C, NTC9682C, NTC9382R, and NTC9682R, the Sall site is downstream in frame with the optimized TPA secretion sequence (SEQ ID NO:25). The TPA ATG start codon is double underlined and the Sall site single underlined. SEQ ID NO:25: TPA secretion sequence atggatgcaatgaagagaggctctgctgtgtgctgctgtgtggagcagtctt cgtttcgcccagcggatccggtaccggtcgcac

For precise cloning, genes are copied by PCR amplification from clones, cDNA, or genomic DNA using primers with Sall (5' end) and BglIII (3' end) sites. Alternatively,

genes are synthesized chemically to be compatible with the unique Sall/BglIII cloning sites in these vectors.

For NTC9385C, NTC9685C, NTC9385R, and NTC9685R, the start codon ATG may immediately follow the Sall site (GTCGACATG) since the Sall site is a high function Kozak sequence. For all vectors one or two stop codons (preferably TAA or TGA) may be included after the open reading frame, prior to the BglIII site. A PCR product or synthetic gene designed for NTC9385C, NTC9685C, NTC9385R, and NTC9685R is compatible with, and can also be cloned into, the NTC9382C, NTC9682C, NTC9382R, and NTC9682R vectors.

EGFP and muSEAP transgene versions NTC9385C, NTC9685C, NTC9385R, and NTC9685R were constructed by standard restriction fragment swaps. The muSEAP gene is secreted using its endogenous secretion signal, while EGFP is cell associated. Expression levels in vitro were determined using EGFP, while expression levels in vivo were determined using muSEAP. Expression levels were compared to the NTC8685 parent vector, the gWIZ vector, and a minicircle comparator.

Adherent HEK293 (human embryonic kidney), A549 (human lung carcinoma), cell lines were obtained from the American Type Culture Collection (Manassas, Va., USA). Cell lines were propagated in Dulbecco's modified Eagle's medium/F12 containing 10% fetal bovine serum and split (0.25% trypsin-EDTA) using Invitrogen (Carlsbad, Calif., USA) reagents and conventional methodologies. For transfections, cells were plated on 24-well tissue culture dishes. plasmids were transfected into cell lines using Lipofectamine 2000 following the manufacturer's instructions (Invitrogen).

Total cellular lysates for EGFP determination were prepared by resuspending cells in cell lysis buffer (BD Biosciences Pharmingen, San Diego, Calif., USA), lysing cells by incubating for 30 min at 37° C., followed by a freeze-thaw cycle at -80° C. Lysed cells were clarified by centrifugation and the supernatants assayed for EGFP by FLX800 microplate fluorescence reader (Bio-Tek, Winooski, Vt., USA). The results are summarized in FIG. 9 and Table 6.

Groups of five mice were injected with plasmid DNA in an IACUC-approved study. Five micrograms of muSEAP plasmid in 25 or 50 µL of phosphate-buffered saline (PBS) was injected intramuscularly (IM) into a tibialis cranialis muscles of female BALB/c mice or ND4 Swiss Webster mice (6 to 8 weeks old) followed by Ichor TriGrid electroporation. SEAP levels in serum were determined using the Phospha-light SEAP Reporter Gene Assay System from Applied Biosystems (Foster City, Calif.) according to the manufacturer's instructions. The results are summarized in FIG. 10 and Table 6.

The NTC9385C, NTC9685C, NTC9385R, and NTC9685R vectors had similar expression to the parent NTC8685 vector in vitro, and higher expression than the gWIZ comparator (FIG. 9). Thus substitution of the R6K or ColE2 replication origin for the pUC origin was not detrimental for eukaryotic cell expression. However, surprisingly, in vivo expression was dramatically improved compared to NTC8685 or gWIZ with the ColE2 and R6K origin vectors (FIG. 10). For example the NTC9385C vector was unexpectedly improved 1.5 to 3.8× that of NTC8385 (Table 6) or NTC8685 (not shown) after IM delivery with EP.

TABLE 6

gWIZ and NTC9385C Nanoplasmid expression compared to NTC8685					
Plasmid	% NTC8685 expression in vitro ^a	% NTC8685 expression T = 7 days BALB/c ^b	% NTC8685 expression T = 7 days ND4 ^b	% NTC8685 expression T = 28 days BALB/c ^b	% NTC8685 expression T = 28 days ND4 ^b
	gWIZ	58	59	57	21
NTC8385	NA	NA	101	NA	101
NTC9385C	92	377	349	150	233
Minicircle ^c	NA	89	NA	40	NA

^a 100 ng/well EGFP transgene vectors transfected with lipofectamine into HEK293 cells

^b murine SEAP (muSEAP) transgene vectors in 8-10 week old BALB/c or ND4 Swiss Webster female mice, 5 µg dose with EP intramuscular into one anterior tibialis muscle followed by Ichor TriGrid electroporation. 25 µL dose for ND4 mice, 50 µL dose for BALB/c.

^c Minicircle equivalent to NTC9385C or NTC9385R, with NheI-KpnI region containing the replication origin and RNA-OUT selection marker (bacterial region) removed from NTC8385-muSEAP by SpeI/NheI digestion, gel purification of the eukaryotic region, in vitro ligation and supercoiling with DNA gyrase. The SpeI site is the same site used to truncate the CMV promoter to make NTC8685 and the NTC9385C-muSEAP vector so the minicircle eukaryotic region is the same as NTC9385C-muSEAP, the difference being the C2 and RNA-OUT region including the KpnI site is deleted in the minicircle.
NA = Not assayed

This improved in vivo expression was not specific to the CMV promoter. Versions of NTC8685-muSEAP and NTC9385C-muSEAP were constructed in which the murine creatine kinase (MCK) promoter (3 copies of the MCK Enhancer upstream of the MCK promoter and 50 bp of the MCK exon 1 leader sequence; Wang B, Li J, Fu F H, Chen C, Zhu X, Zhou L, Jiang X, Xiao X. 2008. Gene Ther 15:1489) was substituted for the CMV promoter. The swaps replaced the entire CMV enhancer CMV promoter-exon 1 leader (NTC8685: from a XbaI site immediately after the SV40 enhancer to a SacII site in the CMV derived exon 1 leader sequence FIG. 1; NTC9385C: from the KpnI site to a SacII site in the CMV derived exon 1 leader sequence FIG. 4) with the MCK enhancer, MCK promoter-exon 1 leader retaining the HTLV-I R portion of exon 1. Purified plasmid DNA from the resultant vectors, NTC8685-MCK-muSEAP (4847 bp) and NTC9385C-MCK-muSEAP (3203 bp), was injected IM into one anterior tibialis muscle of 8-10 week old BALB/c female mice (5 mice/group), 5 µg dose in 50 µL, followed by Ichor TriGrid electroporation as described in Table 6. SEAP levels in serum was determined on day 28 (T=28) post delivery. The NTC9385C-MCK-muSEAP vector (98.4±55.8) had 4.5× higher average expression than NTC8685-muSEAP (22.0±10.9). All 5 NTC9385C-MCK-muSEAP injected mice had higher muSEAP levels than any of the NTC8685-muSEAP mice. This demonstrates that improved in vivo expression with the Nanoplasmid vectors of the invention is not specific to the CMV promoter.

While the basis for expression improvement is unknown, it is not simply due to the size difference between the parent pUC origin vectors and the modified R6K origin-RNA selection marker or Cole2 origin-RNA selection marker vectors of the invention, since expression was not improved with a minicircle comparator vector that contains no bacterial region (Table 6). This demonstrates improved in vivo expression with the R6K origin-RNA selection marker or Cole2 origin-RNA selection marker vectors is not the result of simple elimination of a threshold amount of bacterial region sequences.

Reduction of the vector spacer region size as described herein by replacement of the spacer region replication origin and selection marker with the R6K, Cole2 origin-RNA selection marker vectors of the invention will also increase the duration of in vivo expression since expression duration is improved with plasmid vectors in which the bacterial region is removed (minicircle) or replaced with a spacer region of up to at least 500 bp (Lu J, Zhang F, Xu S, Fire A

20

Z, Kay M A. 2012. Mol Ther 20:2111-9). Thus the replicative minicircle vectors of the invention also have additional utility for applications requiring extended duration expression, such as: liver gene therapy using hydrodynamic delivery with transgenes such as α -1 antitrypsin (AAT) for AAT deficiency, Coagulation Factor VIII for Hemophilia A Therapy or Coagulation Factor IX for Hemophilia B Therapy etc; lung gene therapy with transgenes such as Cystic fibrosis transmembrane conductance regulator (CFTR) for cystic fibrosis etc; muscle gene therapy with transgenes such as the GNE gene for Hereditary inclusion body myopathies (HIBM), or dystrophin or dystrophin minigenes for duchenne muscular dystrophy (DMD).

25

Example 4: Spacer Region and Intron Modified Nanoplasmid Vectors

NTC8685 (SR=1465 bp) has much lower expression than NTC9385R (SR=466 bp) and NTC9385C (SR=281 bp). A minimal pUC origin vector was constructed with an 866 bp spacer region (NTC8385-Min; contains P_{min} minimal pUC origin-RNA-OUT). These vectors were tested for expression in vitro (lipofectamine 2000 delivery) and in vivo after intradermal electroporation delivery. As with Intramuscular injection (Example 3), the results (Table 7) demonstrated Cole2 and R6K origin vector dramatically improved in vivo expression after intradermal delivery compared to NTC8685. For example the NTC9385C vector was unexpectedly improved 2.7 to 3.1× compared to NTC8685 while the NTC9385R vector was unexpectedly improved 5.3 to 6.3× that of NTC8685 (Table 7). The 866 bp minimal pUC origin vector also improved expression to 1.4-1.9× that of NTC8685. This demonstrates improved in vivo expression with the NTC9385C and NTC9385R vectors is not limited to muscle tissue, and is observed also after intradermal delivery. Inclusion of the C2×4 eukaryotic transcription terminator in the NTC9385C vector further improved in vivo expression to 2.9 to 4.1× compared to NTC8685. This demonstrates improved in vivo expression with Nanoplasmid vectors may be obtained with alternative/additional sequences flanking the bacterial region.

A NTC9385R derivative was made in which the RNA-OUT antibiotic free marker was transferred to the intron (NTC9385Ra-O2 SEQ ID NO:39; RNA-OUT SEQ ID NO:23) inserted into the unique HpaI site in the intron (SEQ ID NO: 30). This vector encodes the R6K replication origin in the spacer region (SR=306 bp). To determine splicing

accuracy NTC9385Ra-O2-EGFP was transfected into the A549 cell line and cytoplasmic RNA isolated. The RNA was reverse transcribed using an EGFP specific primer, and PCR amplified using Exon 1 and Exon 2 specific primers. The resultant PCR product (a single band) was determined by sequencing to be the correct spliced exon1-exon2 fragment. This demonstrated that intronic RNA-OUT is accurately removed by splicing and does not interfere with splicing accuracy. NTC9385Ra-O2-EGFP also demonstrated improved in vivo expression compared to NTC8685 (Table 7: 1.6-3.5x). This demonstrates that Nanoplasmid vectors with improved expression of the current invention may encode the RNA selection marker in the intron rather than the spacer region.

The improved expression level after intradermal delivery demonstrates the application of Nanoplasmid vectors of the invention for cutaneous gene therapy applications, for example, for wound healing, burns, diabetic foot ulcer, or critical limb ischemia therapies using growth factors such as hypoxia inducible factor, hypoxia inducible factor 1 α , keratinocyte growth factor, vascular endothelial growth factor (VEGF), fibroblast growth factor-1 (FGF-1, or acidic FGF), FGF-2 (also known as basic FGF), FGF-4, placental growth factor (PlGF), angiotensin-1 (Ang-1), hepatic growth factor (HGF), Developmentally Regulated Endothelial Locus (Del-1), stromal cell derived factor-1 (SDF-1), etc.

vectors <1.2 kb are highly resistant to shear forces used with gene therapy delivery formulation (Catanese et al 2012. Gene Ther 19:94-100).

RNA Pol III Nanoplasmid vectors were made by standard restriction digestion mediated fragment swaps to combine either U6 or HI RNA Pol III promoter-target RNA-TTTTTT terminator (Eukaryotic region) with either the 466 bp Bacterial region [NheI site-trpA terminator-R6K Origin-RNA-OUT-KpnI site; SEQ ID NO:19] for NTC9385R-U6 and NTC9385RE-U6 vectors (Table 8) or the 281 bp Bacterial region [NheI site-ssiA-Cole2 Origin (+7)-RNA-OUT-KpnI site; SEQ ID NO:20] for NTC9385C-U6 and NTC9385CE-U6 vectors (Table 8). Versions were modified to express from the U6 promoter eRNA18, a single stranded RNA the expression of which can be quantified by Reverse transcriptase dependent RT-PCR. Vector performance (U6 promoter mediated eRNA18 RNA expression) was determined in total RNA extracted from HEK293 at either 25 or 48 hrs after lipofectamine 2000 mediated transfection as described (Luke J, Simon G G, Soderholm J, Errett J S, August J T, Gale M Jr, Hodgson C P, Williams J A. 2011. J Virol. 85:1370). These results (Table 9) demonstrate Nanoplasmid RNA Pol III vectors direct dramatically improved RNA expression relative to a plasmid RNA Pol III vector (NTC7485-U6-eRNA18) comparator.

TABLE 7

SR vector expression in vitro and in vivo							
muSEAP Vector ^b	SR ^a	SR (bp) Intron ^a	A549 (A ₄₀₅) ^d	HEK-293 (A ₄₀₅) ^d	ID + EP ^c (pg/mL) T = 4	ID + EP ^c (pg/mL) T = 7	ID + EP ^c (pg/mL) T = 14
NTC8685	T-VA1-BH-P-AF→	1465 HR- β	0.240 \pm 0.029 (1.0x)	3.002 \pm 0.188 (1.0x)	1.9 \pm 1.2 (1.0x)	6.7 \pm 4.1 (1.0x)	5.0 \pm 3.9 (1.0x)
NTC8385-Min ^e	T-P _{min} -AF→	866 HR- β	0.495 \pm 0.027 (2.1x)	2.713 \pm 0.177 (0.9x)	3.7 \pm 2.7 (1.9 x)	12.4 \pm 8.1 (1.9 x)	7.1 \pm 5.2 (1.4 x)
NTC9385R (SEQ ID NO: 2)	T←R-AF→	466 HR- β	0.604 \pm 0.04 (2.5x)	3.036 \pm 0.169 (1.0x)	12.0 \pm 7.4 (6.3 x)	35.5 \pm 31.1 (5.3 x)	29.9 \pm 23.4 (6.0 x)
NTC9385C (SEQ ID NO: 8)	←C-AF→	281 HR- β	0.267 \pm 0.053 (1.1x)	2.720 \pm 0.228 (0.9x)	5.8 \pm 3.0 (3.1 x)	20.8 \pm 9.6 (3.1 x)	13.5 \pm 9.8 (2.7x)
NTC9385C C2x4	←C-AF→	281 HR- β	0.214 \pm 0.017 (0.89x)	2.472 \pm 0.197 (0.82x)	5.6 \pm 2.3 (2.9 x)	27.7 \pm 20.3 (4.1 x)	16.0 \pm 14.3 (3.2x)
NTC9385R a-O2 (SEQ ID NO: 39)	T←R←AF- β	306 HR- β	0.524 \pm 0.071 (2.2x)	3.065 \pm 0.220 (1.0x)	3.6 \pm 2.8 (1.9 x)	23.4 \pm 16.5 (3.5 x)	7.8 \pm 8.0 (1.6 x)

^a Prokaryotic terminator = T; HTLV-IR = HR; B globin 3' acceptor site = β ; RNA-OUT = AF; pUC origin = P; minimal pUC origin = P_{min}; R6K γ origin = R; ColE2-P9 origin = C; C2x4 eukaryotic transcription terminator = C2x4

^b All plasmids produced in XL1Blue dem- host strains. P vectors were produced in dem- XL1Blue NTC54208; R vectors were produced in dem- R6K rep cell line NTC711231 (OL1 G to T); C vectors were produced in dem- ColE2 rep cell line NTC710351 (OL1 G to T).

^c Dose = 50 μ g in 50 μ l saline injected intradermal (ID) with EP on day 0. 6 mice/group. Mean \pm SD pg/mL muSEAP reported for day 4, 7 and 14. () Mean muSEAP standardized to NTC8685

^d muSEAP plasmid DNA transfected with Lipofectamine 2000. Mean \pm SD A₄₀₅ reported 48 hrs post transfection. () Mean A₄₀₅ standardized to NTC8685

^e P_{min} minimal pUC origin (SEQ ID NO: 42) and RNA-OUT (bacterial region = SEQ ID NO: 43)

55

Example 5: RNA Pol III Nanoplasmid Vectors

An example Nanoplasmid vector for RNA Pol III directed expression of RNA is shown in FIG. 11. This vector contains the human H1 RNA Pol III promoter, but an alternative promoter such as the murine U6 promoter can be substituted. This example vector expresses a 22 bp shRNA target RNA, but alternative RNAs may be expressed, including shorter or longer shRNAs, microRNAs, aptamer RNAs, hairpin RNAs, etc. This example vector is very small, with a monomer size of 442 bp. Small size is advantageous, since

Random 22 bp shRNA (KP2F11) versions of NTC9385CE-U6 (903 bp NTC9385CE-U6-KP2F11 shRNA propagated in ColE2 rep cell line NTC710351) and NTC9385R-U6 (855 bp NTC9385R-U6-KP2F11 shRNA propagated in R6K rep cell line NTC711231) were fermented in HyperGRO media as described in Example 1 except fermentation and cultures for inoculations were grown at 37 $^{\circ}$ C. throughout. Final yields were 149 mg/L (NTC9385CE-U6-KP2F11) and 216 mg/L (NTC9385R-U6-KP2F11 shRNA). This demonstrates that Nanoplasmid vectors for RNA Pol III expression (and RNA Pol II; Example

65

3) have superior manufacturing simplicity and yield compared to shRNA expressing minicircle vectors (Zhao et al 2011. Gene Ther 18:220-224). For example, optimal manufacture of minicircle vectors yields only 5 mg of minicircle per liter culture (Kay M A, He C Y, Chen Z Y 2010. Nat Biotechnol 28:1287-1289).

from transfected HEK293 and A549 cells using the protein and RNA isolation system (PARIS kit, Ambion, Austin Tex.) and quantified by A_{260} . Samples were DNase treated (DNA-free DNase; Ambion, Austin Tex.) prior to reverse transcription using the Agpath-ID One step RT-PCR kit (Ambion, Austin Tex.) with a EGFP transgene specific complementary

TABLE 8

RNA Pol III Nanoplasmid vector expression						
Vector	Pol II Enhancer	Size (bp)	Transfection 1: RNA isolated 48 hr post transfection		Transfection 2: RNA isolated 25 hr post transfection	
			HEK pg RNA/100 ng mRNA ^a	HEK Std ^b	HEK pg RNA/100 ng mRNA ^a	HEK Std ^b
NTC9385R-EGFP (negative control)	None	NA	0.0 ± 0.0	0%		
NTC8885MP-U6-eRNA18	SV40	1578 ^c	62.2 ± 5.9 (1.3x)	69%		
NTC9385RE-U6-eRNA18	SV40	1178	119.1 ± 13.9 (2.5x)	98%		
NTC9385R-U6-eRNA18	None	945 ^d	123.7 ± 8.0 (2.6x)	82%		
NTC9385CE-U6-eRNA18	SV40	993	119.0 ± 13.9 (2.5x)	83%		
NTC9385C-U6-eRNA18	None	760 ^e	131.1 ± 10.5 (2.7x)	70%	57.3 ± 2.6 (5.0x)	127%
NTC7485-U6-eRNA18	SV40	2978	48.0 ± 1.3 (1x control)	100% (100% control)	11.5 ± 1.5 (1x)	100% (control)

^apg eRNA18 target/100 ng total RNA isolated post-transfection.

^bStandardized mU6 expression compared to NTC7485-U6 shRNA eRNA18 vector (C) = test vector average pg RNA/C vector average pg RNA × test vector size (bp)/2978 × 100%

^cP_{min} minimal pUC origin (SEQ ID NO: 42) and RNA-OUT (bacterial region = SEQ ID NO: 43) with SV40 enhancer. H1 promoter version (with shRNA and no SV40) is 1035 bp

^dR6K origin and RNA-OUT (bacterial region = SEQ ID NO: 19). H1 promoter version (with shRNA) is 635 bp

^eC2 origin and RNA-OUT (bacterial region = SEQ ID NO: 20). H1 promoter version (with shRNA) is 442 bp (FIG. 11)

Example 6: Alternative RNA Selection Marker Nanoplasmid Vectors

Expression of Nanoplasmid vectors encoding RNA-OUT in the intron (both orientations of RNA-OUT SEQ ID NO:23 inserted into the unique HpaI site in the intron SEQ ID NO:30; NTC9385Ra-O1 dual and NTC9385Ra-O2 dual) demonstrated robust expression with RNA-OUT in either orientation in the intron (Table 9). Consistent with this, similarly high levels of expression are obtained with NTC9385Ra-O1 (SEQ ID NO:40) and NTC9385Ra-O2 (SEQ ID NO:39) which have opposite orientations of intronic RNA-OUT marker and the R6K origin in the spacer region. Nanoplasmid variants with the pMB1 antisense RNA RNAI (SEQ ID NO:31) with promoter and terminator region (RNAI selectable marker: SEQ ID NO:32 flanked by DraIII-KpnI restriction sites for cloning as described previously for RNA-OUT) substituted for RNA-OUT were constructed and tested for expression to determine if alternative selection markers may be utilized in place of RNA-OUT. The results (Table 9) demonstrate alternative RNA selection markers may be substituted for RNA-OUT. Substitution of RNAI for RNA-OUT in the vector backbone (NTC9385Ra-RNAI-O1) or in the intron in either orientation (NTC9385R-RNAI-O1 and NTC9385R-RNAI-O2) did not reduce expression relative to the corresponding RNA-OUT construct. To determine splicing accuracy, NTC9385R-RNAI-O1-EGFP and NTC9385R-RNAI-O2-EGFP were transfected into the A549 cell line and cytoplasmic RNA isolated

strand primer. Intron splicing was determined by PCR amplification of the reverse transcribed cytoplasmic RNA with the exon 1 and exon 2 specific primers. The resultant PCR product (a single band in each case) was determined by sequencing to be the correct spliced exon1-exon2 fragment. This demonstrated that, like intronic RNA-OUT, intronic RNAI in either orientation is accurately removed by splicing and does not interfere with splicing accuracy. This demonstrates that alternative RNA based selection markers could be substituted for RNA-OUT in the spacer region or the intron and that pMB1 RNAI is a preferred RNA based selection marker.

The RNAI transcription unit (SEQ ID NO: 32) may be substituted for the RNA-OUT selection marker (SEQ ID NO: 23) in any of the constructs described in Examples 1-6. Alternatively, the 108 bp RNAI antisense repressor RNA (SEQ ID NO: 31) may be substituted for the 70 bp RNA-OUT antisense repressor RNA (SEQ ID NO: 24) retaining the flanking RNA-OUT transcription control sequences in any of the constructs described in Examples 1-6. RNAI regulated vectors may be grown in RNAII-SacB regulated cell lines further expressing, as required, R6K, Cole2-P9, or Cole2 related rep protein. RNAII-SacB regulated cell lines may be made replacing the RNA-IN sequence in pCAH63-CAT RNA-IN-SacB (P5/6 6/6) with a RNAII target sequence as described in Williams, J A Supra, 2008 included herein by reference. Alternatively, RNAI regulated vectors may be grown in any of the RNAII regulated chromosomal selection marker cell lines disclosed in Grabherr and,

Pfaffenzeller Supra, 2006; Cranenburgh Supra, 2009. These cell lines would be modified for expression, as required, of R6K, Cole2-P9, or Cole2 related rep protein.

Another preferred RNA based selection marker, IncB plasmid RNAI (SEQ ID NO:33; SEQ ID NO:34), is shown in FIG. 12. A cell line for antibiotic free sucrose selection of IncB RNAI expressing plasmid vectors is created by modification of the genomically expressed RNA-IN-SacB cell lines for RNA-OUT plasmid propagation disclosed in Williams, J A Supra, 2008 by replacement of the 68 bp RNA-IN regulator in a PstI-MamI restriction fragment with a 362 bp PstI-MamI IncB RNAII regulator (SEQ ID NO:35). Alternatively, RNA-OUT may be substituted with one of the many RNA based selection markers know in the art.

TABLE 9

High level expression is obtained with pMB1 RNAI or RNA-OUT antisense RNA vectors				
Vector (EGFP)	Spacer region ^a	SR (bp) Intron ^a	A549 FU ^b (T = 48 hr mean + SD)	HEK293 FU ^b (T = 48 hr mean + SD)
NTC8685	T-VA1-BH-P- AF-SV40	1465 HR- β ^c	8546 \pm 1163 (1.0x)	62068 \pm 1760 (1.0x)
NTC8385 (0.85 kb) ^d	T-P _{min} -AF-BE	866 HR- β ^c	9364 \pm 966 (1.10x)	31482 \pm 1822 (0.51x)
NTC9385C (SEQ ID NO: 8)	\leftarrow -C -AF \rightarrow	281 HR- β ^c	8860 \pm 382 (1.04x)	33356 \pm 1489 (0.54x)
NTC9385R (SEQ ID NO: 2)	\leftarrow -R -AF \rightarrow	466 HR- β ^c	16237 \pm 2520 (1.90x)	55919 \pm 6371 (0.90x)
NTC9385Ra- O2 (SEQ ID NO: 39)	\leftarrow -R	306 HR- \leftarrow AF- β	14510 \pm 835 (1.70x)	49526 \pm 2179 (0.80x)
NTC9385Ra- O1 dual	\leftarrow -R -AF \rightarrow	466 HR-AF \rightarrow - β	13929 \pm 1291 (1.63x)	56552 \pm 2714 (0.91x)
NTC9385Ra- O2 dual	\leftarrow -R -AF \rightarrow	466 HR- \leftarrow AF- β	12543 \pm 245 (1.47x)	54379 \pm 1244 (0.89x)
NTC9385Ra- RNAI-O1	\leftarrow -R -RNAI \rightarrow	488 HR-AF \rightarrow - β	15773 \pm 238 (1.85x)	55468 \pm 6619 (0.89x)
NTC9385R- RNAI-O1	\leftarrow -R -AF \rightarrow	466 HR- \leftarrow RNAI - β	14296 \pm 287 (1.67x)	60630 \pm 2176 (0.98x)
NTC9385R- RNAI-O2	\leftarrow -R -AF \rightarrow	466 HR- RNAI \rightarrow - β	12271 \pm 466 (1.44x)	60691 \pm 6482 (0.98x)

^a tnpA term = T; HTLV-IR = HR; B globin 3' acceptor site = β ; RNA-OUT sucrose selection marker = AF; pUC origin RNAI antisense RNA = RNAI; pUC origin = P; R6K origin = R; Cole2 origin = C; BH = PAS-BH UP = upstream pUC plasmid derived DNA.

^b EGFP plasmid DNA transfected with Lipofectamine 2000. Fluorescence units (FU) reported. Mean FU standardized to NTC8685

^c HR β intron is 225 bp

^d P_{min} minimal pUC origin (SEQ ID NO: 42) and RNA-OUT (bacterial region = SEQ ID NO: 43)

Thus, the reader will see that the improved expression vectors of the invention provide for a rational approach to improve plasmid expression.

While the above description contains many examples, these should not be construed as limitations on the scope of the invention, but rather should be viewed as an exemplification of preferred embodiments thereof. Many other variations are possible. For example, the RNA-OUT selectable marker may be substituted with an alternative RNA-OUT sequence variant that functionally binds RNA-IN to repress expression, for example, a CpG free RNA-OUT (SEQ ID NO:36). A CpG free R6K-RNA-OUT bacterial region (SEQ ID NO:37) or CpG free Cole2-RNA-OUT bacterial region (SEQ ID NO: 38) may be utilized. Likewise, the RNA-OUT promoter and/or terminator could be substituted with an alternative promoter and/or terminator. Likewise, the Cole2-P9 or R6K replication origin may be substituted with a Cole2 related replication origin, and propagated in a strain expressing the Cole2 related replication origin replication protein. Likewise, the Cole2-P9 or R6K origin may be substituted with an origin from one of

the numerous additional Rep protein dependent plasmids that are know in the art, for example the Rep protein dependent plasmids described in del Solar et al Supra, 1998 which is included herein by reference. Likewise, the vectors may encode a diversity of transgenes different from the examples provided herein, for example, antigen genes for a variety of pathogens, or therapeutic genes such as hypoxia inducible factor, keratinocyte growth factor, factor IX, factor VIII, etc, or RNA genes such as microRNAs or shRNA. Likewise, the eukaryotic region may express RNA from a RNA Pol III promoter as described herein. The orientation of the various vector-encoded elements may be changed relative to each other. The vectors may optionally contain additional functionalities, such as nuclear localizing

sequences, and/or immunostimulatory RNA elements as disclosed in Williams, Supra, 2008. The vectors may include a boundary element between the bacterial region and the eukaryotic region, for example, the CMV promoter boundary element upstream of the CMV enhancer (or heterologous promoter enhancer) may be included in the vector design (e.g. NTC9385R-BE; SEQ ID NO: 41). The vectors may include a eukaryotic transcriptional terminator between the bacterial region and the eukaryotic region, for example, the 4xC2 terminator or the gastrin terminator. Likewise, the vectors may utilize a diversity of RNA Pol II promoters different from the CMV promoter examples provided herein, for example, constitutive promoters such as the elongation factor 1 (EF1) promoter, the chicken β -actin promoter, the β -actin promoter from other species, the elongation factor-1 α (EF1 α) promoter, the phosphoglycerokinase (PGK) promoter, the Rous sarcoma virus (RSV) promoter, the human serum albumin (SA) promoter, the α -1 antitrypsin (AAT) promoter, the thyroxine binding globulin (TBG) promoter, the cytochrome P450 2E1 (CYP2E1) promoter, etc. The vectors may also utilize combination promoters such as the

chicken β -actin/CMV enhancer (CAG) promoter, the human or murine CMV-derived enhancer elements combined with the elongation factor 1 α (EF1 α) promoters, CpG free versions of the human or murine CMV-derived enhancer elements combined with the elongation factor 1 α (EF1 α) promoters, the albumin promoter combined with an α -feto-protein MERII enhancer, etc, or the diversity of tissue

specific or inducible promoters known in the art such as the muscle specific promoters muscle creatine kinase (MCK), and C5-12 described herein or the liver-specific promoter apolipoprotein A-I (ApoAI).

Accordingly, the scope of the invention should be determined not by the embodiments illustrated, but by the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 44

<210> SEQ ID NO 1

<211> LENGTH: 281

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: R6K gamma Origin

<400> SEQUENCE: 1

```

ggcttggtgt ccacaaccgt taaaccttaa aagctttaa agccttatat attctttttt    60
ttcttataaa acttaaaacc ttagaggcta ttaagttgc tgatttatat taattttatt    120
gttcaaacat gagagcttag tacgtgaaac atgagagctt agtacgtag ccatgagagc    180
ttagtacggt agccatgagg gtttagttcg ttaaacatga gagcttagta cgtaaacaat    240
gagagcttag tacgtactat caacagggtg aactgctgat c                          281

```

<210> SEQ ID NO 2

<211> LENGTH: 1667

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: NTC9385R vector backbone

<400> SEQUENCE: 2

```

ccgcctaag agcgggcttt tttttggctt gttgtccaca accgttaaac cttaaaagct    60
ttaaaagcct tatatattct ttttttctt ataaaactta aaaccttaga ggctatttaa    120
gttgctgatt tatattaatt ttattgttca aacatgagag cttagtagct gaaacatgag    180
agcttagtac gttagccatg agagcttagt acgttagcca tgagggttta gttcgttaaa    240
catgagagct tagtacgta aacatgagag cttagtagct actatcaaca gggtgaactg    300
ctgatccacg ttgtgtaga attggtaaag agagtcgtgt aaaatatcga gttcgcacat    360
cttggtgtct gattattgat ttttggcga accatattgat catatgaca gatgtgtatc    420
taccttaact taatgatttt gataaaaatc attaggtagc ccggctctag ttattaatag    480
taatcaatta cggggtcatt agttcatagc ccatatatgg agttccgctg tacataactt    540
acggtaaag gcccgcctgg ctgaccgccc aacgaccccc gccattgac gtcaataatg    600
acgtatgttc ccatagtaac gccaataggg actttccatt gacgtcaatg ggtggagtat    660
ttacggtaaa ctgccactt ggcagtagat caagtgtatc atatgccaa tagccccct    720
attgacgtca atgacggtaa atggcccggc tggcattatg cccagtagat gaccttatgg    780
gactttccta cttggcagta catctacgta ttagtcatcg ctattaccat ggtgatgcgg    840
ttttggcagt acatcaatgg gcgtggatag cggtttgact cacggggatt tccaagtctc    900
caccatttg acgtcaatgg gagtttgttt tggcaccaaa atcaacggga ctttccaaaa    960
tgtcgtaaca actccgcccc attgacgcaa atgggaggta ggcgtgtacg gtgggaggtc   1020
tatataagca gagctcggtt agtgaaccgt cagatcgctt ggagacgcca tccacgctgt   1080

```


-continued

```

tttgacctcc atagaagaca cgggaccga tccagcctcc gcggctcgca tctctccttc 1140
acgcgcccgc cgccctacct gaggcgcgca tccacgcggg ttgagtcgcg ttctgcccgc 1200
tcccgcctgt ggtgcctcct gaactgcgtc cgccgtctag gtaagttaa agctcaggtc 1260
gagaccgggc ctttgtccgg cgctcccttg gacccctacct agactcagcc ggctctccac 1320
gctttgectg accctgcttg ctcaactcta gttctctcgt taacttaatg agacagatag 1380
aaactggtct tgtagaaaca gagtagtcgc ctgcttttct gccagggtgct gacttctctc 1440
ccctgggctt ttttcttttt ctcagggtga aaagaagaag acgaagaaga cgaagaagac 1500
aaaccgctcg cgacagatct ttttccctct gccaaaatt atggggacat catgaagccc 1560
cttgagcadc tgacttctgg ctaataaagg aaatttattt tcattgcaat agtgtgttgg 1620
aatTTTTTgt gtctctcact cggaaggaca taagggcggc cgctagc 1667

```

<210> SEQ ID NO 3

<211> LENGTH: 2103

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: NTC9685R
vector backbone

<400> SEQUENCE: 3

```

ccgcctaatg agcgggcttt tttttggctt gttgtccaca accgttaaac cttaaaagct 60
ttaaagcct tatatattct ttttttctt ataaaactta aaaccttaga ggctatttaa 120
gttgctgatt tatattaatt ttattgttca aacatgagag cttagtagct gaaacatgag 180
agcttagtac gttagccatg agagcttagt acgttagcca tgagggttta gttcgtaaa 240
catgagagct tagtagctta aacatgagag cttagtagct actatcaaca ggttgaactg 300
ctgatccacg ttgtggtaga attggtaaag agagtcgtgt aaaatatcga gttcgcacat 360
cttgttgtct gattattgat ttttggcgaa accatttgat catatgaaa gatgtgtatc 420
taccttaact taatgatttt gataaaaatc attaggtacc cctgatcact gtggaatgtg 480
tgtcagttag ggtgtggaaa gtccccaggc tccccagcag gcagaagtat gcaaagcatg 540
catctcaatt agtcagcaac cagggtgtga aagtccccag gctccccagc aggcagaagt 600
atgcaaagca tgcactctca ttagtcagca accatagctc cgcccctaac tccgcccctc 660
ccgcccctaa ctccgcccag ttacggggtc attagttcat agcccatata tggagttccg 720
cgttacataa cttacggtaa atggcccgc tggtgaccg cccaacgacc cccgcccatt 780
gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc attgacgtca 840
atgggtggag tatttacggt aaactgccc cttggcagta catcaagtgt atcatatgcc 900
aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgcccagta 960
catgacctta tgggactttc ctacttgcca gtacatctac gtattagtca tcgctattac 1020
catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggtttg actcacgggg 1080
atttccaagt ctccaccca ttgacgtcaa tgggagtttg ttttggcacc aaaatcaacg 1140
ggactttcca aaatgctgta acaactccgc cccattgacg caaatggcg gtaggcgtgt 1200
acggtgggag gtctatataa gcagagctcg ttagtgaaac cgtcagatcg cctggagacg 1260
ccatccacgc tgttttgacc tccatagaag acaccgggac cgatccagcc tccgcccctc 1320
gcatctctcc ttcacgcgcc cgccgcccta cctgagggcg ccatccacgc cggttgagtc 1380
gcgttctgcc gcctcccgcc tgtggtgct cctgaactgc gtccgcccgc taggtaagtt 1440

```

-continued

```

taaagctcag gtcgagaccg ggcctttgtc cggcgctccc ttggagccta cctagactca 1500
gccggctctc cacgctttgc ctgaccctgc ttgetcaact ctagttctct cgtaaactta 1560
atgagacaga tagaaactgg tctttagtaa acagagtagt cgctgcttt tctgccaggt 1620
gctgacttct ctcccctggg cttttttctt tttctcaggt tgaaaagaag aagacgaaga 1680
agacgaagaa gacaaaccgt cgtcgacaga tctttttccc tctgccaaaa attatgggga 1740
catcatgaag ccccttgagc atctgacttc tggctaataa aggaaattta ttttcattgc 1800
aatagtgtgt tggattttt tgtgtctctc actcggagg acataagggc ggccgctagc 1860
ccgcctaatg agcgggcttt tttttcttag ggtgcaaaag gagagcctgt aagcgggcac 1920
tcttccgtgg tctggtggat aaattcgcaa gggatcatg gcggacgacc ggggttcgag 1980
ccccgtatcc ggccgtccgc cgtgatccat gcggttaccg cccgcgtgtc gaaccaggt 2040
gtgcgacgtc agacaacggg ggagtgtctc ttttgcttc cttcccctac cggggcgcgt 2100
agc 2103

```

```

<210> SEQ ID NO 4
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Cole2
Origin (+7)

```

```

<400> SEQUENCE: 4

```

```

caaaagggcg ctgttatctg ataaggctta tctggtctca ttttg 45

```

```

<210> SEQ ID NO 5
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Cole2
Origin (+7, CpG free)

```

```

<400> SEQUENCE: 5

```

```

caaaaggggg ctgttatctg ataaggctta tctggtctca ttttg 45

```

```

<210> SEQ ID NO 6
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Cole2
Origin (Min)

```

```

<400> SEQUENCE: 6

```

```

ggcgctgtta tctgataagg cttatctggt ctcathtt 38

```

```

<210> SEQ ID NO 7
<211> LENGTH: 61
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Cole2
Origin (+16)

```

```

<400> SEQUENCE: 7

```

```

ctgctcaaaa agacgcaaaa agggcgctgt tatctgataa ggcttatctg gtctcathtt 60

```

```

g 61

```


-continued

```

<210> SEQ ID NO 8
<211> LENGTH: 1482
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: NTC9385C
        vector backbone

<400> SEQUENCE: 8
tacaatggct catgtgaaa aaccattggc agaaaaacac ctgccaacag ttttaccaca      60
attgccactt aaccacaaa agggcgctgt tatctgataa ggcttatctg gtctcatttt      120
gcacgttggt gtagaattgg taaagagagt cgtgtaaaat atcgagttcg cacatcttgt      180
tgtctgatta ttgatttttg gcgaaacat ttgatcatat gacaagatgt gtatctacct      240
taacttaatg attttgataa aatcatttag gtaccccggc tctagttatt aatagtaatc      300
aattacgggg tcattagttc atagcccata tatggagttc cgcgttacat aacttacggt      360
aaatggcccc cctggctgac cgcccaacga ccccccca ttgacgtcaa taatgacgta      420
tgttcccata gtaacgcaa tagggacttt ccattgacgt caatgggtgg agtatttacg      480
gtaaactgcc cacttggcag tacatcaagt gtatcatatg ccaagtacgc ccctattga      540
cgtaaatgac ggtaaatggc ccgcctggca ttatgccag tacatgacct tatgggactt      600
tcctacttgg cagtacatct acgtattagt catcgctatt accatggtga tgcggttttg      660
gcagtacatc aatggcgctg gatagcgggt tgactcacgg ggatttcaa gtctccaccc      720
cattgacgtc aatgggagtt tgttttggca ccaaaatcaa cgggactttc caaaatgtcg      780
taacaactcc gcccattga cgcaaatggg cggtaggcgt gtacggtggg aggtctatat      840
aagcagagct cgttttagtga accgtcagat cgctggaga cgccatccac gctgttttga      900
cctccataga agacaccggg accgatccag cctccgcggc tcgcatctct ccttcacgcg      960
cccgcgccc tacctgagge cgccatccac gccggttgag tcgcgttctg ccgcctcccg     1020
cctgtggtgc ctctgaact gcgtccgccc tctaggtaag tttaaagctc aggtcgagac     1080
cgggcctttg tccggcgtc ccttggagcc tacctagact cagecggctc tccacgcttt     1140
gctgaccct gcttgctcaa ctctagttct ctctgtaact taatgagaca gatagaaact     1200
ggtctttagt aaacagagta gtcgctgct tttctgccag gtgctgactt ctctcccctg     1260
ggcttttttc tttttctcag gttgaaaaga agaagacgaa gaagacgaag aagacaaacc     1320
gtcgtcgaca gatcttttc cctctgcaa aaattatggg gacatcatga agccccttga     1380
gcatctgact tctggctaataaaggaaatt tttttcatt gcaatagtgt gttggaattt     1440
tttgtgtctc tactcggaa ggacataagg gcggccgcta gc                               1482

```

```

<210> SEQ ID NO 9
<211> LENGTH: 1918
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: NTC9685C
        vector backbone

```

```

<400> SEQUENCE: 9
tacaatggct catgtgaaa aaccattggc agaaaaacac ctgccaacag ttttaccaca      60
attgccactt aaccacaaa agggcgctgt tatctgataa ggcttatctg gtctcatttt      120
gcacgttggt gtagaattgg taaagagagt cgtgtaaaat atcgagttcg cacatcttgt      180
tgtctgatta ttgatttttg gcgaaacat ttgatcatat gacaagatgt gtatctacct      240

```

-continued

```

taacttaatg attttgataa aatcattag gtaccctga tcaactgtgga atgtgtgtca 300
gtaggggtgt ggaaagtccc caggctcccc agcaggcaga agtatgcaa gcatgcatct 360
caattagtca gcaaccaggt gtggaaagtc cccaggctcc ccaggcaggca gaagtatgca 420
aagcatgcat ctcaattagt cagcaacat agtcccggcc ctaactccgc ccatcccggc 480
cctaactccg cccagttacg gggtcattag ttcatagccc atatatggag ttccgcgta 540
cataacttac ggtaaatggc ccgcctggct gaccgccccaa cgacccccgc ccattgacgt 600
caataatgac gtatgttccc atagtaacgc caatagggac tttccattga cgtcaatggg 660
tggagtattt acggtaaact gccacttgg cagtacatca agtgtatcat atgccaagta 720
cgccccctat tgacgtcaat gacggtaaact ggcccgcctg gcattatgcc cagtacatga 780
ccttatggga ctttctact tggcagtaca tctacgtatt agtcatcgct attacatgg 840
tgatgcgggt ttggcagtac atcaatgggc gtggatagcg gtttgactca cggggatttc 900
caagtctcca cccattgac gtcaatggga gttgttttg gcacaaaat caacgggact 960
ttccaaaatg tcgtaacaac tccgccccat tgacgcaaat gggcggtagg cgtgtacgg 1020
gggaggctta tataagcaga gctcgtttag tgaaccgtca gatcgctgg agacgccatc 1080
cacgtggtt tgacctccat agaagacacc gggaccgatc cagcctccgc ggctcgcatc 1140
tctcttcac gcgcccgcg cctacctga ggccgccatc cagcgggtt gagtcggtt 1200
ctgcccctc ccgctgtgg tgccctcga actgcgtccg ccgtctaggt aagtttaaag 1260
ctcaggctga gaccgggct ttgtccggc ctcccttga gcctacctag actcagccgg 1320
ctctccacgc tttgctgac cctgcttgc caactctagt tctctctga acttaatgag 1380
acagatagaa actggtcttg tagaaacaga gtagtcgct gcttttctgc cagggtgctga 1440
cttctctccc ctgggctttt ttcttttct cagggtgaaa agaagaagac gaagaagacg 1500
aagaagacaa accgtcgtcg acagatcttt ttccctctgc caaaaattat ggggacatca 1560
tgaagcccct tgagcatctg acttctggct aataaaggaa atttattttc attgcaatag 1620
tgtgttgaa tttttgtgt ctctactcg gaaggacata agggcggccg ctagcccgc 1680
taatgagcgg gctttttttt cttagggtgc aaaaggagag cctgtaagcg ggcactcttc 1740
cgtggtctgg tggataaatt cgcaagggtg tcatggcgga cgaccgggt tegagccccg 1800
tatccggccg tccgctgga tccatgcggg taccgcccgc gtgtcgaacc cagggtgtcg 1860
acgtcagaca acgggggagt gctccttttg gcttcttcc cctaccgggg ccgctagc 1918

```

```

<210> SEQ ID NO 10
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: PL Promoter
(-35 to -10)

```

```

<400> SEQUENCE: 10

```

```

ttgacataaa taccactggc ggtgatact 29

```

```

<210> SEQ ID NO 11
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: PL Promoter
OL1-G (-35 to -10)

```


-continued

<400> SEQUENCE: 11

ttgacataaa taccactggc gtgatact

28

<210> SEQ ID NO 12

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: PL Promoter
OL1-G to T(-35 to -10)

<400> SEQUENCE: 12

ttgacataaa taccactggc gttgatact

29

<210> SEQ ID NO 13

<211> LENGTH: 305

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: R6K Rep
protein P42L-P113S

<400> SEQUENCE: 13

Met Arg Leu Lys Val Met Met Asp Val Asn Lys Lys Thr Lys Ile Arg
1 5 10 15His Arg Asn Glu Leu Asn His Thr Leu Ala Gln Leu Pro Leu Pro Ala
20 25 30Lys Arg Val Met Tyr Met Ala Leu Ala Leu Ile Asp Ser Lys Glu Pro
35 40 45Leu Glu Arg Gly Arg Val Phe Lys Ile Arg Ala Glu Asp Leu Ala Ala
50 55 60Leu Ala Lys Ile Thr Pro Ser Leu Ala Tyr Arg Gln Leu Lys Glu Gly
65 70 75 80Gly Lys Leu Leu Gly Ala Ser Lys Ile Ser Leu Arg Gly Asp Asp Ile
85 90 95Ile Ala Leu Ala Lys Glu Leu Asn Leu Pro Phe Thr Ala Lys Asn Ser
100 105 110Ser Glu Glu Leu Asp Leu Asn Ile Ile Glu Trp Ile Ala Tyr Ser Asn
115 120 125Asp Glu Gly Tyr Leu Ser Leu Lys Phe Thr Arg Thr Ile Glu Pro Tyr
130 135 140Ile Ser Ser Leu Ile Gly Lys Lys Asn Lys Phe Thr Thr Gln Leu Leu
145 150 155 160Thr Ala Ser Leu Arg Leu Ser Ser Gln Tyr Ser Ser Ser Leu Tyr Gln
165 170 175Leu Ile Arg Lys His Tyr Ser Asn Phe Lys Lys Lys Asn Tyr Phe Ile
180 185 190Ile Ser Val Asp Glu Leu Lys Glu Glu Leu Ile Ala Tyr Thr Phe Asp
195 200 205Lys Asp Gly Asn Ile Glu Tyr Lys Tyr Pro Asp Phe Pro Ile Phe Lys
210 215 220Arg Asp Val Leu Asn Lys Ala Ile Ala Glu Ile Lys Lys Lys Thr Glu
225 230 235 240Ile Ser Phe Val Gly Phe Thr Val His Glu Lys Glu Gly Arg Lys Ile
245 250 255Ser Lys Leu Lys Phe Glu Phe Val Val Asp Glu Asp Glu Phe Ser Gly
260 265 270

-continued

Asp Lys Asp Asp Glu Ala Phe Phe Met Asn Leu Ser Glu Ala Asp Ala
275 280 285

Ala Phe Leu Lys Val Phe Asp Glu Thr Val Pro Pro Lys Lys Ala Lys
290 295 300

Gly
305

<210> SEQ ID NO 14

<211> LENGTH: 305

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: R6K Rep
protein P42L-P106L-F107S

<400> SEQUENCE: 14

Met Arg Leu Lys Val Met Met Asp Val Asn Lys Lys Thr Lys Ile Arg
1 5 10 15

His Arg Asn Glu Leu Asn His Thr Leu Ala Gln Leu Pro Leu Pro Ala
20 25 30

Lys Arg Val Met Tyr Met Ala Leu Ala Leu Ile Asp Ser Lys Glu Pro
35 40 45

Leu Glu Arg Gly Arg Val Phe Lys Ile Arg Ala Glu Asp Leu Ala Ala
50 55 60

Leu Ala Lys Ile Thr Pro Ser Leu Ala Tyr Arg Gln Leu Lys Glu Gly
65 70 75 80

Gly Lys Leu Leu Gly Ala Ser Lys Ile Ser Leu Arg Gly Asp Asp Ile
85 90 95

Ile Ala Leu Ala Lys Glu Leu Asn Leu Leu Ser Thr Ala Lys Asn Ser
100 105 110

Pro Glu Glu Leu Asp Leu Asn Ile Ile Glu Trp Ile Ala Tyr Ser Asn
115 120 125

Asp Glu Gly Tyr Leu Ser Leu Lys Phe Thr Arg Thr Ile Glu Pro Tyr
130 135 140

Ile Ser Ser Leu Ile Gly Lys Lys Asn Lys Phe Thr Thr Gln Leu Leu
145 150 155 160

Thr Ala Ser Leu Arg Leu Ser Ser Gln Tyr Ser Ser Ser Leu Tyr Gln
165 170 175

Leu Ile Arg Lys His Tyr Ser Asn Phe Lys Lys Lys Asn Tyr Phe Ile
180 185 190

Ile Ser Val Asp Glu Leu Lys Glu Glu Leu Ile Ala Tyr Thr Phe Asp
195 200 205

Lys Asp Gly Asn Ile Glu Tyr Lys Tyr Pro Asp Phe Pro Ile Phe Lys
210 215 220

Arg Asp Val Leu Asn Lys Ala Ile Ala Glu Ile Lys Lys Lys Thr Glu
225 230 235 240

Ile Ser Phe Val Gly Phe Thr Val His Glu Lys Glu Gly Arg Lys Ile
245 250 255

Ser Lys Leu Lys Phe Glu Phe Val Val Asp Glu Asp Glu Phe Ser Gly
260 265 270

Asp Lys Asp Asp Glu Ala Phe Phe Met Asn Leu Ser Glu Ala Asp Ala
275 280 285

Ala Phe Leu Lys Val Phe Asp Glu Thr Val Pro Pro Lys Lys Ala Lys
290 295 300

Gly
305

-continued

<210> SEQ ID NO 15
 <211> LENGTH: 297
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Cole2 Rep
 protein (wild type)

<400> SEQUENCE: 15

```

Met Ser Ala Val Leu Gln Arg Phe Arg Glu Lys Leu Pro His Lys Pro
1           5           10           15

Tyr Cys Thr Asn Asp Phe Ala Tyr Gly Val Arg Ile Leu Pro Lys Asn
          20           25           30

Ile Ala Ile Leu Ala Arg Phe Ile Gln Gln Asn Gln Pro His Ala Leu
          35           40           45

Tyr Trp Leu Pro Phe Asp Val Asp Arg Thr Gly Ala Ser Ile Asp Trp
          50           55           60

Ser Asp Arg Asn Cys Pro Ala Pro Asn Ile Thr Val Lys Asn Pro Arg
65           70           75           80

Asn Gly His Ala His Leu Leu Tyr Ala Leu Ala Leu Pro Val Arg Thr
          85           90           95

Ala Pro Asp Ala Ser Ala Ser Ala Leu Arg Tyr Ala Ala Ala Ile Glu
          100          105          110

Arg Ala Leu Cys Glu Lys Leu Gly Ala Asp Val Asn Tyr Ser Gly Leu
          115          120          125

Ile Cys Lys Asn Pro Cys His Pro Glu Trp Gln Glu Val Glu Trp Arg
          130          135          140

Glu Glu Pro Tyr Thr Leu Asp Glu Leu Ala Asp Tyr Leu Asp Leu Ser
145          150          155          160

Ala Ser Ala Arg Arg Ser Val Asp Lys Asn Tyr Gly Leu Gly Arg Asn
          165          170          175

Tyr His Leu Phe Glu Lys Val Arg Lys Trp Ala Tyr Arg Ala Ile Arg
          180          185          190

Gln Gly Trp Pro Val Phe Ser Gln Trp Leu Asp Ala Val Ile Gln Arg
          195          200          205

Val Glu Met Tyr Asn Ala Ser Leu Pro Val Pro Leu Ser Pro Ala Glu
210          215          220

Cys Arg Ala Ile Gly Lys Ser Ile Ala Lys Tyr Thr His Arg Lys Phe
225          230          235          240

Ser Pro Glu Gly Phe Ser Ala Val Gln Ala Ala Arg Gly Arg Lys Gly
          245          250          255

Gly Thr Lys Ser Lys Arg Ala Ala Val Pro Thr Ser Ala Arg Ser Leu
          260          265          270

Lys Pro Trp Glu Ala Leu Gly Ile Ser Arg Ala Thr Tyr Tyr Arg Lys
          275          280          285

Leu Lys Cys Asp Pro Asp Leu Ala Lys
          290          295

```

<210> SEQ ID NO 16
 <211> LENGTH: 297
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Cole2 Rep
 protein mut (G194D)

<400> SEQUENCE: 16

-continued

Met Ser Ala Val Leu Gln Arg Phe Arg Glu Lys Leu Pro His Lys Pro
 1 5 10 15
 Tyr Cys Thr Asn Asp Phe Ala Tyr Gly Val Arg Ile Leu Pro Lys Asn
 20 25 30
 Ile Ala Ile Leu Ala Arg Phe Ile Gln Gln Asn Gln Pro His Ala Leu
 35 40 45
 Tyr Trp Leu Pro Phe Asp Val Asp Arg Thr Gly Ala Ser Ile Asp Trp
 50 55 60
 Ser Asp Arg Asn Cys Pro Ala Pro Asn Ile Thr Val Lys Asn Pro Arg
 65 70 75 80
 Asn Gly His Ala His Leu Leu Tyr Ala Leu Ala Leu Pro Val Arg Thr
 85 90 95
 Ala Pro Asp Ala Ser Ala Ser Ala Leu Arg Tyr Ala Ala Ala Ile Glu
 100 105 110
 Arg Ala Leu Cys Glu Lys Leu Gly Ala Asp Val Asn Tyr Ser Gly Leu
 115 120 125
 Ile Cys Lys Asn Pro Cys His Pro Glu Trp Gln Glu Val Glu Trp Arg
 130 135 140
 Glu Glu Pro Tyr Thr Leu Asp Glu Leu Ala Asp Tyr Leu Asp Leu Ser
 145 150 155 160
 Ala Ser Ala Arg Arg Ser Val Asp Lys Asn Tyr Gly Leu Gly Arg Asn
 165 170 175
 Tyr His Leu Phe Glu Lys Val Arg Lys Trp Ala Tyr Arg Ala Ile Arg
 180 185 190
 Gln Asp Trp Pro Val Phe Ser Gln Trp Leu Asp Ala Val Ile Gln Arg
 195 200 205
 Val Glu Met Tyr Asn Ala Ser Leu Pro Val Pro Leu Ser Pro Ala Glu
 210 215 220
 Cys Arg Ala Ile Gly Lys Ser Ile Ala Lys Tyr Thr His Arg Lys Phe
 225 230 235 240
 Ser Pro Glu Gly Phe Ser Ala Val Gln Ala Ala Arg Gly Arg Lys Gly
 245 250 255
 Gly Thr Lys Ser Lys Arg Ala Ala Val Pro Thr Ser Ala Arg Ser Leu
 260 265 270
 Lys Pro Trp Glu Ala Leu Gly Ile Ser Arg Ala Thr Tyr Tyr Arg Lys
 275 280 285
 Leu Lys Cys Asp Pro Asp Leu Ala Lys
 290 295

<210> SEQ ID NO 17

<211> LENGTH: 4706

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: pINT pR pL
 R6K Rep piP42L-P106L-F107S (P3-)

<400> SEQUENCE: 17

ctgcaggtga tgattatcag ccagcagaga ttaaggaaaa cagacaggtt tattgagcgc 60
 ttatctttcc ctttattttt gctgcggtta gtcgcataaa aaccattctt cataattcaa 120
 tccatttact atggtatggt ctgaggggag tgaaaattcc cctaattcga tgaagattct 180
 tgctcaattg ttatcageta tgcgcccacc agaacacctt gccgatcagc caaacgtctc 240
 ttcaggccac tgactagcga taactttccc cacaacggaa caactctcat tgcattggat 300

-continued

cattgggtac	tgtgggttta	gtggttgtaa	aaacacctga	ccgctatccc	tgatcagttt	360
cttgaaggta	aactcatcac	cccccaagtct	ggctatgcag	aaatcacctg	gctcaacagc	420
ctgctcaggg	tcaacgagaa	ttaacattcc	gtcaggaaag	cttggettgg	agcctggtgg	480
tgcggtcatg	gaattacctt	caacctcaag	ccagaatgca	gaatcactgg	cttttttggg	540
tgtgcttacc	catctctccg	catcaccttt	ggtaaagggt	ctaagcttag	gtgagaacat	600
cctgcctga	acatgagaaa	aaacagggta	ctcactactca	cttctaagtg	acggctgcat	660
actaacgct	tcatacatct	cgtagatttc	tctggcgatt	gaagggctaa	attcttcaac	720
gctaactttg	agaatttttg	taagcaatgc	ggcgttataa	gcatttaatg	cattgatgcc	780
attaaataaa	gcaccaacgc	ctgactgccc	catccccatc	ttgtctgcca	cagattcctg	840
ggataagcca	agttcatttt	tctttttttc	ataaattgct	ttaaggcgac	gtgcgtcctc	900
aagctgctct	tgtgttaatg	gtttcttttt	tgtgctcata	cgtaaactct	atcaccgcaa	960
gggataaata	tctaaccaccg	tgcgtggtga	ctattttacc	tctggcggtg	ataatggttg	1020
catgtactaa	ggaggttgta	tggaacaacg	cataaccctg	aaagattatg	caatgcgctt	1080
tgggcaaacc	aagacagcta	aagatctctc	acctaccaa	caatgcccc	ctgcaaaaa	1140
taaattcata	taaaaaacat	acagataacc	atctgcggtg	ataaattatc	tctggcggtg	1200
ttgacataaa	taccactggc	ggtgatactg	agcacatcag	caggacgcac	tgaccaccat	1260
gaaggtgacg	ctcttaaaaa	ttaagccctg	aagaagggca	gcattcaaag	cagaaggctt	1320
tggggtgtgt	gatacgaaac	gaagcattgg	gatccagaag	gagatataca	tatgagactc	1380
aaggtcatga	tggacgtgaa	caaaaaaacg	aaaattcgcc	accgaaacga	gctaaatcac	1440
accctggctc	aacttccttt	gcccgcgaaag	cgagtgatgt	atatggcgct	tgctctcatt	1500
gatagcaaag	aacctcttga	acgagggcga	gttttcaaaa	ttagggctga	agaccttgca	1560
gcgctcgcca	aatcacccc	atcgcttgct	tatcgacaat	taaaagaggg	tggtaaatta	1620
cttggtgcca	gcaaaatttc	gctaagaggg	gatgatataca	ttgctttagc	taaagagctt	1680
aacctgctgt	ctactgctaa	aaactcccct	gaagagttag	atcttaacat	tattgagtgg	1740
atagcttatt	caaatgatga	aggatacttg	tctttaaaat	tcaccagaac	catagaacca	1800
tatatctcta	gccttattgg	gaaaaaaaaat	aaattcacia	cgcaattggt	aacggcaagc	1860
ttacgcttaa	gtagccagta	ttcatcttct	ctttatcaac	ttatcaggaa	gcattactct	1920
aattttaaga	agaaaaatta	ttttattatt	tccgttgatg	agttaaagga	agagttaata	1980
gcttatactt	ttgataaaga	tggaaatatt	gagtacaaat	accctgactt	tcctattttt	2040
aaaagggatg	tgttaaataa	agccattgct	gaaattaaaa	agaaaacaga	aatatcgttt	2100
gttgcttca	ctgttcatga	aaaagaaggg	agaaaaatta	gtaagctgaa	gttcgaattt	2160
gtcgttgatg	aagatgaatt	ttctggcgat	aaagatgatg	aagctttttt	tatgaattta	2220
tctgaagctg	atgcagcttt	tctcaaggta	tttgatgaaa	ccgtacctcc	caaaaaagct	2280
aaggggtgag	aattctcatg	tttgacagct	tatcactgat	cagtgaatta	atggcgatga	2340
cgcatcctca	cgataaatatc	cgggtagggc	caatcacttt	cgtctctact	ccgttacaaa	2400
gcgaggctgg	gtatttcccg	gcctttctgt	tatccgaaat	ccactgaaag	cacagcggct	2460
ggctgaggag	ataaataata	aacgaggggc	tgatgcaca	aagcatcttc	tggtgagtta	2520
agaacgagta	tcgagatggc	acatagcctt	gctcaaattg	gaatcaggtt	tgtgccaata	2580
ccagtagaaa	cagacgaaga	agctagctaa	tgctctgtct	caggtcacta	atactatcta	2640
agtagttgat	tcatagtgac	tggatagtt	gcgttttgtc	gcattatgta	gtctatcatt	2700

-continued

```

taaccacaga ttagtgtaat gcgatgattt ttaagtgatt aatggtattt tgtcatcctt 2760
taggtgaata agttgtatat ttaaaatctc ttaattatc agtaaattaa tgtaagtagg 2820
tcattattag tcaaaataaa atcatttgtc gatttcaatt ttgtcccatg gctaattccc 2880
atgtcagccg ttaagtgttc ctgtgtcact caaaattgct ttgagaggct ctaagggctt 2940
ctcagtgcgt tacatccctg gcttggtgtc cacaaccggt aaaccttaaa agctttaaaa 3000
gccttatata ttctttttt tcttataaaa cttaaaacct tagaggctat ttaagttgct 3060
gatttatatt aattttattg ttcaaactg agagcttagt acgtgaaaca tgagagctta 3120
gtacgtagc catgagagct tagtacgta gccatgaggg ttagttcgt taaacatgag 3180
agcttagtac gttaaactg agagcttagt acgtgaaaca tgagagctta gtacgtacta 3240
tcaacagggt gaactgctga tcttcagatc ctctacgccc gacgcatcgt ggccggatct 3300
tgccgcccgt cggcttgaaac gaattgtag acattatttg ccgactacct tggatgctc 3360
gcctttcacg tagtggaaa attcttccaa ctgatctgcg ccgaggcca agcgatcttc 3420
ttcttgcca agataagcct gtctagcttc aagtatgacg ggctgatact gggccggcag 3480
gcgctccatt gccagtcgg cagcgacatc cttcggcggc attttgccg ttactgctt 3540
gtaccaaagc cgggacaacg taagcactac atttcgctca tcgccagccc agtcgggccc 3600
cgagttccat agcgttaagg tttcatttag cgctcaaat agatcctgt caggaaccgg 3660
atcaaagagt tcctccgccc ctggacctac caaggcaacg ctatgttctc ttgctttgt 3720
cagcaagata gccagatcaa tgcgatcgt ggctggctcg aagatactg caagaatgtc 3780
attgctgctc cattctcaa attgcagttc gcgcttagct ggataacgcc acggaatgat 3840
gtcgtcgtgc acaacaatgg tgacttctac agcgcggaga atctcgtct ctccagggga 3900
agccgaagtt tccaaaaggc cgttgatcaa agctcgcgc gttgtttcat caagccttac 3960
ggtcaccgta accagcaaat caatatcact gtgtggcttc aggccgcat ccaactgcga 4020
gccgtacaaa tgtacggcca gcaacgctcg ttcgagatgg cgctcgtga cgccaactac 4080
ctctgatagt tgagtcgata cttcggcgat caccgcttc ctcatgatgt ttaactttgt 4140
tttagggcga ctgcctgct gcgtaacatc gttgctgctc cataacatca aacatcgacc 4200
cacggcgtaa cgcgcttct gcttgatgc ccgaggcata gactgtacc caaaaaaaca 4260
gtcataacaa gccatgaaaa ccgccactgc gccgttacca ccgctgcgtt cggtaaggt 4320
tctggaccag ttgcgtgagc gcatacgtc cttgcattac agcttacgaa ccgaacaggc 4380
ttatgtccac tgggttcgtg ccttcatccg tatcgatgg ccccgatggt agtgtgggg 4440
ctccccatgc gagagtagg aactgccagg catcaaataa aacgaaaggc tcagtcgaaa 4500
gactgggcct ttcgttttat ctggtgttg tcggtgaacg ctctcctgag taggacaaat 4560
ccgccgggag cggatttgaa cgttgcaag caacggccc gaggggtggc ggcaggacgc 4620
ccgccataaa ctgccaggca tcaaattaag cagaaggcca tcctgacgga tggcctttt 4680
gcgtggccag tgccaagctt gcatgc 4706

```

<210> SEQ ID NO 18

<211> LENGTH: 4685

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: pINT pR pL
Cole2 Rep protein mut (G194D)

<400> SEQUENCE: 18

-continued

ctgcaggtga	tgattatcag	ccagcagaga	ttaaggaaaa	cagacaggtt	tattgagcgc	60
ttatctttcc	ctttatTTTT	gctgCGGtaa	gtcgcataaa	aaccattctt	cataattcaa	120
tccatttact	atgTtatgTt	ctgaggggag	tgaaaattcc	cctaattcga	tgaagattct	180
tgctcaattg	ttatcagcta	tgCGccgacc	agaacacctt	gCGatcagc	caaacgtctc	240
ttcaggccac	tgactagcga	taactttccc	cacaacggaa	caactctcat	tgcatgggat	300
cattgggtac	tgtgggttta	gtggTtgtaa	aaacacctga	ccgctatccc	tgatcagttt	360
cttgaaggta	aactcatcac	ccccaaGTct	ggctatgcag	aaatcacctg	gctcaacagc	420
ctgctcaggg	tcaacgagaa	ttaacattcc	gtcaggaaaG	cttggcttgg	agcctgttgg	480
tgCGgtcatg	gaattacctt	caacctcaag	ccagaatgca	gaatcactgg	cttttttggt	540
tgtgcttacc	catctctccg	catcaccttt	ggtaaaggTt	ctaagcttag	gtgagaacat	600
ccctgctga	acatgagaaa	aaacagggta	ctcactactca	cttctaagtG	acggctgcat	660
actaacCGct	tcatacatct	cgtagatttc	tctggcgatt	gaaggGctaa	attcttcaac	720
gctaactttg	agaatTTTTg	taagcaatgc	ggCGttataa	gcatttaatg	cattgatgcc	780
attaaataaa	gcaccaacgc	ctgactgccc	catccccatc	ttgtctgCGa	cagattcctg	840
ggataagcca	agttcatttt	tctttttttc	ataaattgct	ttaaggCGac	gtgCGtctc	900
aagctgctct	tgtgttaatg	gtttctTTTT	tgtgctcata	cgTtaaatct	atcaccgcaa	960
gggataaata	tctaacaccg	tgCGtgTtga	ctattttacc	tctggCGgtg	ataatggTtg	1020
catgtactaa	ggaggttgta	tggaacaacg	cataaccctg	aaagattatg	caatgCGctt	1080
tggGcaaacc	aagacagcta	aagatctctc	acctaccaa	caatgcccc	ctgcaaaaa	1140
taaattcata	taaaaaacat	acagataacc	atctgCGgtg	ataaattatc	tctggCGgtg	1200
ttgacataaa	taccactggc	ggTgatactg	agcacatcag	caggacgcac	tgaccaccat	1260
gaaggtgacg	ctcttaaaaa	ttaagccctg	aagaagggca	gcattcaaag	cagaaggctt	1320
tggggTgtgt	gatacgaaac	gaagcattgg	gatccagaag	gagatataca	tatgagtgcc	1380
gtacttcagc	gcttcagggg	aaaattaccg	cacaaaccgt	actgtacgaa	cgatttcgCG	1440
tacggCGttc	gcattctgCC	gaaaaacatt	gccattcttg	cccgtttcat	ccagcagaac	1500
cagccacatg	cactgtactg	gcttcccttt	gacgtggacc	ggacgggggc	atcaatcgac	1560
tggagCGacc	ggaattgtcc	ggccccgaac	atcacCGtaa	aaaatccccg	taacgggcac	1620
gCGcatctgc	tctacCGct	cgccttccct	gtgagaactg	cgcCGgatgc	atCGgcttcg	1680
gCGctcagat	acgctgcccG	tattgagcgt	gCGTtTgtg	aaaaactggg	cCGggatgtg	1740
aattacagCG	gcctgatctg	caaaaatccg	TGCCaccctg	aatggcagga	agtggaatgg	1800
cCGgaggaac	cctacactct	cGacgaactg	gctgattatc	tCGatttgag	cgcctcagCG	1860
cCGcgtagCG	tCGataaaaa	Ttacgggctg	ggcGaaaact	accatctgtt	cGaaaaggTc	1920
cGtaaattggg	cctacagagc	gattcGtcag	gactggcctg	tattctcaca	atggcttgat	1980
gCGgtgatcc	agcgtgtcga	aatgtacaac	gcatcGcttc	cCGttccgct	ttctccggct	2040
gaatgtcggg	ctattggcaa	gagcattgCG	aaatatacac	acaggaaatt	ctcaccagag	2100
ggattttccg	ctgtacaggc	cGctcGaggt	cGcaagggCG	gaactaaatc	taagcCGcga	2160
gcagttccta	catcagcagc	ttcGctgaag	cCGTgggagg	cattagggat	cagtcGagCG	2220
acgtactacc	gaaaattaaa	atgtgatcca	gatctcGcaa	aatgataaga	attctcatgt	2280
ttgacagctt	atcactgatc	agtgaattaa	TggcGatgac	gcacctcac	gataatatcc	2340

-continued

ggtaggcgc aatcactttc gtctctactc cgttacaaag cgaggctggg tatttcccgg	2400
cctttctggt atccgaaatc cactgaaagc acagcggctg gctgaggaga taaataataa	2460
acgaggggct gtatgcacaa agcatcttct gttgagttaa gaacgagtat cgagatggca	2520
catagccttg ctcaaattgg aatcaggttt gtgccaatac cagtagaaac agacgaagaa	2580
gctagctaat gctctgtctc aggtcactaa tactatctaa gtagttgatt catagtgact	2640
ggatatggtg cgttttgtcg cattatgtag tctatcattt aaccacagat tagtgtaatg	2700
cgatgatatt taagtgatta atgttatttt gtcatccttt aggtgaataa gttgtatatt	2760
taaaatctct ttaattatca gtaaattaat gtaagtaggt cattattagt caaaataaaa	2820
tcatattgtc atttcaattt tgtcccatgg ctaattccca tgtcagccgt taagtgttcc	2880
tgtgtcactc aaaattgctt tgagaggctc taagggcttc tcagtgcgtt acatccctgg	2940
cttgttgctc acaaccgta aaccttaaaa gctttaaag ccttatatat tctttttttt	3000
cttataaaac ttaaaacctt agaggctatt taagttgctg atttatatta attttattgt	3060
tcaaactga gagcttagta cgtgaaacat gagagcttag tacgtagcc atgagagctt	3120
agtacgtag ccatgagggt ttagttcgtt aaacatgaga gcttagtacg ttaaactga	3180
gagcttagta cgtgaaacat gagagcttag tacgtactat caacagggtg aactgctgat	3240
cttcagatcc tctacgccgg acgcatcgtg gccgatctt gcggccgctc ggcttgaacg	3300
aattgttaga cattatttgc cgactacctt ggtgatctcg cctttcacgt agtggacaaa	3360
ttcttccaac tgatctgccc gcgaggccaa gcgatcttct tcttgtcca gataagcctg	3420
tctagcttca agtatgacgg gctgatactg gccggcagg cgctccattg cccagtcggc	3480
agcgacatcc ttcggcgcga ttttgccggt tactgcccgt taccaaatgc gggacaacgt	3540
aagcactaca tttcgctcat cgcagccca gtcgggcggc gagttccata gcgttaaggt	3600
ttcatttagc gcctcaaata gatcctgttc aggaaccgga tcaaagagtt cctccgccgc	3660
tggacctacc aaggcaacgc tatgttctct tgcttttgtc agcaagatag ccagatcaat	3720
gtcgatcgtg gctggctcga agatacctgc aagaatgtca ttgcgctgcc attctccaaa	3780
ttgcagttcg cgcttagctg gataacgcca cggaatgatg tcgtcgtgca caacaatggt	3840
gacttctaca gcgaggagaa tctcgctctc tccaggggaa gccgaagttt ccaaaaggctc	3900
gttgatcaaa gctcgccgcg ttgtttcatc aagccttacg gtcaccgtaa ccagcaaatc	3960
aatatcactg tgtggcttca ggccgccatc cactgcccag ccgtacaaat gtacggccag	4020
caacgtcggg tcgagatggc gctcgatgac gccaaactacc tctgatagtt gagtcgatac	4080
ttcggcgatc accgcttccc tcatgatggt taactttggt ttagggcgac tgccctgctg	4140
cgtaacatcg ttgctgctcc ataacatcaa acatcgaccc acggcgtaac gcgcttgctg	4200
cttggatgcc cgaggcatag actgtacccc aaaaaaacag tcataacaag ccatgaaaac	4260
cgccactgcg ccgttaccac cgctgcgttc ggtcaagggt ctggaccagt tgcgtgagcg	4320
catacgtac ttgcattaca gcttacgaac cgaacaggct tatgtccact gggttcgtgc	4380
cttcatccgt atcgatggc cccgatggta gtgtggggtc tccccatgag agagtggga	4440
actgccaggc atcaataaaa acgaaaggct cagtcgaaag actgggcctt tcgttttatc	4500
tgttgtttgt cgggtaacgc tctcctgagt aggacaaatc cgccgggagc ggatttgaac	4560
gttgcgaagc aacggcccgg aggggtggcg gcaggacgcc cgccataaac tgccaggcat	4620
caaattaagc agaaggccat cctgacggat ggcctttttg cgtggccagt gccaaagctg	4680
catgc	4685

-continued

<210> SEQ ID NO 19
 <211> LENGTH: 466
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: NTC9385R
 and NTC9685R Bacterial region [NheI site-trpA terminator-R6K
 Origin-RNA-OUT-KpnI site]

<400> SEQUENCE: 19

```

gctagcccg ctaatgagcg ggcttttttt tggtttgttg tccacaaccg ttaaacctta    60
aaagctttaa aagccttata tattcttttt tttcttataa aacttaaaac ctagaggct    120
atttaagttg ctgatttata ttaattttat tgttcaaaca tgagagctta gtacgtgaaa    180
catgagagct tagtacgta gccatgagag cttagtacgt tagccatgag ggtttagttc    240
gttaaacatg agagcttagt acgttaaaca tgagagctta gtacgtacta tcaacaggtt    300
gaactgctga tccacgttgt ggtagaattg gtaaagagag tcgtgtaaaa tatcgagttc    360
gcacatcttg ttgtctgatt attgattttt ggcgaaacca tttgatcata tgacaagatg    420
tgtatctacc ttaacttaat gattttgata aaaatcatta ggtacc                    466
  
```

<210> SEQ ID NO 20
 <211> LENGTH: 281
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: NTC9385C
 and NTC9685C Bacterial region [NheI site-ssiA-Cole2 Origin
 (+7)-RNA-OUT-KpnI site]

<400> SEQUENCE: 20

```

gctagctaca atggctcatg tggaaaaacc attggcagaa aaacacctgc caacagtttt    60
accacaattg ccacttaacc cacaaaaggg cgctgttata tgataaggct tatctggctc    120
cattttgcac gttgtgtag aattggtaaa gagagtcgtg taaaatatcg agttcgcaca    180
tcttggtgtc tgattattga tttttggcga aaccatttga tcatatgaca agatgtgtat    240
ctaccttaac ttaatgattt tgataaaaat cattaggtac c                        281
  
```

<210> SEQ ID NO 21
 <211> LENGTH: 76
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: NTC9385C
 and NTC9685C CpG free ssiA [from plasmid R6K]

<400> SEQUENCE: 21

```

tacaatggct catgtggaaa aaccattggc agaaaaacac ctgccaacag ttttaccaca    60
attgccactt aaccca                                             76
  
```

<210> SEQ ID NO 22
 <211> LENGTH: 260
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: CpG free
 R6K origin

<400> SEQUENCE: 22

```

aaaccttaaa accttataaa gccttatata ttcttttttt tcttataaaa cttaaacct    60
  
```

-continued

tagaggctat ttaagttgct gatttatatt aattttattg ttcaaactg agagcttagt	120
acatgaaaca tgagagctta gtacattagc catgagagct tagtacatta gccatgaggg	180
tttagttcat taaacatgag agcttagtac attaaacatg agagcttagt acatactatc	240
aacaggttga actgctgatc	260

<210> SEQ ID NO 23
 <211> LENGTH: 137
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: RNA-OUT
 Selectable Marker from NTC9385C, NTC9685C, NTC9385R, NTC9685R

<400> SEQUENCE: 23

agaattggta aagagagtcg tgtaaaatcg cgagttcgca catcttggtg tctgattatt	60
gatttttggc gaaaccattt gatcatatga caagatgtgt atctacctta acttaatgat	120
tttgataaaa atcatta	137

<210> SEQ ID NO 24
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: RNA-OUT
 Sense strand RNA from NTC9385C, NTC9685C, NTC9385R, NTC9685R,
 NTC9385Ra

<400> SEQUENCE: 24

ttcgcacatc ttgttgctg attattgatt ttggcgaaa ccatttgatc atatgacaag	60
atgtgtatct	70

<210> SEQ ID NO 25
 <211> LENGTH: 87
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: TPA
 secretion sequence

<400> SEQUENCE: 25

atggatgcaa tgaagagagg gctctgctgt gtgctgctgc tgtgtggagc agtcttcggt	60
tgcgccagcg gtaccggatc cgtcgac	87

<210> SEQ ID NO 26
 <211> LENGTH: 57
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: PCR primer
 15061101

<400> SEQUENCE: 26

ggaacgggat ccagaaggag atatacatat gaggccgta cttcagcgt tcagggga	57
---	----

<210> SEQ ID NO 27
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: PCR primer
 15061102

<400> SEQUENCE: 27

-continued

ggaacggaat tcttatcatt ttgcgagatc tggatcacat 40

<210> SEQ ID NO 28
 <211> LENGTH: 32
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Cole2 core
 replication origin
 <400> SEQUENCE: 28

ggcgctgtta tctgataagg cttatctggt ct 32

<210> SEQ ID NO 29
 <211> LENGTH: 121
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 +7(CpG free)-ssiA Cole2 origin
 <400> SEQUENCE: 29

tacaatggct catgtggaaa aaccattggc agaaaaacac ctgccaacag ttttaccaca 60

attgccactt aaccacaaa agggggctgt tatctgataa ggcttatctg gtctcatttt 120

g 121

<210> SEQ ID NO 30
 <211> LENGTH: 228
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: HTLV-
 IR-Rabbit Beta globin hybrid intron
 <400> SEQUENCE: 30

aggtaagttt aaagctcagg tcgagaccgg gcctttgtcc ggcgctccct tggagcctac 60

ctagactcag ccggctctcc acgctttgcc tgaccctgct tgctcaactc tagttctctc 120

gttaacttaa tgagacagat agaaactggc cttgtagaaa cagagtagtc gcttgctttt 180

ctgccagggtg ctgacttctc tcccctgggc tttttctttt ttctcagg 228

<210> SEQ ID NO 31
 <211> LENGTH: 108
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: pMB1 RNAI
 antisense repressor RNA (origin antisense partner of RNAII)
 <400> SEQUENCE: 31

acagtatttg gtatctgcg tctgctgaag ccagttacct tcggaaaaag agttggtagc 60

tcttgatccg gcaaacaaac caccgctggt agcgggtggt tttttggt 108

<210> SEQ ID NO 32
 <211> LENGTH: 156
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: pMB1 RNAI
 selectable Marker, RNAI RNA (Sense strand)
 <400> SEQUENCE: 32

ttgaagtggg ggctaacta cggctacact agaagaacag tatttggtat ctgctgctctg 60

-continued

ctgaagccag ttaccttcgg aaaaagagtt ggtagctctt gatccggcaa acaaaccacc 120
gctggtagcg gtggtttttt tgtttgcaag cagcag 156

<210> SEQ ID NO 33
<211> LENGTH: 73
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: IncB RNAI
antisense repressor RNA (IncB plasmid origin RNAII antisense
partner)

<400> SEQUENCE: 33
gtattctgtg aggccccat tatttttctg cgttccgcca agttcgagga aaaatagtgg 60
gggttttctt tta 73

<210> SEQ ID NO 34
<211> LENGTH: 124
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: IncB RNAI
selectable Marker. DraIII-KpnI restriction fragment.

<400> SEQUENCE: 34
cacgttgtgt tgaatctctg gtacggtttc atatatactt atcccgtatt ctgtgaggcc 60
cccattatth ttctgcggtc cgccaagttc gaggaaaaat agtggggggt ttcctttagg 120
tacc 124

<210> SEQ ID NO 35
<211> LENGTH: 362
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: IncB
RNAII-SacB. PstI-MamI restriction fragment

<400> SEQUENCE: 35
ctgcagttca aagcgggtga aaaggggtat attgcggatc gttattcagt ggcttttggg 60
atcctcgcgg tccgaaagc cagaaaacgg cagaatgcgc cataaggcat tcaggacgta 120
tggcagaaac gacggcagtt tgccggtgcc ggaaggctga aaaaagtttc agaagaccat 180
aaaggaaaac cccactatt tttcctcga cttggcggaa cgcagaaaaa taatgggggc 240
ctcacagaat acgggatagg gcccatgaaa ccgtaccaga gattggggccc tgtgcagtgt 300
ataaatacac ggcacaatcg ctccgccata agcgacagct tgtggcaggt ctgatgaaca 360
tc 362

<210> SEQ ID NO 36
<211> LENGTH: 159
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: CpG free
RNA-OUT selection marker - flanked by KpnI and BglII- EcoRI sites

<400> SEQUENCE: 36
ggtacctggt agaattggtg aagagagttg tgtaaaatat tgagttagca catcttggtg 60
tctgattatt gatttttggg gaaaccattt gatcatatga caagatgtgt atctacctta 120
acttaatgat tttgataaaa atcattaaga tctgaattc 159

-continued

<210> SEQ ID NO 37
 <211> LENGTH: 432
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: CpG free
 R6K gamma - RNA-OUT bacterial region (CpG free R6K origin-CpG free
 RNA-OUT selection marker)-flanked by EcoRI-SphI and BglII-EcoRI
 sites

<400> SEQUENCE: 37

```

gaattcagca tgcaaacctt aaaaccttta aaagccttat atattctttt ttttcttata      60
aaacttaaaa ccttagaggc tatttaagtt gctgatttat attaatttta ttgttcaaac      120
atgagagctt agtacaatgaa acatgagagc ttagtacatt agccatgaga gcttagtaca      180
ttagccatga gggtttagtt cattaaacat gagagccttag tacattaaac atgagagctt      240
agtacatact atcaacaggt tgaactgctg atcggtacct ggtagaattg gtaaagagag      300
ttgtgtaaaa tattgagtta gcacatcttg ttgtctgatt attgattttt ggggaaacca      360
tttgatcata tgacaagatg tgtatctacc ttaacttaat gattttgata aaaatcatta      420
agatctgaat tc                                                                432
  
```

<210> SEQ ID NO 38
 <211> LENGTH: 292
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: CpG free
 Cole2 bacterial region (CpG free ssiA-CpG free Cole2 origin-CpG
 free RNA-OUT selection marker) - - flanked by EcoRI-SphI and
 BglII-EcoRI sites

<400> SEQUENCE: 38

```

gaattcagca tgctacaatg gctcatgtgg aaaaaccatt ggcagaaaaa cacctgccaa      60
cagttttacc acaattgcca cttaacccac aaaagggggc tgttatctga taaggcttat      120
ctggctctcat tttggtacct ggtagaattg gtaaagagag ttgtgtaaaa tattgagtta      180
gcacatcttg ttgtctgatt attgattttt ggggaaacca tttgatcata tgacaagatg      240
tgtatctacc ttaacttaat gattttgata aaaatcatta agatctgaat tc                292
  
```

<210> SEQ ID NO 39
 <211> LENGTH: 1660
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 NTC9385Ra-02 vector backbone

<400> SEQUENCE: 39

```

ccgcctaata agcgggcttt tttttggctt gttgtccaca accgttaaac cttaaaagct      60
ttaaaagcct tatatattct tttttttctt ataaaactta aaaccttaga ggctatttaa      120
gttgctgatt tatattaatt ttattgttca aacatgagag cttagtacgt gaaacatgag      180
agcttagtac gttagccatg agagcttagt acgtagcca tgagggttta gttcgttaaa      240
catgagagct tagtacgta aacatgagag cttagtacgt actatcaaca gggtgaactg      300
ctgatccacc cgggctctag ttattaatag taatcaatta cggggtcatt agttcatagc      360
ccatatatgg agttccgctg tacataactt acggtaaagt gcccgctgg ctgaccgccc      420
aacgaccccc gccattgac gtcaataatg acgtatgttc ccatagtaac gccaataggg      480
  
```

-continued

```

actttccatt gacgtcaatg ggtggagtat ttacggtaaa ctgcccactt ggcagtacat 540
caagtgtatc atatgccaaag tacgccccct attgacgtca atgacggtaa atggccccgcc 600
tggcattatg cccagtacat gaccttatgg gactttccta cttggcagta catctacgta 660
ttagtcatcg ctattaccat ggtgatgegg ttttggcagt acatcaatgg gcgtggatag 720
cggtttgact cacggggatt tccaagtctc caccocattg acgtcaatgg gagtttgttt 780
tggcaccaaaa atcaacggga ctttccaaaa tgtcgtaaca actccgcccc attgacgcaa 840
atggggcggtg ggcgtgtacg gtgggaggtc tatataagca gagctcgttt agtgaaccgt 900
cagatcgect ggagacgcca tccacgctgt tttgacctcc atagaagaca ccgggaccga 960
tccagcctcc gcggtctgca tctctccttc acgcgccccg cgccctacct gagggcgcca 1020
tccacgcggg ttgagtcgcy ttctgcccgc tcccgcctgt ggtgcctcct gaactgcytc 1080
cgccgtctag gtaagttaa agctcaggtc gagaccgggc ctttgtccgg cgctcccttg 1140
gagcctacct agactcagcc ggctctccac gctttgcctg accctgcttg ctcaactcta 1200
gttctctcgt tctaattgatt tttatcaaaa tcattaagtt aaggtagata cacatcttgt 1260
catatgatca aatggtttcg ccaaaaatca ataactcagc aacaagatgt gcgaactcga 1320
tattttacac gactctcttt accaattcta ccacaactta atgagacaga tagaaactgg 1380
tctttagaaa acagagtagt cgctgcttt tctgccaggt gctgacttct ctcccctggg 1440
cttttttctt tttctcaggt tgaaaagaag aagacgaaga agacgaagaa gacaaaaccgt 1500
cgtcgacaga tctttttccc tctgcaaaaa attatgggga catcatgaag ccccttgagc 1560
atctgacttc tggctaataa aggaaattta ttttcattgc aatagtgtgt tggaattttt 1620
tgtgtctctc actcggaagg acataagggc ggccgctagc 1660

```

<210> SEQ ID NO 40

<211> LENGTH: 1660

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:

NTC9385Ra-O1 vector backbone

<400> SEQUENCE: 40

```

ccgcctaatg agcgggcttt tttttggctt gttgtccaca accgttaaac cttaaaagct 60
ttaaaagcct tatatattct tttttttctt ataaaactta aaaccttaga ggctatttaa 120
gttctgatt tatattaatt ttattgttca aacatgagag cttagtacgt gaaacatgag 180
agcttagtac gttagccatg agagcttagt acgttagcca tgagggttta gttcgtaaaa 240
catgagagct tagtacgta aacatgagag cttagtacgt actatcaaca gggtgaactg 300
ctgatccacc ccggctctag ttattaatag taatcaatta cggggtcatt agttcatagc 360
ccatatatgg agttccgctg tacataactt acggtaaatg gcccgctgg ctgaccgccc 420
aacgaccccc gccattgac gtcaataatg acgtatgttc ccatagtaac gccaataggg 480
actttccatt gacgtcaatg ggtggagtat ttacggtaaa ctgcccactt ggcagtacat 540
caagtgtatc atatgccaaag tacgccccct attgacgtca atgacggtaa atggccccgcc 600
tggcattatg cccagtacat gaccttatgg gactttccta cttggcagta catctacgta 660
ttagtcatcg ctattaccat ggtgatgegg ttttggcagt acatcaatgg gcgtggatag 720
cggtttgact cacggggatt tccaagtctc caccocattg acgtcaatgg gagtttgttt 780
tggcaccaaaa atcaacggga ctttccaaaa tgtcgtaaca actccgcccc attgacgcaa 840

```


-continued

```

atgggcggta ggcgtgtacg gtgggaggtc tatataagca gagctcgttt agtgaaccgt    900
cagatcgect ggagacgcca tccacgctgt tttgacctcc atagaagaca ccgggaccga    960
tccagcctcc gcggtctgca tctctccttc acgcgcccgc cgccctacct gaggccgcca   1020
tccacgccgg ttgagtcgcy ttctgcccgc tcccgcctgt ggtgectcct gaactgcgtc   1080
cgccgtctag gtaagttaa agctcaggtc gagaccgggc ctttgtccgg cgctcccttg   1140
gagcctacct agactcagcc ggctctccac gctttgcttg accctgcttg ctcaactcta   1200
gttctctcgt tgtggtagaa ttggtaaaga gagtcgtgta aaatatcgag ttcgcacatc   1260
ttgttgtctg attattgatt tttggcgaaa ccatttgatc atatgacaag atgtgtatct   1320
accttaactt aatgattttg ataaaaatca ttagaactta atgagacaga tagaaactgg   1380
tctttagaaa acagagtagt cgctgcttt tctgccaggt gctgacttct ctcccctggg   1440
cttttttctt tttctcaggt tgaaaagaag aagacgaaga agacgaagaa gacaaaaccgt   1500
cgtcgacaga tctttttccc tctgcaaaaa attatgggga catcatgaag ccccttgagc   1560
atctgacttc tggctaataa aggaaattta ttttcattgc aatagtgtgt tggaaatfff   1620
tgtgtctctc actcggaagg acataagggc ggccgctagc                               1660

```

<210> SEQ ID NO 41

<211> LENGTH: 1763

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: NTC9385R-BE vector backbone

<400> SEQUENCE: 41

```

ccgcctaatg agcgggcttt tttttggctt gttgtccaca accgttaaac cttaaaagct    60
ttaaaagcct tatatattct ttttttctt ataaaactta aaaccttaga ggctatttaa   120
gttgctgatt tatattaatt ttattgttca aacatgagag cttagtagct gaaacatgag   180
agcttagtac gttagccatg agagcttagt acgttagcca tgagggttta gttcgttaaa   240
catgagagct tagtagctta aacatgagag cttagtagct actatcaaca gggtgaactg   300
ctgatccacg ttgtgtaga attggtaaag agagtcgtgt aaaatatcga gttcgcacat   360
cttgttgtct gattattgat ttttggcgaa accatttgat catatgaca gatgtgtatc   420
taccttaact taatgatfff gataaaaatc attaggtacc ccggctctag atggccattg   480
catacgttgt atccatatca taatatgtac atttatattg gctcatgtcc aacattaccg   540
ccatgttgac attgattatt gactagttat taatagtaat caattacggg gtcattagtt   600
catagcccat atatggagtt ccgcgttaca taacttacgg taaatggccc gcctggctga   660
ccgccaacg acccccggcc attgacgtca ataatgacgt atgttcccat agtaacgcca   720
atagggactt tccattgacg tcaatgggtg gagtatttac ggtaaactgc ccacttgcca   780
gtacatcaag tgtatcatat gccaaagtacg ccccctattg acgtcaatga cggtaaatgg   840
cccgcctggc attatgcca gtacatgacc ttatgggact ttctacttg gcagtacatc   900
tacgtattag tcatcgctat taccatgggtg atgcggtfff ggcagtacat caatgggctg   960
ggatagcggg ttgactcacg gggatttcca agtctccacc ccattgacgt caatgggagtt  1020
ttgttttggc accaaaatca acgggacttt ccaaaaatgta gtaacaactc cgccccattg  1080
acgcaaatgg gcggtaggcg tgtacgggtg gaggtctata taagcagagc tcgttttagtg  1140
aaccgtcaga tgcctggag acgcatcca cgctgttttg acctccatag aagacaccgg  1200

```

-continued

```

gaccgatcca gccctccggg ctgcacatctc tccttcacgc gcccgcggcc ctacctgagg 1260
ccgccatcca cgccggttga gtcgcggttct gccgcctccc gcctgtggtg cctcctgaac 1320
tgcgtccgcc gtctaggtaa gtttaaagct caggtcgaga ccgggccttt gtccggcgct 1380
cccttgagac ctacctagac tcagccggct ctccacgctt tgectgacct tgcttgctca 1440
actctagttc tctcgtaac ttaatgagac agatagaaac tggctctgta gaaacagagt 1500
agtcgcctgc tttctgcca ggtgctgact tctctcccct gggctttttt ctttttctca 1560
ggttgaaaag aagaagacga agaagacgaa gaagacaaac cgtcgctgac agatcttttt 1620
ccctctgcca aaaattatgg ggacatcatg aagccccttg agcatctgac ttctggctaa 1680
taaaggaaat ttattttcat tgcaatagtg tgttgaatt ttttgtgtct ctcaactcgga 1740
aggacataag ggcggccgct agc 1763

```

```

<210> SEQ ID NO 42
<211> LENGTH: 678
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Pmin
minimal pUC replication origin

```

```

<400> SEQUENCE: 42

```

```

cgcgttgctg gcgtttttca taggctccgc cccctgacg agcatcacia aaatcgacgc 60
tcaagtcaga ggtggcgaaa cccgacagga ctataaagat accaggcggt tccccctgga 120
agtcacctcg tgcgctctcc tgttccgacc ctgccgctta ccggatacct gtccgccttt 180
ctcccttcgg gaagcgtggc gcttttctcat agctcacgct gtaggtatct cagttcggtg 240
taggtcgttc gctccaagct gggctgtgtg cacgaacccc ccggttcagcc cgaccgctgc 300
gccttatccg gtaactatcg tcttgagtc aaccggtaa gacacgactt atcgccactg 360
gcagcagcca ctggtaacag gattagcaga gcgaggtatg taggcggtgc tacagagttc 420
ttgaagtggg ggcctaacta cggctacact agaagaacag tatttggtat ctgcgctctg 480
ctgaagccag ttaccttcgg aaaaagagtt ggtagctctt gatccggcaa acaaaccacc 540
gctggtagcg gtggtttttt tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct 600
caagaagatc ctttgatctt ttctacgggg tctgacgctc agtggaacga aaactcacgt 660
taagggattt tggatcatg 678

```

```

<210> SEQ ID NO 43
<211> LENGTH: 866
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: pUC (0.85)
Bacterial region [NheI site- trpA terminator-Pmin pUC replication
origin (minimal)-RNA-OUT-KpnI site]

```

```

<400> SEQUENCE: 43

```

```

gctagcccgc ctaatgagcg ggcttttttt tcttaggcct cgcgttgctg gcgtttttca 60
taggctccgc cccctgacg agcatcacia aaatcgacgc tcaagtcaga ggtggcgaaa 120
cccgacagga ctataaagat accaggcggt tccccctgga agtcacctcg tgcgctctcc 180
tgttccgacc ctgccgctta ccggatacct gtccgccttt ctcccttcgg gaagcgtggc 240
gcttttctcat agctcacgct gtaggtatct cagttcggtg taggtcgttc gctccaagct 300
gggctgtgtg cacgaacccc ccggttcagcc cgaccgctgc gccttatccg gtaactatcg 360

```


-continued

```

tcttgagtcc aaccggttaa gacacgactt atcgccactg gcagcagcca ctggtaacag    420
gattagcaga gcgaggtatg taggcggtgc tacagagttc ttgaagtggg ggcctaacta    480
cggctacact agaagaacag tatttggtat ctgcgctctg ctgaagccag ttaccttcgg    540
aaaaagagtt ggtagctctt gatccggcaa acaaaccacc gctggtagcg gtgggttttt    600
tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc ctttgatctt    660
ttctacgggg tctgacgctc agtggaacga aaactcacgt taagggattt tggatcatgg    720
ggtagaattg gtaaagagag tegtgtaaaa tatcgagttc gcacatcttg ttgtctgatt    780
attgattttt ggcgaaacca tttgatcata tgacaagatg tgtatctacc ttaacttaat    840
gattttgata aaaatcatta ggtacc                                     866

```

<210> SEQ ID NO 44

<211> LENGTH: 146

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: PL promoter with locations of the PL promoter OL1, OL2 and OL3 repressor binding sites, -10 and -35 promoter elements for PL1 and PL2 promoters

<400> SEQUENCE: 44

```

agcaattcag atctctcacc taccaaaaa tgccccctg caaaaaataa attcatataa    60
aaaacataca gataaccatc tgcggtgata aattatctct ggcggtgttg acataaatac    120
cactggcggt gatactgagc acatca                                     146

```

What is claimed is:

1. A method for heat inducible production of a Rep protein dependent replication origin plasmid vector, comprising:

- a. cloning the plasmid replication regulating Rep protein into an expression vector to create a P_L -Rep protein expression cassette in which said Rep protein is expressed under the control of the P_L promoter and said P_L promoter further comprises an OL1 mutation in which repressor binding to OL1 is decreased or lost by mutation in OL1, said OL1 mutation comprising either a single base substitution or a single base deletion within the P_L promoter (-35 to -10) SEQ ID NO:10;
- b. integrating said P_L -Rep protein expression cassette into a host strain genome to create a P_L -Rep protein host strain in which expression from the said P_L -Rep protein expression cassette is repressed by a temperature sensitive lambda repressor expressed from the host strain genome;
- c. transforming said P_L -Rep protein host strain with said Rep protein dependent replication origin plasmid vector;
- d. isolating the resultant transformed bacterial cells;
- e. propagating said transformed bacterial cells at 25-32° C. to maintain the said Rep protein dependent plasmid vector at a basal copy number; and
- f. inducing said transformed bacterial cells at 37-42° C. to increase copy number of said Rep protein dependent plasmid vector.

2. The method of claim 1, wherein said P_L promoter OL1 mutation comprising either a single base substitution or a single base deletion within the P_L promoter (-35 to -10) SEQ ID NO:10 is selected from the group consisting of a P_L

35 promoter OL1-G (SEQ ID NO: 11), and a P_L promoter OL1-G to T (SEQ ID NO: 12).

3. The method of claim 1, wherein said Rep protein dependent replication origin is selected from the group consisting of a R6K replication origin, ColE2-P9 replication origin and ColE2 related replication origin.

4. The method of claim 1, wherein said plasmid replication regulating Rep protein comprises an R6K Rep protein mutation selected from the group consisting of P42L-P113S (SEQ ID NO: 13) and P42L-P106L-F107S (SEQ ID NO: 14).

5. The method of claim 1, wherein said plasmid replication regulating Rep protein comprises a ColE2 Rep protein selected from the group consisting of ColE2 Rep protein (SEQ ID NO: 15) and ColE2 Rep protein mutation G194D (SEQ ID NO: 16).

6. The method of claim 1, wherein said P_L -Rep protein host strain further comprises a genomically expressed RNA-IN regulated selection marker.

7. The method of claim 1, wherein said Rep protein dependent plasmid vector for heat inducible production has a vector backbone with at least 95% sequence identity to a sequence selected from the group consisting of: SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 39, SEQ ID NO: 40, and SEQ ID NO: 41.

8. A method for heat inducible production of a Rep protein dependent plasmid vector, comprising:

- a. obtaining bacterial cells transformed with the Rep protein dependent plasmid vector, the bacterial cells comprising a P_L promoter and a Rep protein, wherein the Rep protein dependent plasmid vector comprises a replication origin, wherein the P_L promoter controls the expression of the Rep protein, wherein the P_L promoter

83

comprises the nucleic acid sequence of SEQ ID NO:10, and wherein the P_L promoter comprises an OLI mutation within the nucleic acid sequence of SEQ ID NO: 10, wherein the OLI mutation comprises either a single base substitution or a single base deletion that decreases or prevents repressor binding to OLI;

b. incubating said bacterial cells at 25-32° C. to maintain the Rep protein dependent plasmid vector at a basal copy number; and

c. incubating said bacterial cells at 37-42° C. to increase copy number of said Rep protein dependent plasmid vector.

9. The method of claim 8, wherein said OLI mutation within the nucleic acid sequence of SEQ ID NO:10 is selected from the group consisting of SEQ ID NO: 11 and SEQ ID NO: 12.

10. The method of claim 8, wherein the replication origin is selected from the group consisting of a R6K replication origin, ColE2-P9 replication origin and ColE2 related replication origin.

84

11. The method of claim 8, wherein the replication origin is a R6K replication origin, and wherein said Rep protein comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 13 and SEQ ID NO: 14.

12. The method of claim 8, wherein the replication origin is a ColE2-P9 replication origin, and wherein the Rep protein comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 15 and SEQ ID NO: 16.

13. The method of claim 8, wherein said bacteria cells further comprises a genomically expressed RNA-IN regulated selection marker.

14. The method of claim 8, wherein said Rep protein dependent plasmid vector has a vector backbone with at least 95% sequence identity to a sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 39, SEQ ID NO: 40, and SEQ ID NO: 41.

* * * * *