



US00RE49625E

(19) **United States**
(12) **Reissued Patent**
Sierks et al.

(10) **Patent Number:** **US RE49,625 E**
(45) **Date of Reissued Patent:** **Aug. 29, 2023**

(54) **NANOSCALE PROCESS TO GENERATE REAGENTS SELECTIVE FOR INDIVIDUAL PROTEIN VARIANTS**

(71) Applicant: **Arizona Board of Regents, a Body Corporate of the State of Arizona, Acting For and on Behalf of Arizona State University, Scottsdale, AZ (US)**

(72) Inventors: **Michael Sierks, Ft. McDowell, AZ (US); Mark Hayes, Gilbert, AZ (US)**

(73) Assignee: **ARIZONA BOARD OF REGENTS, A BODY CORPORATE OF THE STATE OF ARIZONA, Scottsdale, AZ (US), ACTING FOR AND ON BEHALF OF ARIZONA STATE UNIVESITY**

(21) Appl. No.: **16/845,761**

(22) Filed: **Apr. 10, 2020**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **9,938,330**
Issued: **Apr. 10, 2018**
Appl. No.: **14/388,209**
Filed: **Sep. 25, 2014**

U.S. Applications:

(60) Provisional application No. 61/617,476, filed on Mar. 29, 2012.

(51) **Int. Cl.**
C07K 14/47 (2006.01)
B03C 5/00 (2006.01)
(Continued)

(52) **U.S. Cl.**
CPC **C07K 14/4711** (2013.01); **B03C 5/005** (2013.01); **B03C 5/026** (2013.01);
(Continued)

(58) **Field of Classification Search**
CPC ... **B03C 5/026**; **G01N 27/447**; **C07K 14/4711**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,014,747 B2 3/2006 Cummings et al.
7,347,923 B2 3/2008 Cummings et al.
(Continued)

FOREIGN PATENT DOCUMENTS

WO 1994010205 A1 5/1994
WO 2004013172 A2 2/2004
(Continued)

OTHER PUBLICATIONS

Nakano. Electrophoresis. Sep. 2011; 32(17): 2314-2322 (Year: 2011).*

(Continued)


Primary Examiner — Padmashri Ponnaluri
Assistant Examiner — Jean C Witz
(74) *Attorney, Agent, or Firm* — Duane Morris LLP

(57) **ABSTRACT**

The present invention provides devices and methods to separate and concentrate target protein species at a micro-liter scale and to generate reagents to those variants with exquisite selectivity for specific protein isoforms using only picograms of target material.

19 Claims, 9 Drawing Sheets

Specification includes a Sequence Listing.



- (51) **Int. Cl.**
B03C 5/02 (2006.01)
G01N 27/447 (2006.01)
B01L 3/00 (2006.01)
- (52) **U.S. Cl.**
 CPC *G01N 27/447* (2013.01); *B01L 3/5027*
 (2013.01); *B03C 2201/26* (2013.01)

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,897,361	B2	3/2011	Westbrook et al.
2008/0107601	A1	5/2008	Lauwereys et al.
2008/0305498	A1*	12/2008	Milewicz G01N 33/6803 435/7.21
2009/0294291	A1	12/2009	Voldman et al.
2011/0044986	A1	2/2011	Biere-Citron et al.
2011/0108422	A1	5/2011	Heller et al.

FOREIGN PATENT DOCUMENTS

WO	2006040153	A2	4/2006
WO	2011031720	A1	3/2011
WO	2012/058334	A1	5/2012

OTHER PUBLICATIONS

Horiuchi et al., (2007). "Electroosmosis with step changes in zeta potential in microchannels." *AICHE Journal* 53: 2521-2533.

Patent Cooperation Treaty, International Searching Authority, Search Report and Written Opinion for PCT/US2013/30563, 10 pages, May 22, 2013.

Nannenga, Brent L., Andleeb Zameer, and Michael R. Sierks. "Anti-oligomeric single chain variable domain antibody differentially affects huntingtin and α -synuclein aggregates." *FEBS Letters* 582. 4 (2008): 517-522.

Kvam E, Sierks MR, Shoemaker CB, Messer A. "Physico-chemical determinants of soluble intrabody expression in mammalian cell cytoplasm." *Protein Eng Des Sel.* 2010;23(6):489-498. with supplement; doi:10.1093/protein/gzq022.

Jones et al., Blood cell capture in a sawtooth dielectrophoretic microchannel, *Anal Bioanal Chem* 401: pp. 2103-2111, available online Aug. 4, 2011.*

Chiti et al., (2006). "Protein misfolding, functional amyloid, and human disease." *Annu. Rev. Biochem.* 75: 333-366.

Selkoe, (2008). "Soluble oligomers of the amyloid beta-protein impair synaptic plasticity and behavior." *Behav. Brain Res.* 192: 106-113.

Cobb et al., (2009). "Prion diseases and their biochemical mechanisms." *Biochemistry* 48: 2574-2585.

Selkoe, (1989). "The deposition of amyloid proteins in the aging mammalian brain: implications for Alzheimer's disease." *Ann Me* 21: 73-76.

Dahlgren et al., (2002). "Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability." *J. Biol. Chem.* 277: 32046-32053.

Deshpande et al., (2006). "Different conformations of amyloid beta induce neurotoxicity by distinct mechanisms in human cortical neurons." *J. Neurosci.* 26: 6011-6018.

Lambert et al., (1998). "Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins." *Proc. Natl. Acad. Sci. USA* 95: 6448-6453.

Lesne et al., (2006). "A specific amyloid-beta protein assembly in the brain impairs memory." *Nature* 440: 352-357.

Selkoe, (2000). "Toward a comprehensive theory for Alzheimer's disease. Hypothesis: Alzheimer's disease is caused by the cerebral accumulation and cytotoxicity of amyloid beta-protein." *Ann. NY Acad. Sci.* 924: 17-25.

Lue et al., (1999). "Soluble amyloid beta peptide concentration as a predictor of synaptic change in Alzheimer's disease." *Am. J. Pathol.* 155: 853-862.

McLean et al., (1999). "Soluble pool of Abeta amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease." *Ann. Neurol.* 46: 860-866.

Walsh et al., (1999). "Amyloid beta-protein fibrillogenesis. Structure and biological activity of protofibrillar intermediates." *J. Biol. Chem.* 274: 25945-25952.

Harper et al., (1997). "Observation of metastable A β amyloid protofibrils by atomic force microscopy." *Chemistry & Biology* 4: 119-125.

Wang et al., (2002). "Soluble oligomers of beta amyloid (1-42) inhibit long-term potentiation but not long-term depression in rat dentate gyrus." *Brain Res.* 924: 133-140.

Walsh et al., (2002). "Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo." *Nature* 416: 535-539.

Chang et al., (2003). "Femtomole immunodetection of synthetic and endogenous amyloid-beta oligomers and its application to Alzheimer's disease drug candidate screening." *J. Mol. Neurosci.* 20: 305-313.

Gong et al., (2003). "Alzheimer's disease-affected brain: presence of oligomeric A beta ligands (ADDLs) suggests a molecular basis for reversible memory loss." *Proc. Natl. Acad. Sci. USA* 100: 10417-10422.

Tsai et al., (2004). "Fibrillar amyloid deposition leads to local synaptic abnormalities and breakage of neuronal branches." *Nat. Neurosci.* 7: 1181-1183.

Koffie et al., (2009). "Oligomeric amyloid beta associates with postsynaptic densities and correlates with excitatory synapse loss near senile plaques." *Proc. Natl. Acad. Sci. USA* 106: 4012-4017.

Cleary et al., (2005). "Natural oligomers of the amyloid-beta protein specifically disrupt cognitive function." *Nat. Neurosci.* 8: 79-84.

Townsend et al., (2006). "Effects of Secreted Oligomers of Amyloid {beta}-Protein on Hippocampal Synaptic Plasticity: A Potent Role for Trimers." *J. Physiol.* 572(Pt 2): 477-492.

Walsh et al., (2005). "The role of cell-derived oligomers of Abeta in Alzheimer's disease and avenues for therapeutic intervention." *Biochem. Soc. Trans.* 33: 1087-1090.

Lambert et al., (2001). "Vaccination with soluble Abeta oligomers generates toxicity-neutralizing antibodies." *J. Neurochem.* 79: 595-605.

Kayed et al., (2003). "Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis." *Science* 300: 486-489.

Lacor et al., (2004). "Synaptic targeting by Alzheimer's-related amyloid beta oligomers." *J. Neurosci.* 24: 10191-10200.

Klein et al., (2004). "Small assemblies of unmodified amyloid beta-protein are the proximate neurotoxin in Alzheimer's disease." *Neurobiol. Aging* 25: 569-580.

Cummings et al., (2003). "Dielectrophoresis in Microchips Containing Arrays of Insulating Posts: Theoretical and Experimental Results." *Anal. Chem.* 75: 4724-4731.

Cummings, (2003). "Streaming Dielectrophoresis for Continuous-Flow Microfluidic Devices." *IEEE Engineering in Medicine and Biology Magazine*, Nov./Dec.: 75-84.

Lapizco-Encinas et al., (2004). "Insulator-based dielectrophoresis for the selective concentration and separation of live bacteria in water." *Electrophoresis* 25: 1695-1704.

Lapizco-Encinas et al., (2004). "Dielectrophoretic Concentration and Separation of Live and Dead Bacteria in an Array of Insulators." *Anal. Chem.* 76: 1571-1579.

Barrett et al., (2005). "Dielectrophoretic manipulation of particles and cells using insulating ridges in faceted prism microchannels." *Analytical Chemistry* 77: 6798-6804.

Lapizco-Encinas et al., (2005). "An insulator-based (electrodeless) dielectrophoretic concentrator for microbes in water." *Journal of Microbiological Methods* 62: 317-326.

Simmons et al., (2006). "The development of polymeric devices as dielectrophoretic separators and concentrators." *MRS Bulletin* 31: 120-124.

Davalos et al., (2008). "Performance impact of dynamic surface coatings on polymeric insulator-based dielectrophoretic particle separators." *Analytical and Bioanalytical Chemistry* 390: 847-855.

(56)

References Cited

OTHER PUBLICATIONS

- Pysher et al., (2007). "Electrophoretic and dielectrophoretic field gradient technique for separating bioparticles." *Analytical Chemistry* 79: 4552-4557.
- Chen et al., (2009). "Insulator-based Dielectrophoretic Separations of Small Particles in Sawtooth Channel." *Electrophoresis* 30: 1441-1448.
- Staton et al., (2010). "Characterization of particle capture in a sawtooth patterned insulating electrokinetic microfluidic device." *Electrophoresis* 31: 3634-3641.
- Weiss et al., (2011). "Dielectrophoretic mobility determination in DC insulator-based dielectrophoresis." *Electrophoresis* 32: 2292-2297.
- Washizu et al., (1996). "Generalized multipolar dielectrophoretic force and electrorotational torque calculation." *Journal of Electrostatics* 38: 199-211.
- Barkhordarian et al., (2006). "Isolating recombinant antibodies against specific protein morphologies using atomic force microscopy and phage display technologies." *Protein Eng. Des. Sel.* 19: 497-502.
- Shlyakhtenko et al., (2007). "Single-molecule selection and recovery of structure-specific antibodies using atomic force microscopy." *Nanomedicine* 3: 192-197.
- Wang et al., (2009). "Characterizing Antibody Specificity to Different Protein Morphologies by AFM." *Langmuir* 25: 912-918.
- Emadi et al., (2004). "Inhibiting Aggregation of alpha-Synuclein with Human Single Chain Antibody Fragments." *Biochemistry* 43: 2871-2878.
- Zhou et al., (2004). "A human single-chain Fv intrabody blocks aberrant cellular effects of overexpressed alpha-synuclein." *Mol. Ther.* 10: 1023-1031.
- Liu et al., (2004). "Single chain variable fragments against beta-amyloid (Abeta) can inhibit Abeta aggregation and prevent abeta-induced neurotoxicity." *Biochemistry* 43: 6959-6967.
- Zameer et al., (2006). "Single Chain Fv Antibodies against the 25-35 Abeta Fragment Inhibit Aggregation and Toxicity of Abeta42." *Biochemistry* 45: 11532-11539.
- Marcus et al., (2008). "Characterization of an antibody scFv that recognizes fibrillar insulin and beta-amyloid using atomic force microscopy." *Nanomedicine* 4: 1-7.
- Emadi et al., (2007). "Isolation of a human single chain antibody fragment against oligomeric alpha-synuclein that inhibits aggregation and prevents alpha-synuclein-induced toxicity." *J. Mol. Biol.* 368: 1132-1144.
- Emadi et al., (2009). "Detecting morphologically distinct oligomeric forms of alpha-synuclein." *J. Biol. Chem.* 284: 11048-11058.
- Zameer et al., (2008). "Anti-oligomeric Abeta single-chain variable domain antibody blocks Abeta-induced toxicity against human neuroblastoma cells." *J. Mol. Biol.* 384: 917-928.
- Kasturirangan et al., (2010). "Nanobody specific for oligomeric beta-amyloid stabilizes non-toxic form." *Neurobiology of Aging* 33(7): 1320-1328.
- Roychaudhuri et al., (2009). "Amyloid beta-Protein Assembly and Alzheimer Disease." *Journal of Biological Chemistry* 284: 4749-4753.
- Picou et al., (2010). "Analysis of monomeric A beta (1-40) peptide by capillary electrophoresis." *Analyst* 135: 1631-1635.
- Picou et al., (2011). "Analysis of A-beta (1-40) and A-beta (1-42) Monomer and Fibrils by Capillary Electrophoresis." *Journal of Chromatography B* 879: 627-632.
- Lapizco-Encinas et al., (2008). "Protein manipulation with insulator-based dielectrophoresis and direct current electric fields." *Journal of Chromatography A* 1206: 45-51.
- Clarke et al., (2005). "Trapping of proteins under physiological conditions in a nanopipette." *Angewandte Chemie-International Edition* 44: 3747-3750.
- Holzel et al., (2005). "Trapping single molecules by dielectrophoresis." *Physical Review Letters* 95: 128102.
- Wang et al., (2008). "Covalent modified hydrophilic polymer brushes onto poly(dimethylsiloxane) microchannel surface for electrophoresis separation of amino acids." *Journal of Chromatography A* 1192: 173-179.
- Lucy et al., (2008). "Non-covalent capillary coatings for protein separations in capillary electrophoresis." *Journal of Chromatography A* 1184: 81-105.
- Mansfield et al., (2007). "Preparation and characterization of cross-linked phospholipid bilayer capillary coatings for protein separations." *Analytical Chemistry* 79: 3135-3141.
- Mohamadi et al., (2007). "Dynamic coating using methylcellulose and polysorbate 20 for nondenaturing electrophoresis of proteins on plastic microchips." *Electrophoresis* 28: 830-836.
- Liu et al., (2007). "Poly(dimethylsiloxane) microchips with two sharpened stretching tips and its application to protein separation using dynamic coating." *Chinese Journal of Chemistry* 25: 190-195.
- MacDonald et al., (2006). "Highly efficient protein separations in capillary electrophoresis using a supported bilayer/diblock copolymer coating." *Journal of Chromatography A* 1130: 265-271.
- Zhang et al., (2006). "Poly(N,N-dimethylacrylamide)-grafted polyacrylamide: A self-coating copolymer for sieving separation of native proteins by CE." *Electrophoresis* 27: 3086-3092.
- Huang et al., (2005). "Coating of poly(dimethylsiloxane) with n-dodecyl-beta-D-maltoside to minimize nonspecific protein adsorption." *Lab on a Chip* 5: 1005-1007.
- Gu et al., (2005). "Preparation and evaluation of poly (polyethylene glycol methyl ether acrylate-co-polyethylene glycol diacrylate) monolith for protein analysis." *Journal of Chromatography A* 1079: 382-391.
- Hellmich et al., (2005). "Poly(oxyethylene) based surface coatings for poly(dimethylsiloxane) microchannels." *Langmuir* 21: 7551-7557.
- Ros et al., (2006). "Bioanalysis in structured microfluidic systems." *Electrophoresis* 27: 2651-2658.
- Lee et al., (2005). "An aqueous-based surface modification of poly(dimethylsiloxane) with poly(ethylene glycol) to prevent biofouling." *Langmuir* 21: 11957-11962.
- Wong et al., (2009). "Surface molecular property modifications for poly(dimethylsiloxane) (PDMS) based microfluidic devices." *Microfluidics and Nanofluidics* 7: 291-306.
- Zhou et al., (2009). "Convenient Method for Modifying Poly(dimethylsiloxane) with Poly(ethylene glycol) in Microfluidics." *Analytical Chemistry* 81: 6627-6632.
- Chen et al., (2008). "Biocompatible polymer materials: Role of protein-surface interactions." *Progress in Polymer Science* 33: 1059-1087.
- Sugiura et al., (2008). "Surface modification of polydimethylsiloxane with photo-grafted poly(ethylene glycol) for micropatterned protein adsorption and cell adhesion." *Colloids and Surfaces B-Biointerfaces* 63: 301-305.
- Jung et al., (2008). "A cell-repellent sulfonated PEG comb-like polymer for highly resolved cell micropatterns." *Journal of Biomaterials Science-Polymer Edition* 19: 161-173.
- Hayes, (1999). "Extension of external voltage control of electroosmosis to high-pH buffers." *Analytical Chemistry* 71: 3793-3798.
- Towns et al., (1990). "Polyethyleneimine-bonded phases in the separation of proteins by capillary electrophoresis." *Journal of Chromatography* 516: 69-78.
- Rosario et al., (2002). "Photon-Modulated Wettability Changes on Spiropyran-Coated Surfaces." *Langmuir* 18: 8062-8069.
- Rosario et al., (2003). "Solvatochromic study of the microenvironment of surface-bound spiropyrans." *Langmuir* 19: 8801-8806.
- Bunker et al., (2003). "Direct Observation of Photo Switching in Tethered Spiropyran Using the Interfacial Force Microscope." *Nano Lett* 3: 1723-1727.
- Rosario et al., (2004). "Lotus Effect Amplifies Light-Induced Contact Angle Switching." *Journal of Physical Chemistry A* 108: 12640-12642.
- Hayes et al., (1999). "Preservation of NADH voltammetry for enzyme-modified electrodes based on dehydrogenase." *Analytical Chemistry* 71: 1720-1727.
- Liu et al., (2004). "Residues 17-20 and 30-35 of beta-amyloid play critical roles in aggregation." *J. Neurosci. Res.* 75: 162-171.

(56)

References Cited

OTHER PUBLICATIONS

Liu et al., (2004). "Proteolytic antibody light chains alter beta-amyloid aggregation and prevent cytotoxicity." *Biochemistry* 43: 9999-10007.

Marcus et al., (2006). "Isolation of an scFv targeting BRG1 using phage display with characterization by AFM." *Biochem. Biophys. Res. Commun.* 342: 1123-1129.

Unger et al., (2000). "Monolithic microfabricated valves and pumps by multilayer soft lithography." *Science* 288: 113-116.

Shankar et al., (2008). "Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory." *Nat. Med.* 14: 837-842.

Walsh et al., (2000). "The oligomerization of amyloid beta-protein begins intracellularly in cells derived from human brain." *Biochemistry* 39: 10831-10839.

Kasturirangan et al., (2010). "Engineered Proteolytic Nanobodies Reduce A beta Burden and Ameliorate A beta-Induced Cytotoxicity." *Biochem.* 49(21): 4501-4508.

Staton et al., (2012). "Manipulation and capture of AB amyloid fibrils and monomers by DC insulator gradient dielectrophoresis (DC-iGDEP)." *Analyst* 137: 3227-3229.

Jones et al., (2011). "Blood cell capture in a sawtooth dielectrophoretic microchannel." *Anal. Bioanal. Chem.* 401(7): 2103-2111.

Srivastava et al., (2011). "DC insulator dielectrophoretic applications in microdevice technology: a review." *Anal. Bioanal. Chem.* 399(1): 301-321.

STATON, (2013). "New methods for biological and environmental protein fingerprinting: from traditional techniques to new technology." PhD Dissertation, Arizona State University. repository.asu.edu/attachments/56754/content/Staton_asu_0010E_10534.pdf.

Kasturirangan et al., (2013). "Isolation and Characterization of Antibody Fragments Selective for Specific Protein Morphologies from Nanogram Antigen Samples." *Biotechnology Progress* 29(2): 463-471.

* cited by examiner




FIG. 1

CDR regions

AMINO ACID

- C6T SYAMS (SEQ ID NO: 16)
- C6T AISGSGGSTYYADSVKG (SEQ ID NO: 17)
- C6T SYGSVKISCFDY (SEQ ID NO: 18)
- C6T KSSQSVLYNSNNKNYLA (SEQ ID NO: 19)
- C6T WASTRES (SEQ ID NO: 20)
- C6T QQFYSTPPT (SEQ ID NO: 21)

DNA

- Agctatgccatgagc (SEQ ID NO: 22)
- Gctattagtggttagtggtggttagcacatactacgcagactccgtgaagggc (SEQ ID NO: 23)
- Agctatggttcagttaaaataagctgctttgactac (SEQ ID NO: 24)
- Aagtccagccagagtgttctttacaactccaacaataagaactacttagct (SEQ ID NO: 25)
- Tgggcatcaaccggaatcc (SEQ ID NO: 26)
- cagcaatattatagctactcctccgact (SEQ ID NO: 27)

>10H (SEQ ID NO:12)

ccatggccgaggtgcagctggtggagtctgggggaggttggtacagcctggggggtccctgagactctcctgtg
cagcctctggattcacctttagcagctatgccatgagctgggtccgccaggctccagggaaagggctggagtggg
tctcaaatattagtagtgcagggaaagggctggagtgggtctcaagtattgatgattctgggtctctacatatt
acgcagactccgtgaagggccggttcaccatctccagagacaattccaagaacacgctgtatctgcaaatgaaca
gcctgagagccgaggacacggcctatattactgtgcgaaagattctgcttctttgactactggggccagggaa
ccctggtcaccgtctcagcgggtggaggcgggtcagggcggaggtggcagcggcgggtggcgggtcgacggacatcc
agatgaccagctctccatcctcctgtctgcacatctgtaggagacagagtcaccatcacttgcgggcaagtcaqa
gcattagcagctatttaaattggtatcagcagaaaccagggaaagcccctaagctcctgatctatactgcaccca
gtttgcaaaagtgggggtcccatcaaggttcagtgccagtggtctgggacagatttcactctcaccatcagcagtc
tgcaacctgaagatcttgcaacttactactgtcaacagctctgctgcttagctccttctacgttcggccaagggacca
aggtggaaatcaaaccggggcggccacatcaccatcaccatcacggggccgcagaacaaaaactctcagaag
nggatcnaangggncgg

>D5Q (SEQ ID NO:11)

ccatggccgaggtgcagctggtggagtctgggggaggttggtacagcctggggggtccctgagactctcctgtg
cagcctctggattcacctttagcagctatgccatgagctgggtccgccaggctccagggaaagggctggagtggg
tctcatcgattggtcagaaggggtggtacacagtagcagactccgtgaagggccggttcaccatctccagag
acaattccaagaacacgctgtatctgcaaatgaacagcctgagagccgaggacacggcctatattactgtgcga
aacatcttgagaatcttgactactggggccagggaaaccctgggtcacctctcagcgggtggaggcgggtcagggc
gaggtggcagcggcgggtggcgggtcgacggacatccagatgaccagctctccatcctcctgtctgeatctgtag
gagacagagtcaccatcacttgcgggcaagtcagagcattagcagctatttaaattggtatcagcagaaaccag
ggaaagcccctaagctcctgatctatgctgcacccatttgcaaaagtgggggtcccatcaaggttcagtgccagtg
gatctgggacagatttcactctcaccatcagcagctctgcaacctgaagatcttgcaacttactactgtcaacaga
cgcgtaggccgcttctacgttcggccaagggaccaaggtggaaatcaaaccggggcggccacatcaccatcacc
atcacggggccgcagaacaaaaactcactctcagaagagaatcactagtgccggccgctgcaggtcgaccata

FIG. 2

>6E (SEQ ID NO:13)

ttgttattactcgcggccagccggccatggccgaggtgcagctgttggagtctggggaggcttggtagcct
ggggggtccctgagactctcctgtgcagcctctggattcacctttagcagctatgccatgagctgggtccgccag
gctccagggaaagggctggagtgggtctcatatattgctagtgggtgataactaaaattacgcagactccgtg
aagggccggttcaccatctccagagacaattccaagaacacgctgtatctgcaaatgaacagcctgagagccgag
gacacggccgtatattactgtgcgaaaggtgcttctgcttttgactactggggccagggaaacctggtcaccgtc
tcgagcgggtggagggcgggttcaggcggaggtggcagcggcggtggcgggtcgacggacatccagatgaccagctc
ccatcctccctgtctgcactctgtaggagacagagtcaccatcacttgccgggcaagtcagagcattagcagctat
ttaaattggtatcagcagaaaccagggaaagcccctaagctcctgatctatgctgcactcctatttgcaagtggg
gtcccatcaaggttcagtgccagtgatctgggacagatttcactctcaccatcagcagctctgcaacctgaagat
tttgcaacttactactgtcaacagagttctaatgatccttatacgttcggccaagggaccaaggtggaaatcaa
cgggcggccgcacatcatcatcaccatcacggggccgcnanaacaaaaactcctctcaannnnntctgaatggg
ggccncatanactggtgaaagtgttttnaaacctctacanaaaantcnnitt

>A4 (SEQ ID NO:8)

ttgttattactcgcggccagccggccatggccgaggtgcagctgttggagtctggggaggcttggtagcct
ggggggtccctgagactctcctgtgcagcctctggattcacctttagcagctatgccatgagctgggtccgccag
gctccagggaaagggctggagtgggtctcagcagttcagcactggtgcggcagacaacttacgcagactccgtg
aagggccggttcaccatctccagagacaattccaagaacacgctgtatctgcaaatgaacagcctgagagccgag
gacacggccgtatattactgtgcgaaagcgtttccggcgtttgactactggggccagggaaacctggtcaccgtc
tcgagcgggtggagggcgggttcaggcggaggtggcagcggcggtggcgggtcgacggacatccagatgaccagctc
ccatcctccctgtctgcactctgtaggagacagagtcaccatcacttgccgggcaagtcagagcattagcagctat
ttaaattggtatcagcagaaaccagggaaagcccctaagctcctgatctatgctgcactccttttgcaagtggg
gtcccatcaaggttcagtgccagtgatctgggacagatttcactctcaccatcagcagctctgcaacctgaagat
tttgcaacttactactgtcaacagcgggagactgggcccnnngttcggncaannggancaangtggaatcaaac
gggcggccgcacatcatcatcaccatcacggggccgcanacaaaaactcctctcanaanaggatctgaatgggg
ccgcatanactggtgaaantgtttancaaacnncatacnnnaaattcattt

>E1 (SEQ ID NO:9)

ttgttattactcgcggccagccggcctggccgaggtgcagctgttggagtctggggaggcttggtagcctg
ggggggtccctgagactctcctgtgcagcctctggattcacctttagcagctatgccatgagctgggtccgccag
ctccagggaaagggctggagtgggtctcatctattcagcctgagggtagggcagcagctacgtagactccgtga
agggccggttcaccatctccagagacaattccaagaacacgctgtatctacaaatgaacagcctgagagccgagg
acacggccgtatattactgtgcgaaaccggcggagaggtttgactactggggccagggaaacctggtcaccgtc
cgagcgggtggagggcgggttcaggcggaggtggcagcggcggtggcgggtcgacggacatccagatgaccagctc
catcctccctgtctgcactctgtaggagacagagtcaccatcacttgccgggcaagtcagagcattagcagctatt
taaattggtatcagcagaaaccagggaaagcccctaagctcctgatctatgctgcactccttttgcaagtgggg
tcccatcaaggttcagtgccagtgatctgggacagatttcactctcaccatcagcagctctgcaacctgaagatt
ttgcaacttactactgtcaacagagttacagtacccttaatacgttcggccaagggaccaaggtggaaatcaaac
gggcggccgcacatcatcatcaccatcacggggccgcaacaaaaactcctctcanaanaggatctgaatgggg
ccgcatagactggtgaaagtgtttancaaacctcatacagaaaattcattt

FIG. 2
CONTINUED

>D10L2 (SEQ ID NO:14)

atggccgaggtgcagctggtggagtctgggggagggctggtccagcctgggaggtccctgagactctcctgtgca
gectctggattcaccttcagtagctatggcatgcactgggtccgccagggcccaggcaaggggctggagtgggtg
gcagttatatcatatgatggaagtaataaataactatgcagactccgtgaagggccgattcaccatctccagagac
aattccaagaacacgctgtatctgcaagtgaacagcctgagagctgaggacacggccgtgtattactgtgcaaga
attaatgcgaagtggggccaaggtaccctggtcaccgtctcagagtggaggggcttcaggcggaggtggctct
ggcggtagtgcacttgacatccagatgacccagctctccatcctccctgtctgcatctgtaggagacagagtcacc
atcacttgccgggcaagtcagagcattagcagctatTTAAATTGGTATCAGCAGAAACCAGGGAAAGCCCTAAG
ctcctgatctatgctgcatccagtttgcaagtggggtccatcaaggttcagtgccagtggtctgggacagat
ttcactctcaccatcagcagctctgcaacctggagatTTTgcaacttactactgtcaacagagttacagtaccccg
acgttcgggcaagggaccaaggtggaaatcaaactgctggggccgcacatcatcaccatcacggggccgcagaa
caaaaactcatctcagaagaggatct

>C6 (SEQ ID NO:10)

ccatggcccaggtacagctgcaggagtcgggggagggcttggtagcagcctggggggtccctgagactctcctgtgc
agcctctggattcacctttagcagctatgccatgagctgggtccgccaggtccagggaaagggctggagtgggt
ctcagctattagtggtagtggtggtagcacatactacgcagactccgtgaagggccgattcaccatctccagaga
caattccaagaacacgctgtatctgcaaatgaacagcctgagagctgaggacacggctgtgtattactgtgcgaa
gagctatgggttcagttaaaataaactgctttgactactggggccagagcaccctggtcaccgtctcctcaggtgg
aggcgggttcaggcggaggtggctctggcgggtggcggatcggaaattgtgctgacgcagctctccagactccctggc
tgtgtctctgggcgagagggccaccatcaactgcaagtcagccagagtggtctttacaactccaacaataagaa
ctacttagcttggtagcagcagaaaccaggacagctctcctgagttgctcatttactgggcatcaaccggggaatc
cgggggtccctgaccgattcagtgccagcgggtctgggacagaattcactettaccatcagcagcctgcagcctga
ggatgtggcagtttattactgtcagcaattttatagtactctccgacttttggccaggggaccaagctggagat
caaactgctggggccgcacatcatcaccatcacggggccgcagaaacaaaactcatctcagaagaggatc

Protein

>10H (SEQ ID NO:5)

MAEVQLLES~~GGGLVQPGGSLRLS~~CAASGFTFSSYAMSWVRQAPGKLEWVSNISSAGKLEWVSSIDDSGASTYY
ADS~~VKGRFTISR~~DNSKNTLYLQMN~~SLRAEDTAVYYCAKDSAS~~FDYWGQGLVTVSSGGGSGGGSGGGSTDIQ
MTQSPSSLSASV~~GRVTITCRASQSISSYL~~NWYQQKPGKAPKLLIY~~TASSLQSGVPSRFSGSGSGTDF~~TLTISSL
QPEDFATYYCOQSAAS~~PSTFGQGTKVEIKRAAAHHHHHGAARQKLI~~SEEDLNCAA*

>6E (SEQ ID NO:6)

MAEVQLLES~~GGGLVQPGGSLRLS~~CAASGFTFSSYAMSWVRQAPGKLEWVSYIASGGDTTNYADSVKGRFTISR~~D~~
NSKNTLYLQMN~~SLRAEDTAVYYCAK~~ASAFDYWGQGLVTVSSGGGSGGGSGGGSTDIQMTQSPSSLSASV~~G~~
RVTITCRASQSISSYL~~NWYQQKPGKAPKLLIYA~~ASYLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCOQ~~S~~
SNDP~~YTFGQGTKVEIKRAAAHHHHHGAARQKLI~~SEEDLNCAA*

FIG. 2
CONTINUED

>D5Q (SEQ ID NO:4)

MAEVQLLESQGGGLVQPGGSLRLSQAASGFTFSSYAMSWVRQAPGKGLEWVSSIGQKGGGTQYADSVKGRFTISRDN
NSKNTLYLQMNLSRAEDTAVYYCAKHFENFDYWGQGTLVTVSSGGGGSGGGSGGGGSTDIQMTQSPSSLSASVGD
RVTITCRASQSISSYLNWYQKPKGKAPKLLIYAASHLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCOQT
RRPPSTFGQGTKVEIKRAAAHHHHHHGAAEQKLISEEDLNCAA*

>A4 (SEQ ID NO:2)

MAEVQLLESQGGGLVQPGGSLRLSQAASGFTFSSYPMSWVRQAPGKGLEWVSAIQHTGAFTTYADSVKGRFTISRDN
NSKNTLYLQMNLSRAEDTAVYYCAKAFPPFDYWGQGTLVTVSSGGGGSGGGSGGGGSTDIQMTQSPSSLSASVGD
RVTITCRASQSISSYLNWYQKPKGKAPKLLIYASLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCOQR
ETGPKAFGQGTKVEIKRAAAHHHHHHGAAEQKLISEEDLNCAA*

>E1 (SEQ ID NO:3)

MAEVQLLESQGGGLVQPGGSLRLSQAASGFTFSSYAMSWVRQAPGKGLEWVSSIQPEGRRTAYVDSVKGRFTISRDN
NSKNTLYLQMNLSRAEDTAVYYCAKPPERFDYWGQGTLVTVSSGGGGSGGGSGGGGSTDIQMTQSPSSLSASVGD
RVTITCRASQSISSYLNWYQKPKGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCOQS
YSTPNTFGQGTKVEIKRAAAHHHHHHGAAEQKLISEEDLNCAA*

>D10L2 (SEQ ID NO:7)

MAEVQLVESGGGVVQPGRSLRLSQAASGFTFSSYGMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFTISRDN
NSKNTLYLQMNLSRAEDTAVYYCARINAKWGQGTLVTVSSGGGGSGGGSGGGSALDIQMTQSPSSLSASVGD
RVTITCRASQSISSYLNWYQKPKGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCOQSYSTP
TFGQGTKVEIKRAAAHHHHHHGAAEQKLISEEDLNCAA*

>C6T (SEQ ID NO:15)

MAQVQLQESGGGLVQPGGSLRLSQAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTISRDN
NSKNTLYLQMNLSRAEDTAVYYCAKSYGSKISCFDYWGQGTLVTVSSGGGGSGGGSGGGGSEIVLTQSPD
SLAVSLGERATINCKSSQSVLYNSNNKNYLAWYQKPKGQSPPELLIYWASTRESGVPDRFSGSGSGTEFTLTISLQAE
DVAVYYCQOFYSTPPTFGQGTKLEIKRAAAHHHHHHGAAEQKLISEEDLNCAA*

FIG. 2
CONTINUED




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8

**NANOSCALE PROCESS TO GENERATE
REAGENTS SELECTIVE FOR INDIVIDUAL
PROTEIN VARIANTS**

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue; a claim printed with strikethrough indicates that the claim was canceled, disclaimed, or held invalid by a prior post-patent action or proceeding.

**[FEDERAL GRANT] GOVERNMENT SUPPORT
CLAUSE**

This invention was made with [Government] government support under [Grants No. 2ROIEB004761-06 and R21IEB010191-01A1,] *R01 EB004761 and R21 EB010191* awarded by the National Institutes of Health. The government has certain rights in the invention.

RELATED APPLICATION

More than one reissue application has been filed for the reissue of U.S. Pat. No. 9,938,330. The reissue applications are U.S. patent application Ser. No. 16/845,761, filed on Apr. 10, 2020, and U.S. patent application Ser. No. 18/134,886, filed Apr. 14, 2023, which is a reissue division of U.S. patent application Ser. No. 16/845,761, all of which are reissues of U.S. Pat. No. 9,938,330.

This application is an application for reissue of U.S. Pat. No. 9,938,330, which issued from U.S. application Ser. No. 14,388,209, filed Mar. 12, 2013, which is a 371 of international application PCT/US2013/030563, filed Mar. 12, 2013, and claims priority to U.S. Provisional Patent Application No. 61/617,476, filed Mar. 29, 2012, [the entirety of which is] which are incorporated herein by reference in their entirety.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on [Feb. 21, 2013] *Apr. 23, 2020*, is named [17555.007WO1_SL.txt] *G811802307_SL25.txt* and is [34,904] *45,584* bytes in size.

BACKGROUND OF THE INVENTION

Protein variants including alternatively processed and misfolded proteins have been associated with many different human diseases. For example, misfolded protein aggregates play a critical role in many devastating human diseases including Alzheimer's (AD) and Parkinson's diseases, diabetes and cancer. However, assessing the variant's role in the onset and progression of different diseases is hampered by a lack of reagents that can distinguish between protein isoforms in vivo. Development of such reagents has been hindered by two main factors: protein variants often occur at only trace levels in vivo and the variants may differ only subtly from the parent form, making them difficult to separate and purify. Novel separation technologies that can tease apart subtle protein variants along with novel molecular recognition protocols are needed to create reagents with sufficient specificity to distinguish between these protein variants. Therefore, there is a continuing need for technolo-

gies capable of generating highly selective reagents to specific protein isoforms isolated from mammalian tissue.

SUMMARY OF THE INVENTION

5

The present invention provides a DC-iGDEP device for separating target protein species in a biological sample based on a variety of chemical and physical parameters comprising an open sawtooth microfluidic channel having an inlet port and an outlet port, and gates between each tooth of the channel, wherein spacing of the gates starts at 50 microns and decreases over two centimeters to 1 micron, and wherein the teeth insulate adjoining gates. As used herein the gap distance of a gate is the distance between teeth across from each other on opposite sides of the device.

10

The present invention provides a method of separating from a biological sample a target species based on various chemical and physical parameters including charge, size, permittivity, deformation, shape and other factors comprising

15

- (a) providing the device for separating target protein species in a biological sample based on a variety of chemical and physical parameters comprising an open sawtooth microfluidic channel having an inlet port and an outlet port, and gates between each tooth of the channel, wherein spacing of the gates starts at 50 microns and decreases over two centimeters to 1 micron, and wherein the teeth insulate adjoining gates,
- (b) loading a loading volume of the sample into the inlet port,
- (c) applying a field to the device to separate particles or molecules in the sample, and
- (d) recovering the target species.

20

In certain embodiments, the target species is an A β aggregate. In certain embodiments, the recovered A β aggregates with an antibody to confirm the size of the A β aggregate. In certain embodiments, the antibody is specific for oligomeric A β aggregates. As used herein, the term "oligomer" refers to a dimer, trimer, or tetramer or larger aggregate. In certain embodiments, the antibody is a nanobody. As used herein, the term "antibody" includes scFv (also called a "nanobody"), humanized, fully human or chimeric antibodies, single-chain antibodies, diabodies, and antigen-binding fragments of antibodies (e.g., Fab fragments). In certain embodiments, the nanobody is a C6, A4, E1, D5, 10H, 6E, D10 or BSEC1 nanobody.

25

30

In certain embodiments, the biological sample has a volume of less than 100 microliters. In certain embodiments, has a volume of about 50 microliters. In certain embodiments, the biological sample is brain tissue, serum, cerebrospinal fluid (CSF), urine or saliva. In certain embodiments, the force is applied for a period of time that is less than 20 minutes, such as between 5-15 minutes. In certain embodiments, target A β aggregate is concentrated by several orders of magnitude. In certain embodiments, the target A β aggregate is concentrated by 10⁶ as compared to the loading volume. In certain embodiments, the protein is p53, islet amyloid polypeptide, beta-amyloid, tau), alpha-synuclein, huntingtin, or superoxide dismutase.

35

40

45

50

55

60

65

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. Schematic showing current DC-iGDEP design (top) used to capture A β fibrils (middle, right) but not monomers (middle, left). The proposed modified device (bottom) will 1) increase maximum DEP force ~700 times to capture monomeric A β , 2) refine the difference in local DEP

force between adjoining “gates,” and 3) enable separation and concentration of various A β aggregates (dimers, trimers, . . . up to fibrils) in separate chambers.

FIG. 2 provides nucleic acid and amino acid sequences for several nanobodies. Underlining indicates CDR regions (or nucleic acids that encode CDR regions).

FIG. 3 depicts an embodiment of the present device. Diagram of the direct current insulator gradient dielectrophoresis (DC-iGDEP) device. An insulated sawtooth pattern is integrated with a tapered microfluidic channel to generate localized non-uniform electric field gradients of increasing strength from left to right.

FIGS. 4A-4D. Fluorescence images of the narrowest portion of the DC-iGDEP for A β monomer samples (30 μ M) with (A) 400 V, (B) 600 V, (C) 800 V, and (D) 1000 V applied. The diffuse light areas show the fluorescently tagged A β monomer distributed throughout the channel. Streaming of monomer is more visually apparent in (C) and (D).

FIG. 5A-5D. Fluorescence images of the narrowest portion of the DCiGDEP (same as in FIG. 4) showing capture of A β fibrils (30 μ M) at (A) 400 V, (B) 600 V, (C) 800 V, and (D) 1000 V applied. The areas of high fluorescence localized between the tips of the sawtooth patterned insulator indicate small zones where the fibrils were captured and concentrated.

FIG. 6. TEM of the mature A β fibrils used in FIG. 5. The scale bar is 100 nm, and the fibril sample was in PBS before buffer exchange to Tris.

FIG. 7. Regions of interest (ROIs) indicated as described in the text. Three different areas were selected to determine the enrichment of fibril concentration in the capture zone using the fluorescence intensities in each region. The areas depicted here are larger than the actual ROIs used for ease of viewing.

FIG. 8. Plot showing computed magnitudes of $\nabla|E|^2$ along the microchannel centerline versus gate width. Vertical axis is plotted using a logarithmic scale. The trend line shows that the relationship between data points can be approximated using a power function.

DETAILED DESCRIPTION OF THE INVENTION

The inventors have developed methods to separate and concentrate protein variants at a microliter scale and to generate reagents to those variants with exquisite selectivity for specific protein isoforms using only picograms of target material. This capability is broadly applicable to protein variants associated with many human diseases. In certain embodiments, this method is used to generate reagents against isoforms of the protein amyloid-beta (A β) that have been implicated in Alzheimer’s disease (AD). Misfolding and assembly of A β into an array of different aggregate species has been linked to the onset and progression of AD. A variety of different length and aggregate forms of A β have been identified in human brain tissue. Unique A β forms present in AD brain tissue are separated and isolated using a novel electric field-based separation process. Antibody-based reagents are utilized that can selectively bind the different A β species and identify which reagents can best distinguish healthy and AD brain tissue.

The present work benefits from the current availability of well characterized post-mortem human AD and healthy brain tissue. The present work utilizes novel electric field-based capabilities, which can quickly isolate and concentrate different protein isoforms using only minimal amounts of

material. This enables the collection in separate nanoliter volumes several distinct A β isoforms isolated from human AD tissue. Novel protocols have been developed that allow the isolation of single chain variable domain antibody fragments (scFvs, also called “nanobodies”) against specific protein morphologies by utilizing Atomic Force Microscopy (AFM) in conjunction with surface display antibody libraries. These protocols enable the isolation of nanobodies to specific aggregate morphologies using only picograms of material without the need for any protein modification. Panning techniques are used to generate nanobodies against the isolated A β isoforms from AD tissue. Protocols have been developed to characterize nanobody binding specificity to different target morphologies without the need to purify or modify the target antigen and again requiring only microliters of protein target solution. Techniques have been developed both to isolate subtle protein variants involved in human disease, in this case A β isoforms isolated from AD brain tissue, and also to generate and characterize nanobody reagents to the key isoforms that distinguish AD from healthy brain tissue.

Understanding the molecular basis of disease progression is often a very challenging and complex problem because the specific biochemical species involved may be extremely difficult to identify. Individual biomolecules may be unstable and differ from each other in only very subtle ways, therefore identification and isolation of different forms is a technologically demanding problem. One critically important and challenging class of closely related protein variants are different folded conformations of the same protein. Misfolded and aggregated protein variants are often indistinguishable by many analytical strategies. Over 30 human health diseases have already been connected to misfolding or misprocessing of proteins including cancer (p53), diabetes (islet amyloid polypeptide), Alzheimer’s (beta-amyloid and tau), Parkinson’s (alpha-synuclein), and Huntington’s diseases (huntingtin), Amyotrophic lateral sclerosis (superoxide dismutase) and prion based diseases. In many of these diseases, specific misfolded protein variants such as small soluble oligomeric forms of the amyloid-beta (A β) protein or a misfolded form of the prion protein have been associated with cell dysfunction and disease progression. While the role of protein expression in disease can be very effectively studied using proteomic and genomic analyses or inhibitory RNA techniques, these methods are generally not capable of distinguishing between misfolded or alternatively processed protein variants. The different protein variants are often metastable and contain only subtle differences. Robust tools that can selectively identify and manipulate aberrant protein forms would be extremely useful in studying and controlling the many diseases associated with protein variants.

Currently, studies to probe the mechanisms underlying these diseases and to develop appropriate therapeutic strategies have been greatly hampered because reagents that can selectively recognize specific protein variants are scarce. In order to address this critically important need, unique electric field-based methods have been developed to efficiently isolate and concentrate subtly different protein variants, and simultaneously developing novel biopanning methods to generate very selective molecular recognition reagents that can be used to identify specific disease related species and mechanisms.

To demonstrate the current technology, reagents were isolated that selectively bind a diverse array of different aggregate species of the protein amyloid-beta (A β). While A β was first implicated in Alzheimer’s Disease (AD) over

20 years ago the role of A β in AD is still unclear and has proven to be much more elusive than originally hoped. Much of the confusion around the role of A β in AD is due to the variety of different A β species that can occur in vivo. Aggregation of A β is a critically important though poorly understood factor in the progression of AD. While the amyloid plaques of AD contain fibrillar A β aggregates, a variety of other smaller aggregate species of A β can also be formed and increasing evidence implicates various small soluble oligomeric A β species in neurotoxicity and loss of synaptic function. Cortical levels of soluble A β correlate well with cognitive impairment and loss of synaptic function. Small soluble aggregates of A β , termed A β -derived diffusible ligands (ADDLs), and spherical or annular aggregates termed protofibrils were shown to be neurotoxic. Oligomeric forms of A β , created in vitro or derived from cell cultures, were shown to inhibit long term potentiation. The concentration of oligomeric forms of A β is also elevated in transgenic mouse models of AD and in AD brain. Disruption of neural connections was shown to occur near A β plaques and fibrils, suggesting a toxic role for the fibrillar form, however the disruption also occurred in regions without fibrillar A β deposition suggesting that the toxicity may be due to small amounts of oligomeric A β , some in equilibrium with the fibrillar form, some existing on their own. A halo of oligomeric A β was shown to surround A β plaques and correlate with synapse loss. Oligomeric A β was also shown to disrupt cognitive function in transgenic animal models of AD.

A major barrier impeding studies of the connection between A β and AD is the lack of suitable reagents to identify and localize the different A β species. Antibodies to selected A β forms have been previously generated by immunization methods using for example soluble oligomeric A β (Lambert, M. P., et al., Vaccination with soluble Abeta oligomers generates toxicity-neutralizing antibodies. *J Neurochem*, 2001. 79: p. 595-605) or molecular mimics of oligomeric A β (Kayed, R., et al., Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science*, 2003. 300: p. 486-9). These reagents have been useful to demonstrate the importance of oligomeric A β in AD for example by confirming the presence of oligomeric A β in AD brains, showing they target synaptic ligands and bind strongly to neuronal dendrites, and that oligomeric A β correlates with synapse loss near amyloid plaques. However, while the current pool of reagents has been useful in pointing out the importance of soluble A β aggregates, their binding specificities are generally not well characterized, the affinities and specificities of the reagents are not sufficient to identify the presence of specific oligomeric species or of low concentration species and are not generally suitable for precise histochemical studies or to localize species to specific cellular locations in situ or in vivo. In addition, they are not suitable for intracellular studies.

Since soluble aggregates of A β , α -synuclein (α -syn) and tau have all been associated with neurodegenerative diseases, reagents that can recognize specific morphologies of specific proteins are needed to elucidate the roles of the different forms of these proteins in disease. Therefore there is a critical need for well characterized reagents that recognize specific A β forms to facilitate studies on the role of A β aggregation in AD, to identify the relevant toxic A β species, and to clarify where and when aggregation of A β begins and how it progresses in AD. This information is vitally important to understand the mechanism of AD, to facilitate diagnoses, and to develop appropriate markers for monitoring

therapeutic interventions. Similar reagents are needed to probe the role of protein variants in other significant human diseases such as Parkinson's, diabetes, and cancer.

To address this need, novel techniques have been developed to separate and purify subtle protein variants and to generate and characterize reagents that recognize specific protein forms. To demonstrate the capabilities of the present technology to generate reagents to specific protein variants, protocols are modified so that the various in vitro generated A β species are separated. These conditions are used to separate the much more complex A β pool present in AD brain.

Subtle protein variants including alternatively processed and folded forms play critical roles in numerous human diseases, many of which cannot be distinguished by genomic and proteomic techniques. Despite the growing importance of protein variants in human disease, few tools are available to study and treat these phenomena because of the difficulty in identifying individual protein isoforms. Since there are often only subtle differences between protein variants, and many may be present only at trace levels or may not be particularly stable, isolating individual protein variants and generating reagents that selectively recognize each form are extremely challenging problems. To overcome these difficulties, protocols have been developed that enable the isolation and concentration of specific protein variant forms, and the generation of reagents that selectively bind specific protein variants using only trace amounts of target.

DC-iGDEP Separation Devices and Methods

One aspect of the present work utilizes the unique abilities of dielectrophoretic (DEP) and electrokinetic (EK) forces to separate protein isoforms. DEP forces typically are used to bifurcate particle systems or uniquely capture specific targets in areas of high gradient (or low gradient, negative versus positive DEP). The approach described herein uniquely harnesses these forces to form a separation scheme not unlike isoelectric focusing (IEF), in that an overarching gradient is formed and specific targets localize according to their unique properties: pI in the case of IEF and DEP/EK force balances in the case of DC-iGDEP. A local electric field gradient can be formed with insulating structures using a static electric field applied from electrodes located in remote inlet and outlet reservoirs (Cummings, E. B. and A. K. Singh, Dielectrophoresis in Microchips Containing Arrays of Insulating Posts: Theoretical and Experimental Results. *Anal. Chem.*, 2003. 75: p. 4724-4731; Cummings, E. B., Streaming Dielectrophoresis for Continuous-Flow Microfluidic Devices. *IEEE Engineering in Medicine and Biology Magazine*, 2003. November/December: p. 75-84; Lapizco-Encinas, B. H., et al., Insulator-based dielectrophoresis for the selective concentration and separation of live bacteria in water. *Electrophoresis*, 2004. 25: p. 1695-1704; Lapizco-Encinas, B. H., et al., Dielectrophoretic Concentration and Separation of Live and Dead Bacteria in an Array of Insulators. *Anal. Chem.*, 2004. 76: p. 1571-1579; Barrett, L. M., et al., Dielectrophoretic manipulation of particles and cells using insulating ridges in faceted prism microchannels. *Analytical Chemistry*, 2005. 77: p. 6798-6804; Lapizco-Encinas, B. H., et al., An insulator-based (electrodeless) dielectrophoretic concentrator for microbes in water. *Journal of Microbiological Methods*, 2005. 62: p. 317-326; Simmons, B. A., et al., The development of polymeric devices as dielectrophoretic separators and concentrators. *MRS Bulletin*, 2006. 31: p. 120-124; Davalos, R. V., et al., Performance impact of dynamic surface coatings on polymeric insulator-based dielectrophoretic particle separators. *Analytical and Bioanalytical Chemistry*, 2008. 390: p. 847-855).

The present method adds a global gradient to the system, allowing the longitudinal separation of mixtures as opposed to a simple bifurcation of two components (Pysher, M. D. and M. A. Hayes, Electrophoretic and dielectrophoretic field gradient technique for separating bioparticles. *Analytical Chemistry*, 2007. 79: p. 4552-4557; Chen, K. P., et al., Insulator-based Dielectrophoretic Separations of Small Particles in Sawtooth Channel. *Electrophoresis*, 2009. 30: p. 1441-1448; Jones, P. V., S. J. R. Staton, and M. A. Hayes, Blood Cell Capture in a Gradient Dielectrophoretic Microchannel. *Anal. Bioanal.* 2011. 401: p. 2103-2111; Staton, S. J. R., et al., Characterization of particle capture in a sawtooth patterned insulating electrokinetic microfluidic device. *Electrophoresis*, 2010. 31: p. 3634-3641; Weiss, N. G., et al., Dielectrophoretic mobility determination in DC insulator-based dielectrophoresis. *Electrophoresis*, 2011. 32, 2292-2297; Staton, S. R., et al., Gradient Insulator Based Dielectrophoresis Isolation and Concentration of A-beta Amyloid Fibrils. *Analyst*, 2012. 137, in press (RSC ID: AN-COM-01-2012-035138)).

The insulating structures are fabricated such that an increasing local gradient is induced along the length of a microfluidic channel (FIG. 1). This sawtooth structure enables separations to be based on high multipole moments. Generally, simple electrophoresis accesses the monopole electric properties of a molecule or particle. For complex molecules and bioparticles, several intricate electrical geometries exist which can be used for separation (see for ex. Jones, T. B. and M. Washizu, Generalized multipolar dielectrophoretic force and electrorotational torque calculation. *Journal of Electrostatics*, 1996. 38: p. 199-211). Protocols have been developed to gently (cells remain viable) separate several targets along a gradient based on specific features of each target. These features can be as subtle as deformability or as simple as size. In addition to earlier work focused on model particles, several species of bacteria and cells from whole human blood have been isolated (along with a model in support of the present work (Chen, K. P., et al., Insulator-based Dielectrophoretic Separations of Small Particles in Sawtooth Channel. *Electrophoresis*, 2009. 30: p. 1441-1448)), and it has been shown that different A β species can be uniquely captured. All devices used to generate these results were created with photolithographic templates fabricated with AZ P4620 photoresist (Pysher, M. D. and M. A. Hayes, Electrophoretic and dielectrophoretic field gradient technique for separating bioparticles. *Analytical Chemistry*, 2007. 79: p. 4552-4557).

The basic approach is demonstrated with polystyrene particles varying in size from 20 nm to 1 micron in diameter being isolated along the increasing gradient (Staton, S. J. R., et al., Characterization of particle capture in a sawtooth patterned insulating electrokinetic microfluidic device. *Electrophoresis*, 2010. 31: p. 3634-3641). The small space between the tips of the insulating structures (gates) on the narrowest portion of the device do not clog, consistent with the inventors' modeling studies showing that the particles never actually penetrate the narrowest zone (Chen, K. P., et al., Insulator-based Dielectrophoretic Separations of Small Particles in Sawtooth Channel. *Electrophoresis*, 2009. 30: p. 1441-1448). Differential behaviors were observed for two populations of the same sized particles, indicating that dielectrophoretic forces act on factors beyond just size—perhaps heterogeneous permittivity (Staton, S. J. R., et al., Characterization of particle capture in a sawtooth patterned insulating electrokinetic microfluidic device. *Electrophoresis*, 2010. 31: p. 3634-3641). Live and dead bacteria were easily separated using this setup without clogging, verifying

the modeled behavior of the micro and nano particles (Pysher, M. D. and M. A. Hayes, Electrophoretic and dielectrophoretic field gradient technique for separating bioparticles. *Analytical Chemistry*, 2007. 79: p. 4552-4557).

It was also possible to separate red blood cells (RBCs) from other types of cell debris or proteins (Blood Cell Capture in a Gradient Dielectrophoretic Microchannel. *Anal. Bioanal.* 2011. 401: p. 2103-2111). A component of this system has also been modeled, which was a series of seven tooth pairs that do not converge and no particle-particle interactions are included (Chen, K. P., et al., Insulator-based Dielectrophoretic Separations of Small Particles in Sawtooth Channel. *Electrophoresis*, 2009. 30: p. 1441-1448). Using just these limiting constraints the ratio of electrokinetic forces (electrophoretic and electroosmotic, $\mu_{ep} + \mu_{eof}$ mobilities) and dielectrophoretic forces was shown to be unique at each gate. The shape of the projected electric field and resulting field gradient is strongly influenced by the shape of the insulators.

Development of DC-iGDEP Device to Isolate a Full Range of In Vitro Generated A β Species.

The unique capabilities of DC-iGDEP are used to extend the production of nanobodies to create a suite of precise probes of A β aggregates specific to AD patients. To accomplish this, a DC-iGDEP device is fabricated that maximizes resolution of the A β aggregate species ranging from dimers to full fibrils, including metastable (several minutes time-scale) intermediates. The location within the DC-iGDEP device of selected isolated and concentrated A β 40 and 42 aggregate species is confirmed, exploiting existing nanobodies with precise affinity for various protein regions and morphologies.

Separation results have been obtained using the open channel sawtooth design on a variety of systems including different size polystyrene particles (Staton, S. J. R., et al., Characterization of particle capture in a sawtooth patterned insulating electrokinetic microfluidic device. *Electrophoresis*, 2010. 31: p. 3634-3641), red blood cells (Blood Cell Capture in a Gradient Dielectrophoretic Microchannel. *Anal. Bioanal.* 2011. 401: p. 2103-2111), and bacteria (Pysher, M. D. and M. A. Hayes, Electrophoretic and dielectrophoretic field gradient technique for separating bioparticles. *Analytical Chemistry*, 2007. 79: p. 4552-4557), and have developed a theoretical model to predict separation properties (Chen, K. P., et al., Insulator-based Dielectrophoretic Separations of Small Particles in Sawtooth Channel. *Electrophoresis*, 2009. 30: p. 1441-1448).

The inventors have also shown that the design can isolate and concentrate fully formed A β fibrils while allowing monomers to pass freely (FIG. 1). In this design, the A β fibrils were captured at a "gate" (closest approach of the insulating teeth) with a 27 micron spacing, resulting in a squared field strength gradient calculated (COMSOL multiphysics) to be approximately 10^{18} V²/m³. These results were obtained within 5-15 minutes after loading the device with approximately 40 microliters of the monomer or fibril solution/colloid. The data demonstrate that this unique separation scheme is well controlled and is ideally suited for the selective isolation and concentration of aggregation intermediates.

According to the basic accepted theories underlying DEP and EK, the net velocity of a particle/molecule is proportional to particle radius (a) squared [$v_{DEP} = \mu_{DEP} \nabla E^2$, $\mu_{DEP} \propto \alpha^2$: v_{DEP} is the velocity of a particle due to DEP forces and E is the local electric field (Chen, K. P., et al., Insulator-based Dielectrophoretic Separations of Small Particles in Sawtooth Channel. *Electrophoresis*, 2009. 30: p. 1441-1448;

Weiss, N. G., et al., Dielectrophoretic mobility determination in DC insulator-based dielectrophoresis. *Electrophoresis*, 2011. 32, 2292-2297]. Using this relationship and noting that monomeric A β nominally are about 1.5 nanometers across and the characteristic length of fibrils is in the 10s to 100s of nanometers (Roychaudhuri, R., et al., Amyloid beta-Protein Assembly and Alzheimer Disease. *Journal Of Biological Chemistry*, 2009. 284: p. 4749-4753), the range of expected DEP forces needed is effectively bracketed [estimate E, ΔE^2 from COMSOL, use A β (1-40) monomer μ_{EP} (10.0 mM TRIS buffer, pH 7.8) of 1.20×10^4 cm²/V s and A β (1-42) monomer μ_{EP} (10.0 mM TRIES buffer, pH 7.8) is 1.072×10^4 cm²/V s (about four peak widths difference between A β 40 and 42) (Picou, R., et al., Analysis of monomeric A beta (1-40) peptide by capillary electrophoresis. *Analyst*, 2010. 135: p. 1631-1635; Picou, R. A., et al., Analysis of A-beta (1-40) and A-beta (1-42) Monomer and Fibrils by Capillary Electrophoresis. *Journal of Chromatography B*, 2011DOI: 10.1016/j.jchromb.2011.01.030)]. A device with the "gate" spacing starting at 50 microns and decreasing over two centimeters to 1 micron effectively captures, at varying points, A β species ranging from monomers to fibrils (FIG. 1, bottom graphic). COMSOL multiphysics was used to design these devices, and the resulting underlying structures are transferred to AutoCAD for creation of photolithographic plates.

With this design, various A β aggregates are separated, although not homogeneously (evenly spaced or uniquely isolated), ranging from monomers to fully formed fibrils. This takes place in a matter of minutes as a typical velocity (EP only, open portions of the device) is $\sim 10^{-4}$ m/s. Further, compared to the loading volume of approximately 50 microliters, specific targets can be concentrated by several orders of magnitude (up 10^6).

Device:

Microchannel geometry consisted of sets of successively larger, equilateral triangular features lining both sides of the channel (FIG. 3). The tip of each triangle corresponded with another opposing triangle on the other side of the channel, forming sequentially narrower gaps along a converging sawtooth pattern. The smallest triangles (located near the entrance to the channel) possessed 6- μ m sides and a 5.2- μ m height. The side-length of the equilateral triangles increased by 40 μ m after every sixth repeated unit. This created a channel with an initial gap pitch of 945 μ m and a final gap pitch of 27 μ m. The whole channel length was approximately 4.1 cm with an average depth of 14 ± 1 μ m.

Materials and Methods:

Microfluidic devices constructed from both glass and polydimethylsiloxane (PDMS) were used to perform the experiments. Platinum electrodes were inserted through small access ports into reservoirs at each end of the microchannel and used to apply potential across the device.

PDMS (Sylgard 184, Dow/Corning, Midland, Mich.) was poured over a photoresist template patterned on a silicon wafer and allowed to cure. Two-millimeter diameter holes were punched through the PDMS at each end of the channel to access the reservoirs. The PDMS, along with a glass slide, were then oxidized via oxygen-plasma treatment and then irreversibly bonded.

Cell samples were typically collected in phosphate buffer and fluorescently stained to aid visualization during experiments (Vybrant DiO, Invitrogen, Inc., Carlsbad, Calif.).

Buffer was pipetted into the inlet reservoir, causing the channel to fill passively via capillary action. After inspecting the device and ensuring uniform fluid distribution, sample was then pipetted into the inlet reservoir. After analyte

bioparticles had entered the device via pressure-driven flow and attained uniform distribution within the channel, buffer was added to the opposite reservoir in order to balance the hydrodynamic pressure. Once the device was prepared in this manner, platinum electrodes (0.404 mm external diameter 99.9% purity, Alfa Aesar, Ward Hill, Mass.) were inserted through the access ports into the reservoirs and connected to a Series 225 DC power supply (Bertan High Voltage Corp., Hicksville, N.Y.). Experiments were observed with an Olympus IX70 microscope. Samples were illuminated using a broad-spectrum mercury lamp (H30 102 w/2, OSRAM) and an Olympus DAPI, FITC, Texas Red triple band-pass cube (Olympus, Center Valley, Pa.).

Generation of Nanobodies.

The inventors have developed novel technology enabling the generation of reagents that recognize specific protein conformations by combining the powerful imaging capabilities of Atomic Force Microscopy (AFM) with the molecular recognition diversity of phage display libraries (Barkhordarian, H., et al., Isolating recombinant antibodies against specific protein morphologies using atomic force microscopy and phage display technologies. *Protein Eng Des Sel*, 2006. 19: p. 497-502; Shlyakhtenko, L. S., et al., Single-molecule selection and recovery of structure-specific antibodies using atomic force microscopy. *Nanomedicine*, 2007. 3: p. 192-7). The inventors have also developed innovative technologies that allow the characterization of binding specificity of these reagents using only nanograms of material again utilizing AFM (Wang, M. S., et al., Characterizing Antibody Specificity to Different Protein Morphologies by AFM. *Langmuir*, 2008). In order to isolate single chain antibody fragments (or nanobodies) to individual aggregate forms, an AFM biopanning technology was developed that allows the visualization of the target protein morphology (Barkhordarian, H., et al., Isolating recombinant antibodies against specific protein morphologies using atomic force microscopy and phage display technologies. *Protein Eng Des Sel*, 2006. 19: p. 497-502; PCT/US11/57887; PCT/US11/57925; PCT/US11/57904).

The protocol is particularly well suited to isolate nanobodies against different protein morphologies since it minimizes protein handling, as the target protein is added to the mica surface without modification, it uses small amounts of protein, low nanogram quantities are more than sufficient, and the target protein does not have to be purified. This basic technology has been utilized to isolate nanobodies that recognize different areas of monomeric A β and a-syn (Emadi, S., et al., Inhibiting Aggregation of alpha-Synuclein with Human Single Chain Antibody Fragments. *Biochemistry*, 2004. 43: p. 2871-2878; Zhou, C., et al., A human single-chain Fv intrabody blocks aberrant cellular effects of overexpressed alpha-synuclein. *Mol Ther*, 2004. 10: p. 1023-31; Liu, R., et al., Single chain variable fragments against beta-amyloid (Abeta) can inhibit Abeta aggregation and prevent abeta-induced neurotoxicity. *Biochemistry*, 2004. 43: p. 6959-67; Zameer, A., et al., Single Chain Fv Antibodies against the 25-35 Abeta Fragment Inhibit Aggregation and Toxicity of Abeta42. *Biochemistry*, 2006. 45: p. 11532-9), fibrillar A β and a-syn (Barkhordarian, H., et al., Isolating recombinant antibodies against specific protein morphologies using atomic force microscopy and phage display technologies. *Protein Eng Des Sel*, 2006. 19: p. 497-502; Marcus, W. D., et al., Characterization of an antibody scFv that recognizes fibrillar insulin and beta-amyloid using atomic force microscopy. *Nanomedicine*, 2008. 4: p. 1-7), two different oligomeric α -syn species (Emadi, S., et al., Isolation of a human single chain antibody

fragment against oligomeric alpha-synuclein that inhibits aggregation and prevents alpha-synuclein-induced toxicity. *J Mol Biol*, 2007. 368: p. 1132-44; Emadi, S., et al., Detecting morphologically distinct oligomeric forms of alpha-synuclein. *J Biol Chem*, 2009. 284: p. 11048-58), and three different oligomeric A β species (Zameer, A., et al., Anti-oligomeric Abeta single-chain variable domain antibody blocks Abeta-induced toxicity against human neuroblastoma cells. *J Mol Biol*, 2008. 384: p. 917-28; Kasturirangan, S., et al., Nanobody specific for oligomeric beta-amyloid stabilizes non-toxic form. *Neurobiol Aging*, 2010. In press; Kasturirangan, S., et al., Isolation and Characterization of a Nanobody that Selectively Binds Brain Derived Oligomeric Beta-Amyloid. (Submitted)).

The different oligomer specific nanobodies do not show cross-reactivity, so the nanobodies binding oligomeric A β do not bind oligomeric α -syn and vice versa. The nanobodies work well in standard ELISA and immunohistochemistry assays as it has been shown that each of the different aggregate species recognized by the different nanobodies naturally occur in human AD or PD tissue, and that the nanobodies can be used to distinguish between AD, PD and healthy brain tissue, and block toxicity of different aggregate species (Emadi, S., et al., Isolation of a human single chain antibody fragment against oligomeric alpha-synuclein that inhibits aggregation and prevents alpha-synuclein-induced toxicity. *J Mol Biol*, 2007. 368: p. 1132-44; Emadi, S., et al., Detecting morphologically distinct oligomeric forms of alpha-synuclein. *J Biol Chem*, 2009. 284: p. 11048-58;

Zameer, A., et al., Anti-oligomeric Abeta single-chain variable domain antibody blocks Abeta-induced toxicity against human neuroblastoma cells. *J Mol Biol*, 2008. 384: p. 917-28; Kasturirangan, S., et al., Nanobody specific for oligomeric beta-amyloid stabilizes non-toxic form. *Neurobiol Aging*, 2010. In press). The nanobodies currently developed have several significant advantages over conventional antibodies including: 1) they target specific morphologies of a single protein target for example recognizing a selected A β oligomer form, but not any α -syn oligomer forms, 2) they can be affinity matured to femtomolar levels; 3) they can be genetically modified with targeting or tag sequences; 4) they can be expressed intracellularly as intrabodies to identify intracellular A β species; and 5) their specificities can be carefully characterized.

Combination of the novel separation and molecular recognition technologies enables the ability to identify and concentrate specific protein isoforms connected with diseased human tissue and to generate nanobody reagents that selectively recognize the different protein isoforms characteristic of the disease. The resulting panel of nanobodies provide extremely powerful tools in this case for the AD community to better define the roles of the different A β species in the progression of AD. This effort provides a clear and immediately useful example of the overall strategy of highly refined reagent development.

In certain embodiments, the antibody fragments that can be used in the present invention are those listed in Table 1 below (See also FIG. 2):

TABLE 1

Antibody Fragment	Library source	Specificity	Assays to validate	Applications demonstrated	SEQ ID NO
A4	Tomlinson (MRC)	Oligomeric Abeta 3-day aggregates	Dot blot, time course, ELISA, AFM (Note: 3-day oligomeric target is not stable for westerns)	Human AD brain tissue, Human CSF, Mouse AD brain tissue, (Dot blot assays), Immunohistochemistry	amino acid SEQ ID NO: 2; nucleic acid SEQ ID NO: 8
E1	Tomlinson (MRC)	Oligomeric Abeta 1-day aggregates	Dot blot, time course, ELISA, AFM (Note: 1-day oligomeric target is not stable for westerns)	Human AD brain tissue, Human CSF, Mouse AD brain tissue, (Dot blot assays)	amino acid SEQ ID NO: 3; nucleic acid SEQ ID NO: 9
C6	Sheets (UCSF)	Oligomeric Abeta: brain derived	AFM	Human AD brain tissue, Human CSF, Mouse AD brain tissue, (Dot blot assays)	amino acid SEQ ID NO: 1; nucleic acid SEQ ID NO: 10 and amino acid SEQ ID NO: 15
D5	Tomlinson (MRC)	Oligomeric a-synuclein 3-day aggregates	Dot blot, time course, ELISA, AFM, western blot analysis	Human PD brain tissue, Human CSF, Mouse PD brain tissue, (Dot blot assays, western blot, Immunohistochemistry with tissue and cells)	amino acid SEQ ID NO: 4; nucleic acid SEQ ID NO: 11

TABLE 1-continued

Antibody Fragment	Library source	Specificity	Assays to validate	Applications demonstrated	SEQ ID NO
10H	Tomlinson (MRC)	Oligomeric a-synuclein 7-day aggregates	Dot blot, time course, ELISA, AFM, western blot analysis	Human PD brain tissue, Mouse PD brain tissue, (Dot blot assays, western blot, Immunohistochemistry with tissue and cells	amino acid SEQ ID NO: 5; nucleic acid SEQ ID NO: 12
6E	Tomlinson (MRC)	Fibrillar aggregates (likely not protein specific)	Dot blot, time course, ELISA, AFM	Human PD brain tissue, Human CSF, Mouse PD brain tissue, (Dot blot assays, western blot, Immunohistochemistry with tissue and cells	amino acid SEQ ID NO: 6; nucleic acid SEQ ID NO: 13
D10	Tomlinson (MRC)	All forms of a-synuclein	Dot blot, time course, ELISA, AFM, western	Human PD brain tissue, Human CSF, Mouse PD brain tissue, (Dot blot assays, western blot, Immunohistochemistry with tissue and cells	amino acid SEQ ID NO: 7; nucleic acid SEQ ID NO: 14
BSEC1	PNRL (Pacific National Research Labs?) Yeast library	BACE1 cleavage site on APP (Does not bind soluble Abeta	BACE-1 catalytic assay	Cell toxicity, cell culture assays	

In certain embodiments, the C6 nanobody has a sequence of SEQ ID NO:1:

```

EXPIAYGSRWIVITRGPAGHGPGTAAGVGGGLVQPGGSLRLSCAASGF
TFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTISRDN
KNTLYLQMNLSRAEDTAVYYCAKSYGKSVKISCFDYWGQSTLVTVSSG
GGGSGGGGGGGSEIVLTQSPDSLAVSLGERATINCKSSQSVLYNSN
NKNYLAWYQQKPGQSPPELLIYWASTRESGVPDRFSGSGSGTEFTLTIS
LQAEDVAVYYCQQFYSTPPTFGQGTKLEIKRAAAHHHHHGAEEQKL
ISEED

```

In certain embodiments, the antibody fragment comprises amino acid residues 16-292 of SEQ ID NO:1. In certain embodiments, the antibody fragment comprises or consists of amino acid sequence SEQ ID NO:1. In certain embodiments, the antibody fragment is less than 500 amino acids in length, such as between 200-450 amino acids in length, or less than 300 amino acids in length.

Identification of Isolated A β 40/42 Fractions that React with Existing Morphology-Specific Nanobodies.

Several different nanobodies to different A β morphologies including ones that selectively bind only fibrillar A β (Marcus, W. D., et al., Characterization of an antibody scFv that recognizes fibrillar insulin and beta-amyloid using atomic force microscopy. *Nanomedicine*, 2008. 4: p. 1-7) or three different oligomeric species of A β (Zameer, A., et al., Anti-oligomeric Abeta single-chain variable domain antibody blocks Abeta-induced toxicity against human neuroblastoma cells. *J Mol Biol*, 2008. 384: p. 917-28; Kasturirangan, S., et al., Nanobody specific for oligomeric beta-amyloid stabilizes non-toxic form. *Neurobiol Aging*, 2010. In press; Kasturirangan, S., et al., Isolation and Characterization of a Nanobody that Selectively Binds Brain Derived

30 Oligomeric Beta-Amyloid. (Submitted)) have been developed by the inventors. This existing pool of morphology specific nanobodies is used to verify that distinct A β species are separated in the DC-iGEP protocols, and to identify 35 novel A β species for which there are not selective reagents.

Separation and Collection of A β Fractions.

A β fractions are isolated and concentrated along the sawtooth gates. Each A β fraction is captured within the so called "cell" (the area between the gates) (Staton, S. J. R., et al., Characterization of particle capture in a sawtooth patterned insulating electrokinetic microfluidic device. *Electrophoresis*, 2010. 31: p. 3634-3641). An additional design feature is needed to keep the collected targets within a confined volume, yet accessible to various reagents. Lateral channels are used that are "electrically silent" when the 45 electric field is placed across the long axis of the device. By placing electrodes in reservoirs placed above and below the device and corresponding channels, the captured A β species for subsequent analysis is recovered.

Reactivity of A β Fractions with Nanobody.

There are numerous techniques that are used to determine binding specificity of each of the nanobodies isolated against the different target A β morphologies depending on the availability and stability of the target antigen. In characterizing binding specificity to the different A β species isolated by DC-iGDEP, most A β species are present in only very small amounts and low volume. For these cases, reactivity of the existing species is analyzed using AFM, which utilizes minimal material without the need for any modification. For those A β morphologies that are obtained in reasonable quantity, binding specificity is determined by ELISA, western or dot blot, depending on how easy it is to purify the target aggregate morphology. The protocols for each of these assays are routinely used (Emadi, S., et al., Inhibiting 65 Aggregation of alpha-Synuclein with Human Single Chain Antibody Fragments. *Biochemistry*, 2004. 43: p. 2871-2878; Zhou, C., et al., A human single-chain Fv intrabody blocks

aberrant cellular effects of overexpressed alpha-synuclein. *Mol Ther*, 2004. 10: p. 1023-31; Liu, R., et al., Single chain variable fragments against beta-amyloid (Abeta) can inhibit Abeta aggregation and prevent abeta-induced neurotoxicity. *Biochemistry*, 2004. 43: p. 6959-67; Zameer, A., et al., Single Chain Fv Antibodies against the 25-35 Abeta Fragment Inhibit Aggregation and Toxicity of Abeta42. *Biochemistry*, 2006. 45: p. 11532-9; Emadi, S., et al., Isolation of a human single chain antibody fragment against oligomeric alpha-synuclein that inhibits aggregation and prevents alpha-synuclein-induced toxicity. *J Mol Biol*, 2007. 368: p. 1132-44; Liu, R., et al., Residues 17-20 and 30-35 of beta-amyloid play critical roles in aggregation. *J Neurosci Res*, 2004. 75: p. 162-71; Liu, R., et al., Proteolytic antibody light chains alter beta-amyloid aggregation and prevent cytotoxicity. *Biochemistry*, 2004. 43: p. 9999-10007). In certain circumstances, the nanobody specificity for some of the oligomeric A β samples may not be able to be determined by conventional means such as western blot as described above. Several different AFM based methods are used to determine antibody specificity for antigen targets that are not suitable for analysis as described above, or that are available in only limited amounts. For example, nanobody specificity by height distribution analysis (Wang, M. S., et al., Characterizing Antibody Specificity to Different Protein Morphologies by AFM. *Langmuir*, 2008) or by recognition imaging (Marcus, W. D., et al., Isolation of an scFv targeting BRG1 using phage display with characterization by AFM. *Biochem Biophys Res Commun*, 2006. 342: p. 1123-9).

At least five A β fractions are separated, including monomeric, several oligomeric and a fibrillar form. In certain situations if the lateral channels are not sufficiently electrically silent or they are accessible under the diffusive conditions during processing, a simple pneumatic valve concept is used (Unger, M. A., et al., Monolithic microfabricated valves and pumps by multilayer soft lithography. *Science*, 2000. 288: p. 113-116), where the roof of the lateral channels are closed by applying pressure because the device is made of elastic polymers. Pressure is nominal, and can be applied mechanically.

The invention will now be illustrated by the following non-limiting Examples.

Example 1

Here we report a novel method for the manipulation and concentration of A β amyloid fibrils, implicated in Alzheimer's disease, using DC insulating gradient dielectrophoresis (DC-iGDEP). Fibril enrichment was found to be ~400%. Simulations suggest that capture of the full range of amyloid protein aggregates is possible with optimized device design.

Protein aggregates have been associated with more than 25 human diseases, including Alzheimer's disease, Parkinson's disease and type 2 diabetes. As a result, researchers in medical, biochemical and analytical chemistry fields are striving to understand the mechanisms of formation of amyloid aggregates and to identify toxic species. Protofibrils, transient metastable oligomeric aggregates that develop prior to the formation of amyloid fibrils, have been implicated in disease pathogenesis and are considered to be cytotoxic. Amyloid aggregation has been shown to form a heterogeneous mixture of oligomeric species in vitro, but the specific structure or structures of the cytotoxic species, their relationship to amyloid fibril formation and the mechanism of toxicity are unknown. Techniques capable of manipulating and concentrating various aggregate structures are of paramount importance because specific knowledge of all

constituent aggregate species is needed to effectively develop therapeutics based on disrupting or altering the aggregation process. Each of the different aggregate species from monomer to mature fibrils can have unique chemical and physical properties providing a basis for differential toxicity.

Traditional means of isolating various oligomers and fibrils of A β amyloid and other misfolded proteins have relied heavily on techniques such as sedimentation, size-exclusion chromatography (SEC) and capillary electrophoresis (CE). Sedimentation typically uses large volume preparations and cannot effectively separate oligomers into subpopulations, given their structural similarities and the typical small volumes of aggregate samples. While CE uses small volume samples and provides greater resolution compared to other separation techniques, target populations generally are diluted rather than concentrated during separation. Similarly, SEC has proven effective for isolating relatively stable protofibril species, but as with CE, samples are generally diluted during separation, and it has been shown to disrupt some protein aggregates. Several other methods used to isolate and characterize A β aggregates include HPLC, gel electrophoresis, and transmission electron microscopy (TEM); however, these methods do not provide a comprehensive account of the coexisting oligomeric populations during amyloid fibril formation. TEM measurements are made after extensive sample preparation and generally only capture stable end-point species. Considering amyloid fibril formation as a series of reversible reactions suggests that all separation schemes can alter the populations of aggregate species present before separation. For instance, protofibrils can convert back to monomers. To provide the most realistic view of the population of aggregates in a sample, a separation-based technique should be as rapid and gentle as possible.

The technique described here, direct current based insulator gradient dielectrophoresis (DC-iGDEP), rapidly resolves particulates by balancing electrophoresis and dielectrophoresis within a single separation channel. (S. J. R. Staton, K. P. Chen, T. J. Taylor, J. R. Pacheco and M. A. Hayes, *Electrophoresis*, 2010, 31, 3634-3641.) Relative to a capillary electrophoretic separation, the addition of dielectrophoretic forces increases the analyte-specific separation vectors to include the permeability and conductivity of the particle as well as the same parameters for the surrounding medium, which can be tuned. Combining these forces in opposition translates into discrete collection points that also concentrate the target species. This achieves both the separation and concentration of A β amyloid structures in a short time frame for direct quantification or use in downstream research on the properties of those structures. The separation is not based on the interaction of the aggregates with a stationary phase, reducing the impact of the separation process on the aggregate structures.

Previously, it has been demonstrated that DC-iGDEP can separate and concentrate spherical synthetic polymer nano- and microparticles from 20 nm to 1 μ m. (S. J. R. Staton, K. P. Chen, T. J. Taylor, J. R. Pacheco and M. A. Hayes, *Electrophoresis*, 2010, 31, 3634-3641; M. D. Pysher and M. A. Hayes, *Anal. Chem.* 2007, 79, 4552-4557.) It was not known if DC-iGDEP could be used successfully to manipulate species with the size, shape and chemical properties of amyloid aggregates. Here we present initial studies demonstrating that DC-iGDEP is able to rapidly and selectively concentrate A β amyloid aggregates, and has the potential to analyze populations of smaller protofibrillar aggregates with further tailoring of the channel design.

The design and fabrication of the DC-iGDEP device is detailed in Staton et al. (S. J. R. Staton, K. P. Chen, T. J. Taylor, J. R. Pacheco and M. A. Hayes, *Electrophoresis*, 2010, 31, 3634-3641) Briefly, the DC-iGDEP microfluidic channel was fabricated in polydimethylsiloxane (PDMS) with a glass cover slide (FIG. 3), and 20 μ L of A β sample was introduced into the reservoir at the broader end of the channel. The A β monomer and fibrils in PBS were prepared as described in detail previously. (B. O’Nuallain, A. K. Thakur, A. D. Williams, A. M. Bhattacharyya, S. Chen, T. Geetha and R. Wetzel, *Methods Enzymol.*, 2006, 413, 34-74; R. Picou, J. P. Moses, A. D. Wellman, I. Kheterpal and S. D. Gilman, *Analyst*, 2010, 135, 1631-1635; I. Kheterpal, K. D. Cook and R. Wetzel, *Methods Enzymol.* 2006, 39, 584-593; R. A. Picou, I. Kheterpal, A. D. Wellman, M Minnamreddy, G. Ku and S. D. Gilman, *J. Chromatogr. B*, 2011, 879, 627-632.) Additional information about the fabrication of the DC-iGDEP and A β sample preparation can be found in Example 2. The monomer and fibril samples were tested separately by DC-iGDEP. The results reported here were consistent and representative of multiple experiments using three independent preparations of the monomer and fibrils and multiple devices. Experiments were performed by applying voltages between 400 to 1000 V (over the entire channel length) for 1 to 15 min for each experiment.

The DC-iGDEP experiments were monitored by fluorescence microscopy. When introduced into the DC-iGDEP channel, the monomer consistently created streaming patterns at all of the applied voltages (400 to 1000 V, FIG. 4), which were more apparent at higher voltages. The streamlines appearing along the centerline of the channel indicate significant influence of electrophoretic forces; however, none of the conditions tested resulted in the monomers being captured.

In contrast, when the A β amyloid fibril samples were tested using the DC-iGDEP channel under similar conditions, separation and concentration of the fibrils were observed (FIG. 5). A TEM of the fibril sample used in this experiment is presented in FIG. 6 and clearly shows the presence of mature amyloid fibrils. Thioflavin T (10 μ M) fluorescence measured that aggregation was complete after 7 days. Images of experiments below 400 V applied over the length of the channel indicated movement of fibrils without capture. At 400-1000 V applied, the A β fibrils were captured and concentrated at the narrow points in the channel with similar local fluorescence intensity. Previous studies with synthetic nanospheres have demonstrated that particle capture typically does not obstruct the channels, and nanoparticles below the capture threshold size will pass through an intersection containing larger particles that have been concentrated and captured. (S. J. R. Staton, K. P. Chen, T. J. Taylor, J. R. Pacheco and M. A. Hayes, *Electrophoresis*, 2010, 31, 3634-3641.)

In order to differentiate between nonspecific and DC-iGDEP controlled capture, the applied voltage was removed after the capture and concentration of A β fibrils, and bright areas were monitored for dispersion away from the capture area. The bright fluorescence regions shown in FIG. 5 rapidly dissipated due to diffusion and slight convection of the A β fibrils away from the capture regions. Following a short period of time, the voltage was reapplied to allow selective capture to occur again. This process was repeated several times to verify that the capture was the result of the DC-iGDEP as well as to demonstrate the ability of the technique to reproducibly manipulate the amyloid aggregates within a given experimental session. The fibrils controllably collected near tips of the sawtooth-patterned insu-

lating channel. The position of capture indicates that, under these experimental conditions, the fibrils exhibited properties consistent with positive dielectrophoresis, meaning that the collection points were where the electric field intensity is greatest. (S. J. R. Staton, K. P. Chen, T. J. Taylor, J. R. Pacheco and M. A. Hayes, *Electrophoresis*, 2010, 31, 3634-3641; H. A. Pohl, *Dielectrophoresis*, Cambridge University Press, 5 Cambridge, 1978; K. P. Chen, J. R. Pacheco, M. A. Hayes and S. J. R. Staton, *Electrophoresis*, 2009, 30, 1441-1448.) According to classic theory, particles that undergo positive dielectrophoretic capture are less permeable than the surrounding medium. Some evidence of nonspecific adsorption of the amyloid aggregates was observed along the channel surface. Without surface treatments or coatings nonspecific protein adsorption is common for PDMS channels.

The ability of the DC-iGDEP method to concentrate fibrils was examined semi-quantitatively by noting the fluorescence intensity at the collection points compared to background levels in the channel prior to collection and areas where no detectable capture had occurred. The resulting enrichment of the fibril material ranged from about 350% to over 500% depending on the applied voltage (see Example 2). However, under more ideal conditions with lower fibril loads, the potential capture efficiency could be as high as 600%, simply by reducing the amount of background fluorescence. The maximum observed enrichment was 520% at 600 V.

The overall goal of this work is to separate, capture and concentrate the full range of aggregate structures generated during mature amyloid fibril formation. The experimental results presented here demonstrate that mature A β fibrils, but not A β monomer, are captured and concentrated using the current device design and the described experimental conditions. COMSOL (finite element multiphysics modeling software) calculations allow for directed device design development by exploring alternative device design parameters not yet tested experimentally. Modeling with COMSOL confirmed that for the current DC-iGDEP device design, A β monomer should not be captured for any reasonable experimental conditions. Details of the models used to evaluate the particle capture potential of the device can be found in Example 2. Modeling also predicts that by reducing the smallest sawtooth gap distance from 27 μ m in the current design to \sim 10 nm, the increased field gradient strength required to allow monomer capture along with all intermediate species could be generated. By modeling the amyloid aggregates, oligomers, and monomers based on their unique electrophoretic and dielectrophoretic properties, the speed of the device design evolution can be accelerated while also being tailored for capturing particular bioanalytes and positioning them along the length of the channel.

In conclusion, this study demonstrates that DC-iGDEP can be used to manipulate and selectively capture A β amyloid fibrils, while influencing but not capturing A β monomer. DC-iGDEP successfully combines high enrichment of fibrils (up to 520%) with short analysis times (1-15 min) on a cost effective platform. Production and operation of the DC-iGDEP microdevices using PDMS are very simple. This initial study and related simulations indicate that this technique has the potential to rapidly isolate and concentrate various A β aggregate structures intermediate between monomer and mature fibrils. Development of new rapid and gentle separation techniques for isolating and characterizing amyloid aggregates is essential for understanding the role of protein aggregation in amyloid-linked diseases.

A) Details of DC-iGDEP Device Fabrication

The DC-iGDEP devices were fabricated utilizing standard photolithography, fabrication, and bonding techniques. (C. Mack, *Fundamental Principles of Optical Lithography: The Science of Microfabrication*, Wiley, Hoboken, 2008.) Photomasks were designed in AutoCAD (Autodesk; San Rafael, Calif., USA), and photolithographic positive stamps were made using AZ P4620 photoresist (AZ Electronic Materials; Branchburg, N.J., USA) and contrast enhancement material CEM388SS (Shin-Etsu MicroSi; Phoenix, Ariz., USA). Device channels were fabricated from polydimethylsiloxane (PDMS) with a microscope slide coverplate. The PDMS channels were cast using Sylgard 184 silicone elastomer kit PDMS (Dow/Corning; Midland, Mich., USA). Shortly after the PDMS portion of the device was fabricated, access holes were made using a hole punch (3 mm diameter through 0.5-1 cm of PDMS), and then the PDMS portion of the device was sealed to the glass cover plate by plasma oxidation followed by contact sealing. (K. Haubert, T. Drier and D. Beebe, *Lab Chip*, 2006, 6, 1548-1549.) The geometry of the separatory portion of the DC-iGDEP channel consisted of successive triangular units that extended into the open volume to induce local electric field gradients. The insulating PDMS 60° triangles began with a base length of 6 μm and a height of 5.2 μm. Their side length and width increased by 40 μm after every six repeats (FIG. 3), resulting in an initial gap distance of 945 μm and a final gap distance of 27 μm. The separatory portion of the DC-iGDEP channel connected the two reservoirs created by the hole punch, where sample and buffer were introduced into the channel. The channel depth ranged from 13 to 16 μm.

Sample was introduced via the reservoir at the end of the channel with the larger gap distance. After sample introduction, platinum wire electrodes (0.404 mm diameter, 99.9% purity; Alfa Aesar; Ward Hill, Mass., USA) were placed in each of the reservoirs in contact with the solution and attached to a power supply (Series 225, Bertram). The voltage was applied at a potential between 0 and 1000 V, for 1-15 min depending upon the experiment. Visualization was achieved using an Olympus inverted IX70 microscope with a mercury short arc H30 103 w/2 light source from OSRAM and an Olympus DAPI, FITC, Texas Red triple band pass cube (Olympus; Center Valley, Pa., USA).

Videos and still images were collected with a monochrome QICAM cooled CCD camera (QImaging, Inc.; Surrey, BC, Canada) and Streampix III image capture software (Norpix, Inc.; Montreal, QC, Canada). The fluorescence intensity was then analyzed with Image? (NIH; Bethesda, Md.). Three different regions of interest (ROI) were selected (FIG. 7). The ROIs represent the capture zone (gate), a background area within the channel (recess), and a background outside of the channel (PDMS). Table 2 uses the same ROIs described in FIG. 7. The data presented in Table 3 represent multiple replicates ($n > 3$) with isolation events occurring in the same position. Ratios of the various ROIs were utilized to determine enrichment of fluorescent fibrils in the capture zone when compared to the rest of the microfluidic channel filled with sample as well as a background of the device outside of the channel. Each ROI represents equally sized areas.

Fluorescence Intensity Values in ROIs for Fibril Samples.				
	Photo label	Intensity/Area	Ratio	
400 V				
Gate	1	212.8		
Recess	2	52.2	Gate/Recess	4.1
PDMS	3	35.9	Gate/PDMS	5.9
600 V				
Gate	1	207.0		
Recess	2	40.0	Gate/Recess	5.2
PDMS	3	36.1	Gate/PDMS	5.7
800 V				
Gate	1	214.5		
Recess	2	60.0	Gate/Recess	3.6
PDMS	3	36.1	Gate/PDMS	5.9
1000 V				
Gate	1	211.7		
Recess	2	61.2	Gate/Recess	3.5
PDMS	3	36.1	Gate/PDMS	5.9

B) Preparation of A β (1-40) Monomers and Fibrils

Both A β (1-40) monomer and fibril samples were prepared as described in detail previously. (R. Picou, J. P. Moses, A. D. Wellman, I. Kheterpal and S. D. Gilman, *Analyst*, 2010, 135, 1631-1635; R. A. Picou, I. Kheterpal, A. D. Wellman, M. Minnamreddy, G. Ku and S. D. Gilman, *J Chromatogr. B*, 2011, 879, 627-632; B. O'Nuallain, A. K. Thakur, A. D. Williams, A. M. Bhattacharyya, S. Chen, G. Thiagarajan and R. Wetzel, in *Methods Enzymol.* 2006, 413, 34-74.) Briefly, A β (1-40) peptide (W. M Keck Foundation Biotechnology Research Laboratory, Yale University; New Haven, Conn.) and carboxyfluorescein (FAM) labeled A β (1-40) peptide (Anaspec Inc.; Fremont, Calif.) were first treated with trifluoroacetic acid (TFA) and hexafluoroisopropanol (HFIP) to remove any preexisting aggregates. For A β monomer samples, the solvent was evaporated off, and the peptides were dissolved in 10.00 mM Tris at pH 7.79. FAM-A β (1-40) monomer was mixed with A β (1-40) monomer at a mass ratio of 1:4. The total A β concentration of the monomer solution was determined to be 30 μM using a Shimadzu HPLC-UV instrument with detection at 215 nm as described previously. (R. Picou, J. P. Moses, A. D. Wellman, I. Kheterpal and S. D. Gilman, *Analyst*, 2010, 135, 1631-1635; R. A. Picou, I. Kheterpal, A. D. Wellman, M. Minnamreddy, G. Ku and S. D. Gilman, *J Chromatogr. B*, 2011, 879, 627-632; B. O'Nuallain, A. K. Thakur, A. D. Williams, A. M. Bhattacharyya, S. Chen, G. Thiagarajan and R. Wetzel, in *Methods Enzymol.* 2006, 413, 34-74.)

For the A β (1-40) fibril samples, TFA was evaporated off, and the peptides were dissolved in HFIP. The concentration of each peptide was determined using HPLC-UV. FAM-A β (1-40) monomer was mixed at a mass ratio of 1:4 with A β (1-40) monomer. HFIP was evaporated off, and the peptide mixture was dissolved stepwise in equal volumes of 2.0 mM NaOH and 2× phosphate buffered saline (PBS) containing 22.8 mM phosphate, 274 mM NaCl, 5.4 mM KCl and 0.1% NaN₃ at pH 7.4. The samples were centrifuged at 50,000 g for a minimum of 10 hat 4° C. Fibril formation was initiated by addition of a small quantity (0.1% by weight of total A β monomer) of fibrillar aggregates to the supernatant from a previous fibril synthesis. The mixture was then incubated at 37° C. for 7 d. Depletion in monomer during fibril formation was monitored using HPLC-UV with detection at 215 nm as described previously. (R. Picou, J. P. Moses, A. D. Wellman,

I. Kheterpal and S. D. Gilman, *Analyst*, 2010, 135, 1631-1635; R. A. Picou, I. Kheterpal, A. D. Wellman, M. Minnamreddy, G. Ku and S. D. Gilman, *J Chromatogr. B*, 2011, 879, 627-632; B. O’Nuallain, A. K. Thakur, A. D. Williams, A. M. Bhattacharyya, S. Chen, G. Thiagarajan and R. Wetzal, in *Methods Enzymol.* 2006, 413, 34-74.) Fibril growth was monitored using ThT fluorescence until complete (5-7 days), and the quality of fibrils was assessed by electron microscopy. For direct comparison with A β (1-40) monomer samples in DC-iGDEP analysis, fibril samples were buffer exchanged from PBS to 10.00 mM Tris electrophoresis buffer at pH 7.79. (I. Kheterpal, S. Zhou, K. D. Cook and R. Wetzal, *Proc. Natl. Acad. Sci. USA*, 2000, 97, 13597-13601.) The monomer-equivalent concentrations of all samples were determined to be 30 μ M by HPLC-UV.

C) COMSOL Mathematical Modeling

In order to aid in understanding and interpreting particle behavior within the sawtooth-patterned microchannel, the electric field distribution within the microdevice was determined numerically and plotted. This was accomplished using finite element analysis software COMSOL Multiphysics 4.1 (COMSOL, Inc., Burlington, Mass.) in the “conductive media DC” mode. Models created in this mode are time-independent calculations of field strength within an aqueous medium. COMSOL computes the electric field by solving the Laplace equation, $\nabla^2(\varphi)=0$, for potential distribution at various points throughout the channel, along a predefined mesh. The boundary conditions are defined as distinct potentials at the channel inlet and outlet and electrical insulators at the channel walls.

A properly scaled model of the main channel geometry was produced and imported into the COMSOL environment. In order to simplify the model, a 2D approximation of the channel was utilized. Due to the placement of electrodes in distant reservoirs and the high width to depth ratio of the channel, the effects of channel depth on electrical potential were neglected. The effects of particles on the electric field distribution were also neglected.

Reservoir-channel junctions were set to predetermined potentials similar to those used during experiments. V_{inlet} and V_{outlet} were 0 V and -550 V, respectively. All other boundaries (representing PDMS and glass walls) were set to be perfect insulators; an approximation justified by the large difference in conductivity between the fluid medium and the channel walls. The conductivity and relative permittivity of the medium were set at 1.2 S/m and 78, respectively. This conductivity is consistent with values that can be obtained using standard phosphate-buffered saline.

A triangular mesh was applied to the entire channel area. The mesh contained approximately 97,000 triangular elements and 680 vertex elements. Through finite element analysis, the software approximates the electric potential at each mesh point. From these numerical values other useful parameters relevant to electrokinesis and dielectrophoresis may be determined, such as electric field strength, $|E|$, and the gradient of square of the electric field, $\nabla|E|^2$. Built-in tools enable graphic representation of the resulting data.

Since dielectrophoretic force is proportional to $\nabla|E|^2$, the magnitude of this term along the channel centerline was extracted from the COMSOL model using a 2D cut line. Localized maxima from this dataset correspond to channel constrictions or gates along the sawtooth pattern. When these values of $\nabla|E|^2$ are plotted as a function of gate width, the relationship can be reasonably approximated with a power function (FIG. 8). Visualizing the data in this manner illustrates the rapid scaling of dielectrophoretic force with increasing geometric constriction of the channel.

Amongst other things, dielectrophoretic force is proportional to particle size. Thus, capture of smaller particles such as protein monomers will require larger values of $\nabla|E|^2$. FIG. 8 illustrates that drastically larger values of $\nabla|E|^2$ are attainable by increasing the constriction ratio at gates. Using the following assumptions and the $\nabla|E|^2$ values determined in FIG. 8, it was found that a gate width of ~10 nm would be sufficient to capture A β monomers. The assumptions used in the calculation were that there was negligible electroosmotic flow, the A β monomer is spherical with a diameter of ~2.5 nm, the monomer μ EP is 1.2×10^{-4} cm²/(Vs), the viscosity and permittivity of the medium are similar to that of water, and the conductivity of the A β monomer is significantly less than that of the fluid medium.

Although the foregoing specification fully discloses and enables the present invention, it is not intended to limit the scope of the invention, which is defined by the claims appended hereto.

All publications, patents and patent applications are incorporated herein by reference. While in the foregoing specification this invention has been described in relation to certain embodiments thereof, and many details have been set forth for purposes of illustration, it will be apparent to those skilled in the art that the invention is susceptible to additional embodiments and that certain of the details described herein may be varied considerably without departing from the basic principles of the invention.

The use of the terms “a” and “an” and “the” and similar referents in the context of describing the invention are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms “comprising,” “having,” “including,” and “containing” are to be construed as open-ended terms (i.e., meaning “including, but not limited to”) unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

Embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Variations of those embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 79

<210> SEQ ID NO 1

<211> LENGTH: 292

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (2)..(2)

<223> OTHER INFORMATION: Any amino acid

<400> SEQUENCE: 1

Glu Xaa Pro Ile Ala Tyr Gly Ser Arg Trp Ile Val Ile Thr Arg Gly
1 5 10 15

Pro Ala Gly His Gly Pro Gly Thr Ala Ala Gly Val Gly Gly Gly Leu
20 25 30

Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe
35 40 45

Thr Phe Ser Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys
50 55 60

Gly Leu Glu Trp Val Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr
65 70 75 80

Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser
85 90 95

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr
100 105 110

Ala Val Tyr Tyr Cys Ala Lys Ser Tyr Gly Ser Val Lys Ile Ser Cys
115 120 125

Phe Asp Tyr Trp Gly Gln Ser Thr Leu Val Thr Val Ser Ser Gly Gly
130 135 140

Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Ile Val
145 150 155 160

Leu Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly Glu Arg Ala
165 170 175

Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Asn Ser Asn Asn
180 185 190

Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Glu
195 200 205

Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val Pro Asp Arg
210 215 220

Phe Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser
225 230 235 240

Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Phe Tyr Ser
245 250 255

Thr Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Ala
260 265 270

Ala Ala His His His His His His Gly Ala Ala Glu Gln Lys Leu Ile
275 280 285

Ser Glu Glu Asp
290

<210> SEQ ID NO 2

<211> LENGTH: 268

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 2

Met Ala Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro
 1 5 10 15

Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 20 25 30

Ser Tyr Pro Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu
 35 40 45

Trp Val Ser Ala Ile Gln His Thr Gly Ala Pro Thr Thr Tyr Ala Asp
 50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
 65 70 75 80

Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
 85 90 95

Tyr Cys Ala Lys Ala Phe Pro Pro Phe Asp Tyr Trp Gly Gln Gly Thr
 100 105 110

Leu Val Thr Val Ser Ser Gly Gly Gly Ser Gly Gly Gly Ser
 115 120 125

Gly Gly Gly Gly Ser Thr Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
 130 135 140

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 145 150 155 160

Gln Ser Ile Ser Ser Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys
 165 170 175

Ala Pro Lys Leu Leu Ile Tyr Ser Ala Ser Ser Leu Gln Ser Gly Val
 180 185 190

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 195 200 205

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 210 215 220

Arg Glu Thr Gly Pro Lys Ala Phe Gly Gln Gly Thr Lys Val Glu Ile
 225 230 235 240

Lys Arg Ala Ala Ala His His His His His His Gly Ala Ala Glu Gln
 245 250 255

Lys Leu Ile Ser Glu Glu Asp Leu Asn Gly Ala Ala
 260 265

<210> SEQ ID NO 3

<211> LENGTH: 268

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 3

Met Ala Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro
 1 5 10 15

Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 20 25 30

Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu
 35 40 45

Trp Val Ser Ser Ile Gln Pro Glu Gly Arg Arg Thr Ala Tyr Val Asp
 50 55 60

-continued

```

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
65          70          75          80
Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
85          90          95
Tyr Cys Ala Lys Pro Pro Glu Arg Phe Asp Tyr Trp Gly Gln Gly Thr
100         105         110
Leu Val Thr Val Ser Ser Gly Gly Gly Ser Gly Gly Gly Gly Ser
115         120         125
Gly Gly Gly Gly Ser Thr Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
130         135         140
Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
145         150         155         160
Gln Ser Ile Ser Ser Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys
165         170         175
Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val
180         185         190
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
195         200         205
Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
210         215         220
Ser Tyr Ser Thr Pro Asn Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
225         230         235         240
Lys Arg Ala Ala Ala His His His His His His Gly Ala Ala Glu Gln
245         250         255
Lys Leu Ile Ser Glu Glu Asp Leu Asn Gly Ala Ala
260         265

```

```

<210> SEQ ID NO 4
<211> LENGTH: 268
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide
<400> SEQUENCE: 4

```

```

Met Ala Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro
1          5          10          15
Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
20         25         30
Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu
35         40         45
Trp Val Ser Ser Ile Gly Gln Lys Gly Gly Gly Thr Gln Tyr Ala Asp
50         55         60
Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
65          70          75          80
Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
85          90          95
Tyr Cys Ala Lys His Phe Glu Asn Phe Asp Tyr Trp Gly Gln Gly Thr
100         105         110
Leu Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
115         120         125
Gly Gly Gly Gly Ser Thr Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
130         135         140
Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser

```


-continued

145	150	155	160
Gln Ser Ile Ser Ser Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys	165	170	175
Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser His Leu Gln Ser Gly Val	180	185	190
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr	195	200	205
Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln	210	215	220
Thr Arg Arg Pro Pro Ser Thr Phe Gly Gln Gly Thr Lys Val Glu Ile	225	230	235
Lys Arg Ala Ala Ala His His His His His His Gly Ala Ala Glu Gln	245	250	255
Lys Leu Ile Ser Glu Glu Asp Leu Asn Gly Ala Ala	260	265	

<210> SEQ ID NO 5
 <211> LENGTH: 281
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polypeptide

<400> SEQUENCE: 5

Met Ala Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro	1	5	10	15
Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser	20	25	30	
Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu	35	40	45	
Trp Val Ser Asn Ile Ser Ser Ala Gly Lys Gly Leu Glu Trp Val Ser	50	55	60	
Ser Ile Asp Asp Ser Gly Ala Ser Thr Tyr Tyr Ala Asp Ser Val Lys	65	70	75	80
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu	85	90	95	
Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala	100	105	110	
Lys Asp Ser Ala Ser Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr	115	120	125	
Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly	130	135	140	
Gly Ser Thr Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala	145	150	155	160
Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile	165	170	175	
Ser Ser Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys	180	185	190	
Leu Leu Ile Tyr Thr Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg	195	200	205	
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser	210	215	220	
Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Ala Ala	225	230	235	240

-continued

Ser Pro Ser Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Ala
 245 250 255

Ala Ala His His His His His His Gly Ala Ala Glu Gln Lys Leu Ile
 260 265 270

Ser Glu Glu Asp Leu Asn Gly Ala Ala
 275 280

<210> SEQ ID NO 6
 <211> LENGTH: 268
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polypeptide

<400> SEQUENCE: 6

Met Ala Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro
 1 5 10 15

Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 20 25 30

Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu
 35 40 45

Trp Val Ser Tyr Ile Ala Ser Gly Gly Asp Thr Thr Asn Tyr Ala Asp
 50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
 65 70 75 80

Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
 85 90 95

Tyr Cys Ala Lys Gly Ala Ser Ala Phe Asp Tyr Trp Gly Gln Gly Thr
 100 105 110

Leu Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 115 120 125

Gly Gly Gly Gly Ser Thr Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
 130 135 140

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 145 150 155 160

Gln Ser Ile Ser Ser Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys
 165 170 175

Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Tyr Leu Gln Ser Gly Val
 180 185 190

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 195 200 205

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 210 215 220

Ser Ser Asn Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
 225 230 235 240

Lys Arg Ala Ala Ala His His His His His His Gly Ala Ala Glu Gln
 245 250 255

Lys Leu Ile Ser Glu Glu Asp Leu Asn Gly Ala Ala
 260 265

<210> SEQ ID NO 7
 <211> LENGTH: 263
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polypeptide

-continued

<400> SEQUENCE: 7

```

Met Ala Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro
1           5           10           15

Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
20           25           30

Ser Tyr Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu
35           40           45

Trp Val Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp
50           55           60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
65           70           75           80

Leu Tyr Leu Gln Val Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
85           90           95

Tyr Cys Ala Arg Ile Asn Ala Lys Trp Gly Gln Gly Thr Leu Val Thr
100          105          110

Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Ser
115          120          125

Ala Leu Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser
130          135          140

Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser
145          150          155          160

Ser Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu
165          170          175

Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe
180          185          190

Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu
195          200          205

Gln Pro Gly Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Ser Thr
210          215          220

Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Ala Ala Ala
225          230          235          240

His His His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser Glu
245          250          255

Glu Asp Leu Asn Gly Ala Ala
260

```

<210> SEQ ID NO 8

<211> LENGTH: 877

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (715)..(718)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (725)..(725)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (729)..(729)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (733)..(733)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<220> FEATURE:

-continued

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (737)..(737)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (789)..(789)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (807)..(807)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (810)..(810)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (833)..(833)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (844)..(844)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (852)..(852)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (858)..(859)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (865)..(867)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 8

ttgttattac tcgcgcccca gccggccatg gccgaggtgc agctggtgga gtctggggga      60
ggcttggtac agcctggggg gtccttgaga ctctctgtg cagcctctgg attcaccttt      120
agcagctatc ccatgagctg ggtccgccag gctccagggg aggggctgga gtgggtctca      180
gcgattcagc atactggtgc gccgacaact tacgcagact ccgtgaaggg ccggttcacc      240
atctccagag acaattccaa gaacacgctg tatctgcaaa tgaacagcct gagagccgag      300
gacacggccg tatattactg tgcgaaagcg tttccgccgt ttgactactg gggccagggg      360
accctggtca ccgtctcgag cgggtggaggc ggttcaggcg gaggtggcag cggcggtggc      420
gggtcgacgg acatccagat gacctcagtct ccctcctccc tgtctgcatc tgtaggagac      480
agagtcacca tcacttgccg ggcaagtcag agcattagca gctatttaaa ttggtatcag      540
cagaaaccag ggaaagcccc taagctctctg atctattctg catcctcttt gcaaagtggg      600
gtcccatcaa gggtcagtg cagtggatct gggacagatt tcactctcac catcagcagt      660
ctgcaacctg aagatthtgc aacttactac tgtcaacagc gggagactgg gcctnnnngt      720
tcggncaang gancaangtg gaaatcaaac gggcgccgcg acatcatcat caccatcacg      780
gggccgcana acaaaaactc atctcanaan aggatctgaa tggggccgca tanactggtg      840
aaanttgttt ancaaacnnc atacnnaaaa ttcattt                                877

<210> SEQ ID NO 9
<211> LENGTH: 877
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (807)..(807)

```


-continued

```

<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (810)..(810)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (852)..(852)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 9

ttgttattac tcgcgcccca gccggcctgg ccgaggtgca gctggtggag tctgggggag      60
gcttggtaca gcctgggggg tccctgagac tctcctgtgc agcctctgga ttcaccttta      120
gcagctatgc catgagctgg gtccgccagg ctccagggaa ggggctggag tgggtctcat      180
ctattcagcc tgagggtagg cggacagcgt acgtagactc cgtgaagggc cggttcacca      240
tctccagaga caattccaag aacacgctgt atctacaaat gaacagcctg agagccgagg      300
acacggccgt atattactgt gcgaaaccgc cggagaggtt tgactactgg ggccagggaa      360
ccctggtcac cgtctcgagc ggtggaggcg gttcaggcgg aggtggcagc ggcggtgggc      420
ggtcgacgga catccagatg acccagtctc catcctccct gtctgcatct gtaggagaca      480
gagtcaccat cacttgccgg gcaagtcaga gcattagcag ctatttaaat tggtatcagc      540
agaaaaccagg gaaagcccct aagctcctga tctatgctgc atccagtttg caaagtgggg      600
tcccatcaag gttcagtggc agtggatctg ggacagattt cactctcacc atcagcagtc      660
tgcaacctga agattttgca acttactact gtcaacagag ttacagtacc cctaatacgt      720
tcggccaagg gaccaaggtg gaaatcaaac gggcgccgcg acatcatcat caccatcacg      780
gggccgcaga acaaaaactc atctcanaan aggatctgaa tggggccgca tagactgttg      840
aaagttgttt ancaaacctc atacagaaaa ttcattt      877

```

```

<210> SEQ ID NO 10
<211> LENGTH: 821
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

```

```

<400> SEQUENCE: 10

ccatggccca ggtacagctg caggagtcgg gggaggcttg gtacagcctg gggggtccct      60
gagactctcc tgtgcagcct ctggattcac ctttagcagc tatgccatga gctgggtccg      120
ccaggctcca ggaagggggc tggagtgggt ctcagctatt agtggtagtg gtggtagcac      180
atactacgca gactccgtga agggccgatt caccatctcc agagacaatt ccaagaacac      240
gctgtatctg caaatgaaca gcctgagagc tgaggacacg gctgtgtatt actgtgcgaa      300
gagctatggg tcagttaaaa taagctgctt tgactactgg ggccagagca ccctggtcac      360
cgtctcctca ggtggaggcg gttcaggcgg aggtggctct ggcggtgggc gatcggaat      420
tgtgtgacg cagtctccag actccctggc tgtgtctctg ggcgagaggg ccaccatcaa      480
ctgcaagtcc agccagagtg ttctttacaa ctccaacaat aagaactact tagcttggtg      540
ccagcagaaa ccaggacagt ctctgaggtt gctcatttac tgggcatcaa cccgggaatc      600
cggggtccct gaccgattca gtggcagcgg gtctgggaca gaattcactc ttaccatcag      660
cagcctgcag gctgaggatg tggcagttta ttactgtcag caattttata gtactcctcc      720
gacttttggc caggggacca agctggagat caaacgtgcg gccgcacatc atcatcacca      780

```


-continued

 tcacggggcc gcagaacaaa aactcatctc agaagaggat c 821

<210> SEQ ID NO 11
 <211> LENGTH: 822
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polynucleotide

<400> SEQUENCE: 11

```

ccatggccga ggtgcagctg ttggagtctg ggggaggctt ggtacagcct ggggggtccc   60
tgagactctc ctgtgcagcc tctggattca cctttagcag ctatgccatg agctgggtcc   120
gccaggctcc agggaagggg ctggagtggg tctcatcgat tggtcagaag ggtggtggta   180
cacagtacgc agactccgtg aagggccggt tcacatctc cagagacaat tccaagaaca   240
cgctgtatct gcaaatgaac agcctgagag ccgaggacac ggccgtatat tactgtgcca   300
aacattttga gaattttgac tactggggcc agggaaccct ggtcaccgtc tcgagcgggtg   360
gaggcggttc aggcggaggt ggcagcggcg gtggcgggtc gacggacatc cagatgaccc   420
agtctccatc ctccctgtct gcatctgtag gagacagagt caccatcact tgccgggcaa   480
gtcagagcat tagcagctat ttaaattggt atcagcagaa accagggaaa gccccctaagc   540
tctgatcta tgctgcatcc catttgcaaa gtgggggtccc atcaaggttc agtggcagtg   600
gatctgggac agatttcact ctcacatca gcagtctgca acctgaagat tttgcaactt   660
actactgtca acagacgcgt aggccgcctt ctacgttcgg ccaagggacc aaggtggaaa   720
tcaaacgggc ggccgcacat catcatcacc atcacggggc cgagaacaa aaactcatct   780
cagaagagaa tcactagtgc ggccgcctgc aggtcgacca ta                       822
  
```

<210> SEQ ID NO 12
 <211> LENGTH: 843
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polynucleotide

<220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (816)..(816)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (826)..(826)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (832)..(833)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (836)..(836)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (840)..(840)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 12

```

ccatggccga ggtgcagctg ttggagtctg ggggaggctt ggtacagcct ggggggtccc   60
tgagactctc ctgtgcagcc tctggattca cctttagcag ctatgccatg agctgggtcc   120
gccaggctcc agggaagggg ctggagtggg tctcaaatat tagtagtgca gggaaggggc   180
tggagtgggt ctcaagtatt gatgattctg gtgcttctac atattacgca gactccgtga   240
  
```


-continued

```

agggccgggt caccatctcc agagacaatt ccaagaacac gctgtatctg caaatgaaca 300
gctgagagac cgaggacacg gccgtatatt actgtgcgaa agattctgct tcttttgact 360
actggggcca ggaaccctg gtcaccgtct cgagcgggtg aggcgggtca ggcggaggtg 420
gcagcggcgg tggcgggtcg acggacatcc agatgaccca gtctccatcc tccctgtctg 480
catctgtagg agacagagtc accatcactt gccgggcaag tcagagcatt agcagctatt 540
taaattggta tcagcagaaa ccagggaaag ccctaagct cctgatctat actgcatcca 600
gtttgcaaag tgggggtcca tcaaggttca gtggcagtgg atctgggaca gatttcactc 660
tcaccatcag cagtctgcaa cctgaagatt ttgcaactta ctactgtcaa cagtctgctg 720
ctagtccttc tacgttcggc caagggacca aggtggaaat caaacgggcg gccgcacatc 780
accatcacca tcacggggcc gcagaacaaa aactcntctc agaagnggat cnaangggg 840
ccg 843

```

```

<210> SEQ ID NO 13
<211> LENGTH: 879
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (784)..(784)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (788)..(789)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (791)..(791)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (811)..(815)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (830)..(830)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (835)..(835)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (853)..(854)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (862)..(863)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (867)..(867)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (872)..(872)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (875)..(876)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 13

```

```

ttgttattac tcgcggcca gccggccatg gccgaggtgc agctgttga gtctggggga 60

```


-continued

```

ggcttggtac agcctggggg gtcctgaga ctctcctgtg cagcctctgg attcaccttt 120
agcagctatg ccatgagctg ggtccgccag gctccagga aggggctgga gtgggtctca 180
tatattgcta gtggtggtga tactacaaat tacgcagact ccgtaaggg ccggttcacc 240
atctccagag acaattccaa gaacacgctg tatctgcaa tgaacagcct gagagccgag 300
gacacggccg tatattactg tgcgaaaggt gcttctgctt ttgactactg gggccagga 360
accctggtca cgtctcgag cgggtggaggc ggttcaggcg gaggtggcag cggcgggtggc 420
gggtcgacgg acatccagat gaccagctct ccatcctccc tgtctgcatc tgtaggagac 480
agagtacca tcacttgccg ggcaagtacg agcattagca gctatttaa ttggtatcag 540
cagaaaccag ggaaagcccc taagctcctg atctatgctg catcctattt gcaaagtggg 600
gtcccatcaa ggttcagtgg cagtggatct gggacagatt tcactctcac catcagcagt 660
ctgcaacctg aagattttgc aacttactac tgtcaacaga gttctaata tccttatacg 720
ttcggccaag ggaccaagt ggaaatcaaa cggcggccg cacatcatca tcaccatcac 780
ggngccna naacaaaaac tcactcaaa nnnntctga atggggccn catanactgt 840
tgaaagttgt ttnnaacct cntacanaaa antcnnttt 879

```

```

<210> SEQ ID NO 14
<211> LENGTH: 776
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 14

```

```

atggccgagg tgcagctggt ggagtctggg ggaggcgtgg tccagcctgg gaggtccctg 60
agactctect gtgcagcctc tggattcacc ttcagtagct atggcatgca ctgggtccgc 120
caggccccag gcaaggggct ggagtgggtg gcagttatat catatgatgg aagtaataaa 180
tactatgcag actccgtgaa gggccgattc accatctcca gagacaattc caagaacacg 240
ctgtatctgc aagtgaacag cctgagagct gaggacacgg ccgtgtatta ctgtgcaaga 300
attaatgcga agtggggcca aggtaccctg gtcaccgtct cgagtgggtg aggcggttca 360
ggcggagggtg gctctggcgg tagtgcactt gacatccaga tgaccagtc tccatcctcc 420
ctgtctgcat ctgtaggaga cagagtcacc atcacttgcg gggcaagtca gagcattagc 480
agctatttaa attggtatca gcagaaacca gggaaagccc ctaagctcct gatctatgct 540
gcatccagtt tgcaaagtgg ggtcccatca aggttcagtg gcagtggatc tgggacagat 600
ttcactctca ccatcagcag tctgcaacct ggagattttg caacttacta ctgtcaacag 660
agttacagta ccccgacgtt cgggcaaggg accaaggtgg aatcaaacg tgccggccgca 720
catcatcatc accatcacgg ggccgcagaa caaaaactca tctcagaaga ggatct 776

```

```

<210> SEQ ID NO 15
<211> LENGTH: 278
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polypeptide

```

```

<400> SEQUENCE: 15

```

```

Met Ala Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Pro
1           5           10           15

```


-continued

Gly

<210> SEQ ID NO 18
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 peptide

<400> SEQUENCE: 18

Ser Tyr Gly Ser Val Lys Ile Ser Cys Phe Asp Tyr
 1 5 10

<210> SEQ ID NO 19
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 peptide

<400> SEQUENCE: 19

Lys Ser Ser Gln Ser Val Leu Tyr Asn Ser Asn Asn Lys Asn Tyr Leu
 1 5 10 15

Ala

<210> SEQ ID NO 20
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 peptide

<400> SEQUENCE: 20

Trp Ala Ser Thr Arg Glu Ser
 1 5

<210> SEQ ID NO 21
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 peptide

<400> SEQUENCE: 21

Gln Gln Phe Tyr Ser Thr Pro Pro Thr
 1 5

<210> SEQ ID NO 22
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 22

agctatgccca tgagc

15

<210> SEQ ID NO 23
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 23

gctattagtg gtagtggtgg tagcacatac tacgcagact ccgtgaaggg c 51

<210> SEQ ID NO 24
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 24

agctatgggt cagttaaaat aagctgcttt gactac 36

<210> SEQ ID NO 25
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 25

aagtcagcc agagtgttct ttacaactcc aacaataaga actacttagc t 51

<210> SEQ ID NO 26
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 26

tgggcatcaa cccgggaatc c 21

<210> SEQ ID NO 27
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 27

cagcaatttt atagtactcc tccgact 27

<210> SEQ ID NO 28
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 28

Ser Tyr Pro Met Ser
 1 5

<210> SEQ ID NO 29
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 29

Ala Ile Gln His Thr Gly Ala Pro Thr Thr Tyr Ala Asp Val Ser Lys
 1 5 10 15

Gly

<210> SEQ ID NO 30

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 30

Ala Phe Pro Pro Phe Asp Tyr
 1 5

<210> SEQ ID NO 31

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 31

Arg Ala Ser Gln Ser Ile Ser Ser Tyr Leu Asn
 1 5 10

<210> SEQ ID NO 32

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 32

Ser Ala Ser Ser Leu Gln Ser
 1 5

<210> SEQ ID NO 33

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 33

Gln Gln Arg Glu Thr Gly Pro Ile Leu
 1 5

<210> SEQ ID NO 34

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 34

Ser Ile Gln Pro Glu Gly Arg Arg Thr Ala Tyr Val Asp Ser Val Lys
 1 5 10 15

Gly

<210> SEQ ID NO 35

<211> LENGTH: 7

-continued

<212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 35

Pro Pro Glu Arg Phe Asp Tyr
 1 5

<210> SEQ ID NO 36
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 36

Ala Ala Ser Ser Leu Gln Ser
 1 5

<210> SEQ ID NO 37
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 37

Gln Gln Ser Tyr Ser Thr Pro Asn Thr
 1 5

<210> SEQ ID NO 38
 <211> LENGTH: 30
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 38

Asn Ile Ser Ser Ala Gly Lys Gly Leu Glu Trp Val Ser Ser Ile Asp
 1 5 10 15
 Asp Ser Gly Ala Ser Thr Tyr Tyr Ala Asp Ser Val Lys Gly
 20 25 30

<210> SEQ ID NO 39
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 39

Asp Ser Ala Ser Phe Asp Tyr
 1 5

<210> SEQ ID NO 40
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 40

Thr Ala Ser Ser Leu Gln Ser
 1 5

-continued

<210> SEQ ID NO 41
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 41

Gln Gln Ser Ala Ala Ser Pro Ser Thr
 1 5

<210> SEQ ID NO 42
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 42

agctatccca tgagc 15

<210> SEQ ID NO 43
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 43

gcgattcagc atactggtgc gccgacaact tacgcagact ccgtgaaggg c 51

<210> SEQ ID NO 44
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 44

gcgtttccgc cgtttgacta c 21

<210> SEQ ID NO 45
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 45

cgggcaagtc agagcattag cagctattta aat 33

<210> SEQ ID NO 46
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 46

tctgcatcct ctttgcaaag t 21

<210> SEQ ID NO 47
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

-continued

<220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (22)..(25)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

 <400> SEQUENCE: 47

 caacagcggg agactgggcc tnnnng 26

 <210> SEQ ID NO 48
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

 <400> SEQUENCE: 48

 tctattcagc ctgagggtag gcgacagcg tacgtagact ccgtgaaggg c 51

 <210> SEQ ID NO 49
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

 <400> SEQUENCE: 49

 ccgccggaga ggtttgacta c 21

 <210> SEQ ID NO 50
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

 <400> SEQUENCE: 50

 gctgcatcca gtttgcaaag t 21

 <210> SEQ ID NO 51
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

 <400> SEQUENCE: 51

 caacagagtt acagtacccc taatagc 27

 <210> SEQ ID NO 52
 <211> LENGTH: 90
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

 <400> SEQUENCE: 52

 aatattagta gtgcagggaa ggggctggag tgggtctcaa gtattgatga ttctggtgct 60

 tctacatatt acgcagactc cgtgaagggc 90

 <210> SEQ ID NO 53
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

-continued

<400> SEQUENCE: 53

gattctgctt cttttgacta c

21

<210> SEQ ID NO 54

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 54

actgcatcca gtttgcaaag t

21

<210> SEQ ID NO 55

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 55

caacagtctg ctgctagtcc ttctacg

27

<210> SEQ ID NO 56

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 56

Ser Ile Gly Gln Lys Gly Gly Gly Thr Gln Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> SEQ ID NO 57

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 57

His Phe Glu Asn Phe Asp Tyr
1 5

<210> SEQ ID NO 58

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 58

Ala Ala Ser His Leu Gln Ser
1 5

<210> SEQ ID NO 59

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 59

-continued

Gln Gln Thr Arg Arg Pro Pro Ser Thr
1 5

<210> SEQ ID NO 60
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 60

Tyr Ile Ala Ser Gly Gly Asp Thr Thr Asn Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> SEQ ID NO 61
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 61

Gly Ala Ser Ala Phe Asp Tyr
1 5

<210> SEQ ID NO 62
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 62

Ala Ala Ser Tyr Leu Gln Ser
1 5

<210> SEQ ID NO 63
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 63

Gln Gln Ser Ser Asn Asp Pro Tyr Thr
1 5

<210> SEQ ID NO 64
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 64

Ser Tyr Gly Met His
1 5

<210> SEQ ID NO 65
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 65

-continued

Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> SEQ ID NO 66
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 66

Ile Asn Ala Lys
 1

<210> SEQ ID NO 67
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 67

Gln Gln Ser Tyr Ser Thr Pro
 1 5

<210> SEQ ID NO 68
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 68

tcgattggtc agaagggtgg tggtagacag tacgcagact ccgtgaaggg c 51

<210> SEQ ID NO 69
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 69

cattttgaga attttgacta c 21

<210> SEQ ID NO 70
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 70

gctgcatccc atttgcaaag t 21

<210> SEQ ID NO 71
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 71

caacagacgc gtaggccgcc ttctacg 27

-continued

<210> SEQ ID NO 72
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 72

tatattgcta gtggtggtga tactacaaat tacgcagact ccgtgaaggg c 51

<210> SEQ ID NO 73
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 73

ggtgcttctg cttttgacta c 21

<210> SEQ ID NO 74
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 74

gctgcatcct atttgcaaag t 21

<210> SEQ ID NO 75
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 75

caacagagtt ctaatgatcc ttatacg 27

<210> SEQ ID NO 76
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 76

agctatggca tgcac 15

<210> SEQ ID NO 77
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 77

gttatatcat atgatggaag taataaatac tatgcagact ccgtgaaggg c 51

<210> SEQ ID NO 78
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 78

attaatgcga ag

12

<210> SEQ ID NO 79

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 79

caacagagtt acagtacccc gacg

24

What is claimed is:

1. A direct current insulating gradient dielectrophoresis (DC-iGDEP) device for separating a target protein species in a biological sample based on a chemical or physical parameter, the device comprising: a sawtooth microfluidic channel configured to separate a target protein species from a biological sample, the channel having an inlet port and an outlet port, the sawtooth microfluidic channel including a plurality of pairs of opposing teeth lining both sides of the microfluidic channel, wherein the teeth have a triangular, asymmetric, crescent, starred or rounded shape, each of the plurality of pairs of opposing teeth including a gap between each of the plurality of pairs of opposing teeth, each of the gaps associated with each of the pairs of opposing teeth decreasing from about [1 mm] 50 μm to about [0.5 mm] 1 μm along the sawtooth microfluidic channel.

2. The device of claim 1, wherein the teeth have a triangular shape.

3. The device of claim 2, wherein the teeth have an equilateral triangular shape having a base and a height.

4. The device of claim 3, wherein the smallest teeth have a base length of 2-10 μm and a height of 2-10 μm , and wherein after every 1-10 repeats successive teeth are 25-50 μm larger in their side length and width.

5. The device of claim 1, wherein the spacing of the initial gaps has a distance of about 945 pm and the spacing of the final gaps has a distance of about 27 pm.]

6. The device of claim 1, wherein the [spacing of the initial gaps has a distance of] *gaps associated with each of the pairs of opposing teeth decrease from about 50 [pm and the spacing of the final gaps has a distance of] μm to about 1 nm.*

7. The device of claim 1, wherein the spacing of the initial gaps has a distance of about 50 pm and the spacing of the final gaps has a distance of about 1 pm.]

8. The device of claim 1, wherein the microfluidic channel has a depth of about 10 to 20 μm .

9. A method of separating from a biological sample a target species based on a chemical or physical parameter comprising,

(a) providing the device of claim 1,

(b) loading a loading volume of the sample into the inlet port,

(c) applying a field to the device to separate particles or molecules in the sample based on the chemical or physical parameter, and

(d) recovering the target species.

10. The method of claim 9, wherein the chemical or physical parameter is charge, size, permittivity, deformation, or shape.

11. The method of claim 9, wherein the target species is an A β aggregate.

12. The method of claim 11, further comprising contacting the recovered A β aggregates with an antibody to confirm the size of the A β aggregate.

13. The method of claim 12, wherein the antibody is specific for oligomeric A β aggregates.

14. The method of claim 12, wherein the antibody is a nanobody, and the nanobody is a C6, A4, E1, D5, 10H, 6E, D10 or BSEC1 nanobody.

15. The method of claim 9, wherein the biological sample has a volume of less than 100 microliters.

16. The method of claim 9, wherein the biological sample is brain tissue, serum, cerebrospinal fluid (CSF), urine or saliva.

17. The method of claim 9, wherein the field is applied for a period of time that is less than 20 minutes.

18. The method of claim 9, wherein the target species is concentrated by several orders of magnitude as compared to the loading volume.

19. The method of claim 9, wherein the target species is concentrated by 10^6 as compared to the loading volume.

20. The method of claim 9, wherein the target species is a protein.

21. The method of claim 20, wherein the protein is p53, islet amyloid polypeptide, beta-amyloid, tau, alpha-synuclein, huntingtin, or superoxide dismutase.

* * * * *